

Antipsychotic Medications: Metabolic and Cardiovascular Risk

John W. Newcomer, M.D.

Individuals with serious mental illness experience excess morbidity and mortality, including an increased prevalence of diabetes mellitus and cardiovascular disease. Cardiovascular disease is the leading cause of death in persons with serious mental illness, and the elevated prevalence of obesity in this population is of particular concern. Obesity is an independent cardiometabolic risk factor that impacts morbidity and mortality and contributes to the development of other cardiometabolic risk factors, such as dyslipidemia and hypertension. In addition, obesity is a major risk factor for type 2 diabetes, with the relative risk of diabetes increasing with body mass index. Increased abdominal fat is strongly associated with insulin resistance, which can lead to impaired glucose regulation. Abdominal obesity, hyperglycemia, hypertension, and dyslipidemia are key components of the metabolic syndrome, a constellation of cardiometabolic risk factors linked by their common association with insulin resistance. Evidence from large clinical samples indicates a high prevalence of metabolic syndrome and all of its components in persons with serious mental illness, particularly in patients with schizophrenia. In addition, psychotropic agents, including some antipsychotic medications, are associated with substantial weight gain, as well as with adiposity-dependent and possibly adiposity-independent changes in insulin sensitivity and lipid metabolism, which increase the risk of diabetes and cardiovascular disease. Among the second-generation antipsychotics, clozapine and olanzapine are associated with the highest risk of substantial weight gain, similar to the weight gain potential associated with low-potency first-generation antipsychotics such as thioridazine or chlorpromazine, as well as with an increased risk of diabetes and dyslipidemia. Various strategies for monitoring cardiometabolic risk factors in patients with mental illness are discussed in this review.

(J Clin Psychiatry 2007;68[suppl 4]:8-13)

Compared to persons without psychiatric disorders, individuals with serious mental illness experience increased morbidity and mortality, especially due to cardiovascular illness.¹⁻³ In a meta-analysis¹ of 152 mortality studies, patients with schizophrenia (N = 35,815) demonstrated a 2-fold greater risk of premature death from medical causes compared to the general population. Recent U.S. public sector data confirm that cardiovascular disease is the leading cause of death in persons with major mental illness.⁴ Further, a 2000 study² from Sweden that assessed mortality after a first hospitalization for schizophrenia indicated that death from cardiovascular causes increased dramatically during the period from 1976 to 1995. This increase was greatest from 1991 to 1995, with male and female patients having an 8.3-fold and 5-fold increase in cardiovascular mortality, respectively. Among those with serious mental illness, however, increased mortality is not limited to patients with schizophrenia. In a study³ of 15,386 individuals with bipolar disorder, the standardized mortality ratio (observed deaths in study population/

expected deaths in general population) for all deaths from medical causes was increased by a factor of 1.9 among men and 2.1 among women.

Recognition of the increased cardiometabolic risk associated with serious mental illness has prompted interest in the factors that contribute to this problem. Key risk factors for cardiovascular disease that have been established by large-scale population-based studies⁵ include obesity, dyslipidemia, hypertension, smoking, and hyperglycemia. Table 1 lists the prevalence of these key modifiable risk factors in patients with schizophrenia and bipolar disorder.⁶⁻¹⁴ While poverty, urbanization, and reduced access to medical care continue to be concerns for persons with serious mental illness, there is growing interest in the contribution of psychotropic treatment to the prevalence of such risk factors as obesity, hyperglycemia, and dyslipidemia.¹⁵ The purpose of this review is to evaluate the clinical importance of cardiometabolic risk factors among persons with mental disorders, address the contribution of antipsychotic medications to increased cardiometabolic risk, and suggest monitoring strategies for modifiable risk factors relevant to the treatment of serious mental illness.

THE PROBLEM OF OBESITY

As defined by World Health Organization criteria,¹⁶ a body mass index (BMI) between 18.5 and 25 kg/m² is considered "normal," while a BMI from 25 to 30 kg/m² is "overweight" and a BMI \geq 30 kg/m² is "obese," although thresholds can vary with ethnicity (e.g., Asians). Obesity has become a major clinical focus, based on the association of overweight and obesity with other cardiovascular risk factors^{17,18} and because it is an independent risk factor for both morbidity and mortality. The U.S. Department of Health and Human Services¹⁹ recently reported that following tobacco use, obesity is the second leading cause of preventable death in the United States. Even in patients who are not obese, increasing adiposity is associated with an increasing risk of significant medical morbidity, producing significant risk beginning in the range of moderate-to-severe overweight related to the

From the Department of Psychiatry, Center for Clinical Studies, Washington University School of Medicine, St. Louis, Mo.

This article was derived from the editorial planning roundtable "Strategies to Integrate Physical Health Care Into Mental Health," which was held August 28, 2006, in Philadelphia, Pa., and supported by an educational grant from Bristol-Myers Squibb Co., Princeton, N.J., and Otsuka America Pharmaceutical, Inc., Rockville, Md.

Dr. Newcomer has served as a consultant for AstraZeneca, Bristol-Myers Squibb, GlaxoSmithKline, Janssen, Solway, Pfizer, and Wyeth and as a consultant for litigation; has received grant/research support from Bristol-Myers Squibb, Janssen, Pfizer, the National Alliance for Research on Schizophrenia and Depression, the National Institute of Mental Health, and the Sidney R. Baer Jr. Foundation; has served on the Data Safety Monitoring Committee for Organon; and has received product development royalties (for a metabolic screening form) from Compact Clinicals.

Corresponding author and reprints: John W. Newcomer, M.D., Department of Psychiatry, Washington University School of Medicine, 600 South Euclid Ave., Campus Box 8134, St. Louis, MO 63110-1093 (e-mail: newcomerj@wustl.edu).

Table 1. Cardiometabolic Disease Risk Factors in Schizophrenia and Bipolar Disease^a

Modifiable Risk Factor	Schizophrenia		Bipolar Disease, Estimated Prevalence of Risk Factor (%)
	Estimated Prevalence of Risk Factor (%)	Relative Risk ^b	
Obesity	45–55	1.5–2 ×	26
Smoking	50–80	2–3 ×	55
Diabetes	10–14	2 ×	10
Hypertension	≥ 18		15
Dyslipidemia		Up to 5 ×	

^aData from references 6–14.^bCompared with the general population.

increased risk of diabetes mellitus, coronary heart disease (CHD), and other conditions.¹⁷

In addition to the association between obesity and overall morbidity and mortality, there is a strong relationship between obesity and the risk of type 2 diabetes, with the relative risk of diabetes increasing sharply with increasing BMI in comparison to the adiposity-related increase in risk associated with other common conditions (e.g., hypertension, CHD, and cholelithiasis).⁵ For instance, in women, a dramatic rise in the relative risk of diabetes occurs as BMI increases from the “overweight” (25 to 30 kg/m²) to the “obese” range (≥ 30 kg/m²).¹⁸ The prevalence of both obesity and diabetes has been rising dramatically in the United States, increasing among U.S. adults by 74% and 61%, respectively, from 1991 to 2001.²⁰

DIABETES, INSULIN RESISTANCE, AND OBESITY

In patients at risk for type 2 diabetes, increased insulin resistance and inefficient glucose utilization work together to exhaust β-cell insulin reserves, resulting in the hyperglycemia of diabetes.²¹ In patients with type 2 diabetes, the risk of a cardiovascular event is equivalent to that in individuals with established CHD.²² This risk equivalence impacts the need for monitoring and treatment, with the National Cholesterol Education Program Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel [ATP] III) guidelines²³ indicating that in persons with diabetes, risk factors such as blood pressure and plasma lipids should be addressed as aggressively as they are in persons who have already suffered a prior myocardial infarction.

Underlying the association between obesity and diabetes described earlier is substantial evidence that increased abdominal fat, in particular, visceral adipose tissue volume, is related strongly to reductions in insulin sensitivity (i.e., insulin resistance). Insulin sensitivity has been shown to decrease as the amount of visceral adipose tissue increases.²⁴ In addition, the relationship of visceral adiposity to diabetes risk extends to the elderly.²⁵ In a study²⁵ of 2964 elderly subjects (mean age: 73.6 years), intermuscular fat and visceral abdominal fat were significantly greater in subjects with type 2 diabetes and impaired glucose tolerance as compared to fat measurements in those with normal glucose tolerance. The importance of adiposity was underscored by the finding that greater intermuscular fat and visceral abdominal fat were associated with higher fasting insulin levels (indicating insulin resistance) in both subjects with a normal BMI (between 18.5 and 25 kg/m²) and obese subjects, suggesting that regional adipose tissue distribution is a key determinant of insulin resistance and altered glucose homeostasis. Therefore, even elderly people with

a “normal” BMI may be at cardiometabolic risk if they have increased visceral abdominal and/or intramuscular fat.

With respect to recommended clinical assessments, the ATP III criteria define abdominal obesity as a waist circumference > 102 cm (> 40.2 in) in men and > 88 cm (> 34.6 in) in women.²⁶ However, either an increased waist circumference or an elevated BMI can serve as a predictor of insulin resistance.²⁷

THE METABOLIC SYNDROME

Abdominal obesity and increased fasting plasma glucose levels constitute 2 components of the metabolic syndrome, a constellation of cardiovascular risk factors that includes metabolic abnormalities associated with insulin resistance and/or compensatory hyperinsulinemia.²⁸ Among patients with metabolic syndrome, the relative risk for diabetes and CHD ranges from 1.5 to 5 times that of the general population.²⁹ In particular, patients' CHD risk increases with the number of metabolic syndrome risk factors present.^{30,31}

Metabolic syndrome is diagnosed when ≥ 3 of the following abnormalities are present: (1) increased waist circumference (> 40 in for men and > 35 in for women), (2) elevated triglyceride levels (≥ 150 mg/dL), (3) low high-density lipoprotein cholesterol (< 40 mg/dL in men and < 50 mg/dL in women), (4) blood pressure ≥ 130/85 mm Hg, and (5) fasting glucose ≥ 110 mg/dL,³² with recent modification of the fasting glucose criteria to ≥ 100 mg/dL.³³ Increases in adiposity,^{34,35} reductions in fitness,³⁶ and genetic factors^{37–42} may all contribute to insulin resistance and hyperinsulinemia. Hyperinsulinemia, in turn, has been implicated in the development of all of the following: abnormal glucose and uric acid metabolism; hemodynamic abnormalities, such as hypertension, increased sympathetic nervous system activity, and sodium retention; atherogenic dyslipidemia; and prothrombotic and proinflammatory states.⁴³

Metabolic syndrome and its components are prevalent among persons with mental illness. Analyzing data from the National Health and Nutrition Examination Survey (NHANES) III (N = 17,689), Allison and colleagues⁴⁴ reported that patients with schizophrenia, especially women, had a significantly higher BMI than did women without schizophrenia (BMI: 27.36 vs. 24.50 kg/m², *p* < .001). Highlighting the increase in risk of metabolic syndrome associated with increases in BMI, Park and colleagues⁴⁵ observed that among men who participated in NHANES III, metabolic syndrome was present in 4.6%, 22.4%, and 59.6% of normal weight, overweight, and obese individuals, respectively, with a similar distribution in women. Compared to age-matched individuals in the general population, a significantly higher prevalence of metabolic syndrome was observed in persons with schizophrenia in the National Institute of Mental Health–sponsored Clinical Antipsychotic Trials of Intervention Effectiveness (CATIE) schizophrenia trial,⁴⁶ in which metabolic syndrome was estimated to be present in 51.6% of women and 36.0% of men.

In general, individuals are more likely to develop metabolic syndrome if they have the following risk factors: older age (≥ 65 years), postmenopausal status, Mexican-American ethnicity, higher BMI, current smoking, low household income, high carbohydrate intake, no alcohol consumption, and physical inactivity.⁴⁵ Mental illness is associated with an increased prevalence of smoking, reduced income, and overweight and obesity, as well as medication effects that include sedation and weight gain.

ANTIPSYCHOTICS AND OBESITY

In the long term, weight gain is a potential problem for many patients receiving antipsychotic therapy. "Clinically significant" weight gain is commonly defined as a $\geq 7\%$ increase in body weight from baseline. Using this definition, the pooled incidence of clinically significant weight gain reported in the U.S. product labels for second-generation antipsychotics (SGAs) is as follows: olanzapine 29%,⁴⁷ quetiapine 23%,⁴⁸ risperidone 18%,⁴⁹ ziprasidone 10%,⁵⁰ and aripiprazole 8%.⁵¹

Consistent with the manufacturers' percentages reported above, long-term data (up to 52 weeks) from several clinical trials⁵²⁻⁵⁴ have suggested that olanzapine has the highest propensity to produce weight gain among the SGAs. This effect may be somewhat dose-related if one includes very low doses outside the antipsychotic dose range, with 1-year mean weight increases of approximately 6 kg for pooled doses of olanzapine from 2.5 to 17.5 mg/day⁵³ compared to a 1-year mean of 12 kg for commonly used antipsychotic doses (12.5 to 17.5 mg/day).⁵⁴ With olanzapine treatment, as with other agents, higher levels of weight gain are associated with lower starting weights at treatment initiation.

Pooled weight gains associated with the other SGAs over 52 weeks in other clinical trials conducted during their developmental phase have also been documented. Quetiapine has been reported⁵⁵ to cause a mean weight increase of approximately 3.6 kg at 52 weeks; however, it should be noted that this result was documented from an "observed case" analysis in contrast to "last observation carried forward" analyses used for the other medications discussed. Risperidone treatment has been associated with a mean weight increase of 2.2 kg at 52 weeks, whereas aripiprazole and ziprasidone treatment each were associated with a mean weight increase of approximately 1 kg over 52 weeks.⁵⁶ In a randomized double-blind trial⁵⁷ comparing aripiprazole and olanzapine over 52 weeks, the effect of aripiprazole on weight gain was significantly less than for olanzapine at all time points ($p < .001$). In an effort to rank SGAs according to their relative propensity to cause weight gain, Tandon and Halbreich⁵⁸ tentatively expressed the order as follows: clozapine > olanzapine > risperidone = quetiapine > ziprasidone = aripiprazole. Weight gain is not only a problem associated with second-generation antipsychotic medicines. Increased risk of weight gain has long been associated with treatment using low-potency first-generation antipsychotic medications like chlorpromazine, thioridazine, and mesoridazine.⁴⁴ High-potency first-generation medications like haloperidol and fluphenazine are associated with minimal mean increases in weight during treatment.⁴⁴

Evidence regarding the mechanism(s) underlying antipsychotic-induced weight gain includes studies⁵⁹ of the relationship between weight gain and drug interactions with various neurotransmitter receptors. The level of H_1 antagonism associated with different antipsychotic medications is hypothesized to modulate feeding behavior (increased appetite and decreased sensation of satiety), based on the significant association of weight gain and the binding affinity for this receptor. Antipsychotics with minimal affinity for H_1 receptors, such as aripiprazole, ziprasidone, and haloperidol, are associated with limited weight gain, while antipsychotics with a high affinity for H_1 receptors, such as clozapine, olanzapine, thioridazine, and chlorpromazine, are associated with clinically significant increases in weight. Based on the differential association of various medications with mechanisms that

support weight gain, it can be hypothesized that a switch from a weight gain-inducing antipsychotic (such as olanzapine) to an antipsychotic with a limited risk for weight gain (such as aripiprazole and ziprasidone) would result in reductions in mean weight.^{60,61} This effect has been observed in several studies, including a 2003 report⁶² in which a switch to aripiprazole from certain SGAs produced a mean weight loss of 1.3 kg to 1.7 kg over a period of 8 weeks. An extended discussion of the effect of antipsychotic "switching" on weight and plasma lipids is presented by Weiden⁶³ elsewhere in this supplement.

ANTIPSYCHOTIC EFFECT ON DIABETES RISK

An increased prevalence of impaired glucose tolerance and diabetes mellitus has been observed in persons with schizophrenia. Type 2 diabetes has been reported⁹ to affect 10% to 14% of this patient population. In addition, impaired glucose tolerance has been documented⁶⁴ in 15% of a small sample of drug-naïve, first-episode patients as compared to 0% in matched control subjects, although cohort-specific effects and hospitalization- and acute illness-related increases in plasma cortisol may have contributed to this finding. Dixon and colleagues⁹ examined Medicare and Medicaid data from the pre-SGA era and reported elevated prevalence rates of diabetes, ranging from 5.1% to 14.9% in patients with schizophrenia compared to nonschizophrenia patients.

Overall, published findings from case reports, chart reviews, database analyses, and clinical trials demonstrate differing metabolic effects with the various first-generation antipsychotic (FGA) and SGA medications.⁵⁶ Evidence is strongest for clozapine and olanzapine, with findings from published reports, including controlled experimental studies and randomized clinical trials, indicating that olanzapine and clozapine are associated with an increased risk of diabetes. Weight gain is a drug-induced adverse event that occurs to a differing degree among SGAs, with clozapine and olanzapine commonly associated with clinically significant weight gain. There is a reasonable hypothesis that much of the effect on glucose regulation observed with the different SGAs can be explained as a function of their effect on weight and adiposity,⁶⁵ although drug effects independent of adiposity have been reported and may add to the effects of adiposity.⁶⁶ Evidence of the effects of risperidone treatment on the risk of diabetes is less extensive than for clozapine and olanzapine, with similar numbers of database analyses, but fewer chart reviews and observational reports and fewer controlled studies. The published data do not support the suggestion that risperidone treatment is associated with a reliable increase in diabetes risk.⁵⁶ Even fewer data are available concerning the relationship between quetiapine treatment and diabetes risk.⁵⁶ Studies with ziprasidone and aripiprazole do not support the hypothesis that these agents adversely affect blood glucose regulation in treated patients.⁵⁶

To test the hypothesis that some antipsychotic medications may have adiposity-independent effects on glucose metabolism, Newcomer and colleagues⁶⁶ used modified oral glucose tolerance tests in groups of antipsychotic-treated but adiposity-matched patients with schizophrenia. These investigators⁶⁶ demonstrated that patients treated with clozapine or olanzapine had significant glucose elevations at all time points compared with patients receiving FGAs and untreated healthy control subjects ($p < .05$). Similarly, Henderson's group⁶⁷ used a frequently sampled

intravenous glucose tolerance test to analyze the effects of SGAs on glucose metabolism in patients with schizophrenia, again matching treated patient groups for adiposity. They reported⁶⁷ significant reductions in insulin sensitivity in patients treated with clozapine ($p < .001$) and olanzapine ($p = .001$) compared to those treated with risperidone.

ANTIPSYCHOTIC EFFECT ON DYSLIPIDEMIA

Dyslipidemia, another modifiable cardiovascular risk factor, also may be affected by antipsychotic therapy.⁶⁸ Koro and colleagues¹⁴ reported the results from a large population-based study that found that olanzapine, but not risperidone, was associated with a nearly 5-fold increase in the risk of developing hyperlipidemia compared with no antipsychotic exposure and a more than 3-fold increase compared with FGA treatment. More recently, results from the randomized CATIE schizophrenia trial⁶⁹ and Stroup et al.⁷⁰ demonstrated that the greatest risk for dyslipidemia was associated with olanzapine therapy. A 52-week open-label trial⁷¹ comparing the effects of aripiprazole (not included in CATIE) and olanzapine on mean changes in serum triglycerides found that while baseline values were similar for both aripiprazole and olanzapine treatment groups (137 and 128 mg/dL, respectively), patients treated with olanzapine had higher mean changes in fasting triglycerides compared with those treated with aripiprazole at week 52 (29.87 vs. 0.6 mg/dL) and at endpoint (24.78 vs. 4.91 mg/dL). These reports are representative of the available literature⁵⁶ and consistent with conclusions from the 2004 joint American Diabetes Association/American Psychiatric Association Consensus Development Statement,¹⁵ which found olanzapine and clozapine to be associated with the greatest risk of causing dyslipidemia among the SGAs, whereas aripiprazole and ziprasidone were noted to show no evidence of an effect on serum lipid levels.

MONITORING CARDIOMETABOLIC RISK

Several consensus panels have recommended approaches for monitoring cardiometabolic risk factors in patients receiving antipsychotics. For example, the Expert Consensus Guideline for the Treatment of Schizophrenia⁷² indicates concern regarding the issue of SGA-related weight gain and the risk of type 2 diabetes and cardiovascular disease in patients with mental illness. The Expert Consensus Guideline⁷² concluded that it is important to monitor for all treatment-related complications in patients taking SGAs, with 60% and 56% of the experts in this group stating that obesity and diabetes, respectively, were the most important complications that should be monitored.

Similarly, 4 professional organizations—the American Diabetes Association, the American Psychiatric Association, the American Association of Clinical Endocrinologists, and the North American Association for the Study of Obesity—cosponsored a Joint Consensus Statement¹⁵ based on an examination of the relationship between SGAs and weight gain, dyslipidemia, and the risk of diabetes (Table 2). The Consensus Statement¹⁵ issued a monitoring schedule for the assessment of cardiometabolic risk in patients receiving SGAs, regardless of the patients' initial level of perceived risk (see Sernyak's⁷³ article elsewhere in this supplement). This schedule includes a review of family history and measurement of weight, BMI, waist circumference, blood pressure, fasting plasma glucose level, and fasting lipid profile at or near baseline during initiation of antipsychotic treatment. Once

Table 2. Metabolic Disturbances Associated With Antipsychotic Agents^a

Drug	Weight Gain	Diabetes Risk	Dyslipidemia
Clozapine	+++	+	+
Olanzapine	+++	+	+
Risperidone	++	D	D
Quetiapine	++	D	D
Aripiprazole	+/-	-	-
Ziprasidone	+/-	-	-

^aAdapted with permission from the American Diabetes Association.¹⁵

Abbreviation: D = discrepant results.

Symbols: + = increased effect, - = no effect.

the patient has begun SGA treatment, the monitoring schedule calls for a reassessment of weight at 4, 8, and 12 weeks, and quarterly thereafter. Blood pressure, fasting plasma glucose level, and fasting lipid profile measurements should be repeated at least 12 weeks after SGA initiation. In the absence of abnormalities, blood pressure, plasma glucose, and plasma lipids should be monitored annually, although more frequent monitoring is suggested for patients with a higher baseline cardiometabolic risk and those with rising values during treatment or in whom interventions are being made. The presence of symptoms such as polyuria or polydipsia, indicators of hyperglycemia, or additional symptoms of diabetic ketoacidosis, such as nausea, vomiting, or altered levels of consciousness, require immediate evaluation and treatment. Weight gain $\geq 5\%$ of baseline body weight or the emergence of dyslipidemia or hyperglycemia should prompt intervention, including consideration of a switch to a medication with lower weight gain liability and lower metabolic risk.

A similar monitoring schedule was proposed by the Mount Sinai Conference.⁷⁴ According to this expert group,⁷⁴ all patients receiving an SGA should have BMI and weight assessments at every visit for the first 6 months after starting a new medication. A gain of 1 BMI unit should prompt consideration of a switch to an agent with a lower weight gain liability. Similarly, a baseline glucose level measurement should be collected from all patients before initiating antipsychotic therapy. Subsequently, patients with significant risk factors for diabetes should have fasting glucose or HbA_{1c} levels assessed at 4 months postinitiation and yearly thereafter. In addition, psychiatrists should ensure that lipids are monitored and any abnormalities are treated appropriately.

CONCLUSION

Individuals with schizophrenia have an increased risk for obesity, type 2 diabetes, and other cardiometabolic risk factors. In particular, obesity can have a important negative impact on patients' overall health in relation to changes in insulin sensitivity and the associated risk of hypertension, hyperglycemia, and hyperlipidemia—all additional cardiometabolic risk factors. Based on substantial evidence that some treatments can increase adiposity, alter plasma lipids, and increase the risk of hyperglycemia, clinicians must be alert for potential negative effects on cardiometabolic risk. Clinicians should monitor and appropriately treat cardiometabolic risk in patients with schizophrenia.

Drug names: aripiprazole (Abilify), chlorpromazine (Thorazine and others), clozapine (FazaClo, Clozaril, and others), olanzapine (Zyprexa), quetiapine (Seroquel), risperidone (Risperdal), ziprasidone (Geodon).

Disclosure of off-label usage: The author has determined that, to the best of his knowledge, no investigational information about pharmaceutical agents that is outside U.S. Food and Drug Administration–approved labeling has been presented in this article.

REFERENCES

- Harris EC, Barraclough B. Excess mortality of mental disorder. *Br J Psychiatry* 1998;173:11–53
- Osby U, Correia N, Brandt L, et al. Mortality and causes of death in schizophrenia in Stockholm County, Sweden. *Schizophr Res* 2000;45:21–28
- Osby U, Brandt L, Correia N, et al. Excess mortality in bipolar and unipolar disorder in Sweden. *Arch Gen Psychiatry* 2001;58:844–850
- Colton CW, Manderscheid RW. Congruencies in increased mortality rates, years of potential life lost, and causes of death among public mental health clients in eight states. *Prev Chronic Dis* 2006;3:A42
- Wilson PW, D'Agostino RB, Levy D, et al. Prediction of coronary heart disease using risk factor categories. *Circulation* 1998;97:1837–1847
- Allison DB, Fontaine KR, Heo M, et al. The distribution of body mass index among individuals with and without schizophrenia. *J Clin Psychiatry* 1999;60:215–220
- Allebeck P. Schizophrenia: a life-shortening disease. *Schizophr Bull* 1989;15:81–99
- Davidson S, Judd F, Jolley D, et al. Cardiovascular risk factors for people with mental illness. *Aust N Z J Psychiatry* 2001;35:196–202
- Dixon L, Weiden P, Delahanty J, et al. Prevalence and correlates of diabetes in national schizophrenia samples. *Schizophr Bull* 2000;26:903–912
- Herran A, de Santiago A, Sandoya M, et al. Determinants of smoking behaviour in outpatients with schizophrenia. *Schizophr Res* 2000;41:373–381
- McElroy SL, Frye MA, Suppes T, et al. Correlates of overweight and obesity in 644 patients with bipolar disorder. *J Clin Psychiatry* 2002;63:207–213
- Ucok A, Polat A, Bozkurt O, et al. Cigarette smoking among patients with schizophrenia and bipolar disorders. *Psychiatry Clin Neurosci* 2004;58:434–437
- Cassidy F, Ahearn E, Carroll BJ. Elevated frequency of diabetes mellitus in hospitalized manic-depressive patients. *Am J Psychiatry* 1999;156:1417–1420
- Koro CE, Fedder DO, L'Italien GJ, et al. An assessment of the independent effects of olanzapine and risperidone exposure on the risk of hyperlipidemia in schizophrenic patients. *Arch Gen Psychiatry* 2002;59:1021–1026
- American Diabetes Association, American Psychiatric Association, American Association of Clinical Endocrinologists, North American Association for the Study of Obesity. Consensus Development Conference on antipsychotic drugs and obesity and diabetes. *Diabetes Care* 2004;27:596–601
- World Health Organization. Body mass index (BMI). Available at: www.euro.who.int/nutrition/20030507_1. Accessed March 20, 2004
- Willett WC, Dietz WH, Colditz GA. Guidelines for healthy weight. *N Engl J Med* 1999;341:427–434
- Colditz GA, Willett WC, Stampfer MJ, et al. Weight as a risk factor for clinical diabetes in women. *Am J Epidemiol* 1990;132:501–513
- Gregg EW, Cheng YJ, Cadwell BL, et al. Secular trends in cardiovascular disease risk factors according to body mass index in US adults. *JAMA* 2005;293:1868–1874
- Mokdad AH, Ford ES, Bowman BA, et al. Prevalence of obesity, diabetes, and obesity-related health risk factors, 2001. *JAMA* 2003;289:76–79
- Bergental RM, Nag SS, Reusch JE, et al. Macrovascular risk factors in patients with diabetes: physician treatment strategies and extent of control. *Endocr Pract* 2005;11:172–179
- Haffner SM, Lehto S, Ronnemaa T, et al. Mortality from coronary heart disease in subjects with type 2 diabetes and in nondiabetic subjects with and without prior myocardial infarction. *N Engl J Med* 1998;339:229–234
- National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation* 2002;106:3143–3421
- Banerji MA, Lebowitz J, Chaiken RL, et al. Relationship of visceral adipose tissue and glucose disposal is independent of sex in black NIDDM subjects. *Am J Physiol* 1997;273:E425–E432
- Goodpaster BH, Krishnaswami S, Resnick H, et al. Association between regional adipose tissue distribution and both type 2 diabetes and impaired glucose tolerance in elderly men and women. *Diabetes Care* 2003;26:372–379
- Morantz C, Torrey B. Practice guideline briefs: definition of metabolic syndrome. *Am Fam Physician* 2005;71:189
- Farin HM, Abbasi F, Reaven GM. Body mass index and waist circumference correlate to the same degree with insulin-mediated glucose uptake. *Metabolism* 2005;54:1323–1328
- Chen NG, Holmes M, Reaven GM. Relationship between insulin resistance, soluble adhesion molecules, and mononuclear cell binding in healthy volunteers. *J Clin Endocrinol Metab* 1999;84:3485–3489
- American Heart Association, National Heart, Lung, and Blood Institute. AHA and NHLBI Scientific Statement. Metabolic syndrome: new guidance for prevention and treatment. Sept 13, 2005. Available at: <http://www.americanheart.org>. Accessed Nov 6, 2006
- Sattar N, Gaw A, Scherbakova O, et al. Metabolic syndrome with and without C-reactive protein as a predictor of coronary heart disease and diabetes in the West of Scotland Coronary Prevention Study. *Circulation* 2003;108:414–419
- Ridker PM, Buring JE, Cook NR, et al. C-reactive protein, the metabolic syndrome, and risk of incident cardiovascular events: an 8-year follow-up of 14,719 initially healthy American women. *Circulation* 2003;107:391–397
- Pasternak R. Adult Treatment Panel II versus Adult Treatment Panel III: what has changed and why? *Am J Cardiol* 2002;89:3C–7C
- Grundey SM, Brewer HB, Cleeman JI, et al. Definition of metabolic syndrome: report of the National Heart, Lung, and Blood Institute/American Heart Association conference on scientific issues related to definition. *Circulation* 2004;109:433–438
- Karter AJ, Mayer-Davis EJ, Selby JV, et al. Insulin sensitivity and abdominal obesity in African-American, Hispanic, and non-Hispanic white men and women: the Insulin Resistance and Atherosclerosis Study. *Diabetes* 1996;45:1547–1555
- Macor C, Ruggeri A, Mazzone P, et al. Visceral adipose tissue impairs insulin secretion and insulin sensitivity but not energy expenditure in obesity. *Metabolism* 1997;46:123–129
- Reaven G. Age and glucose intolerance: effect of fitness and fatness. *Diabetes Care* 2003;26:539–540
- Valle T, Tuomilehto J, Bergman RN, et al. Mapping genes for NIDDM: design of the Finland-United States Investigation of NIDDM Genetics (FUSION) Study. *Diabetes Care* 1998;21:949–958
- Lebovitz HE. Type 2 diabetes: an overview. *Clin Chem* 1999;45:1339–1345
- Watanabe RM, Valle T, Hauser ER, et al. Familiality of quantitative metabolic traits in Finnish families with non-insulin-dependent diabetes mellitus: Finland-United States Investigation of NIDDM Genetics (FUSION) Study Investigators. *Hum Hered* 1999;49:159–168
- Elbein SC, Wegner K, Kahn SE. Reduced beta-cell compensation to the insulin resistance associated with obesity in members of Caucasian familial type 2 diabetic kindreds. *Diabetes Care* 2000;23:221–227
- Baroni MG, Arca M, Sentinelli F, et al. The G972R variant of the insulin receptor substrate-1 (IRS-1) gene, body fat distribution and insulin-resistance. *Diabetologia* 2001;44:367–372
- McIntyre EA, Walker M. Genetics of type 2 diabetes and insulin resistance: knowledge from human studies. *Clin Endocrinol (Oxf)* 2002;57:303–311
- Devaraj S, Rosenson RS, Jialal I. Metabolic syndrome: an appraisal of the pro-inflammatory and procoagulant status. *Endocrinol Metab Clin North Am* 2004;33:431–453
- Allison DB, Mentore JL, Heo M, et al. Antipsychotic-induced weight gain: a comprehensive research synthesis. *Am J Psychiatry* 1999;156:1686–1696
- Park YW, Zhu S, Palaniappan L, et al. The metabolic syndrome: prevalence and associated risk factor findings in the US population from the Third National Health and Nutrition Examination Survey, 1988–1994. *Arch Intern Med* 2003;163:427–436
- McEvoy JP, Meyer JM, Goff DC, et al. Prevalence of the metabolic syndrome in patients with schizophrenia: baseline results from the Clinical Antipsychotic Trials of Intervention Effectiveness (CATIE) schizophrenia trial and comparison with national estimates from NHANES III. *Schizophr Res* 2005;80:19–32
- Zyprexa [package insert]. Indianapolis, Ind: Eli Lilly and Company; 2004
- Seroquel [package insert]. Wilmington, Del: AstraZeneca Pharmaceuticals LP; 2006
- Risperdal [package insert]. Titusville, NJ: Janssen LP; 2006
- Geodon [package insert]. New York, NY: Pfizer Inc; 2005
- Abilify [package insert]. Princeton, NJ: Bristol-Myers Squibb Company and Rockville, Md: Otsuka America Pharmaceutical, Inc; 2006
- Jones B, Basson BR, Walker DJ, et al. Weight change and atypical antipsychotic treatment in patients with schizophrenia. *J Clin Psychiatry* 2001;62(suppl 2):41–44

53. Kinon BJ. The routine use of atypical antipsychotic agents: maintenance treatment. *J Clin Psychiatry* 1998;59(suppl 19):18–22
54. Nemeroff CB. Dosing the antipsychotic medication olanzapine. *J Clin Psychiatry* 1997;58(suppl 10):45–49
55. Brecher M, Zukin S, Leong R, et al. Long-term weight change with quetiapine treatment in schizophrenia: a comprehensive data review. *Neuropsychopharmacology* 2004;29(suppl 1):S109
56. Newcomer JW. Second-generation (atypical) antipsychotics and metabolic effects: a comprehensive literature review. *CNS Drugs* 2005;19:1–93
57. McQuade RD, Stock E, Marcus R, et al. A comparison of weight change during treatment with olanzapine or aripiprazole: results from a randomized, double-blind study. *J Clin Psychiatry* 2004;65(suppl 18):47–56
58. Tandon R, Halbreich U. The second generation “atypical” antipsychotics: similar improved efficacy but different neuroendocrine side effects. *Psychoneuroendocrinology* 2003;28(suppl 1):1–7
59. Kroeze WK, Hufeisen SJ, Popadak BA, et al. H-1 histamine receptor affinity predicts short-term weight gain for typical and atypical antipsychotic drugs. *Neuropsychopharmacology* 2003;28:519–526
60. Chue P. The assessment and management of antipsychotic-associated metabolic disturbances from a psychiatric perspective. *Can J Psychiatry* 2004;49:200–207
61. Nemeroff CB, Lieberman JA, Weiden PJ, et al. From clinical research to clinical practice: a 4-year review of ziprasidone. *CNS Spectr* 2005;10:S1–S20
62. Casey DE. Switching patients to aripiprazole from other antipsychotic agents: a multicenter randomized study. *Psychopharmacology (Berl)* 2003;166:391–399
63. Weiden PJ. Switching antipsychotics as a treatment strategy for antipsychotic-induced weight gain and dyslipidemia. *J Clin Psychiatry* 2007;68(suppl 4):34–39
64. Ryan MC, Collins P, Thakore JH. Impaired fasting glucose tolerance in first-episode, drug-naïve patients with schizophrenia. *Am J Psychiatry* 2003;160:284–289
65. Haupt DW, Fahnestock PA, Flavin KA, et al. Adiposity and insulin sensitivity derived from intravenous glucose tolerance tests in antipsychotic-treated patients. *Neuropsychopharmacology* 2007 March 21 (epub ahead of print)
66. Newcomer JW, Haupt DW, Fucetola R, et al. Abnormalities in glucose regulation during antipsychotic treatment of schizophrenia. *Arch Gen Psychiatry* 2002;59:337–345
67. Henderson DC, Cagliero E, Copeland PM, et al. Glucose metabolism in patients with schizophrenia treated with atypical antipsychotic agents: a frequently sampled intravenous glucose tolerance test and minimal model analysis. *Arch Gen Psychiatry* 2005;62:19–28
68. Newcomer JW. Insulin resistance and metabolic risk [poster 37B]. Presented at the 158th annual meeting of the American Psychiatric Association; May 21–26, 2005; Atlanta, Ga
69. Lieberman JA, Stroup TS, McEvoy JP, et al. Effectiveness of antipsychotic drugs in patients with chronic schizophrenia. *N Engl J Med* 2005;353:1209–1223
70. Stroup TS, Lieberman JA, McEvoy JP, et al. Effectiveness of olanzapine, quetiapine, risperidone, and ziprasidone in patients with chronic schizophrenia following discontinuation of a previous atypical antipsychotic. *Am J Psychiatry* 2006;163:611–622
71. Stock E, Nyilas M, McQuade R, et al. Aripiprazole versus olanzapine in schizophrenia: a 52-week, open-label extension study [poster NR267]. Presented at the 158th annual meeting of the American Psychiatric Association; May 21–26, 2005; Atlanta, Ga
72. Kane JM, Leucht S, Carpenter D, et al. Expert Consensus Guideline Series: Optimizing Pharmacologic Treatment of Psychotic Disorders. Introduction: methods, commentary, and summary. *J Clin Psychiatry* 2003;64(suppl 12):5–19
73. Sernyak MJ. Implementation of monitoring and management guidelines for second-generation antipsychotics. *J Clin Psychiatry* 2007;68(suppl 4):14–18
74. Marder SR, Essock SM, Miller AL, et al. Physical health monitoring of patients with schizophrenia. *Am J Psychiatry* 2004;161:1334–1349