

Neurochemistry in the Pathophysiology of Autism

Christopher J. McDougle, M.D.; Craig A. Erickson, M.D.;
Kimberly A. Stigler, M.D.; and David J. Posey, M.D.

Significant progress has been made in the search for underlying pathophysiologic mechanisms in autism over the past 50 years. The cause of the disorder, however, remains largely unknown. This article reviews neurochemical contributions to the pathophysiology of autism with a focus on monoamines, glutamate/ γ -aminobutyric acid systems, and neuropeptides. As these efforts move forward, it will be important to begin to integrate genetic studies with those involving neuroimaging and post-mortem research in each of these 3 areas, as well as with pharmacologic treatment approaches.

(J Clin Psychiatry 2005;66[suppl 10]:9-18)

Research into the pathophysiology and etiology of autistic disorder (autism) has been ongoing for nearly a half century. Despite these significant efforts, the cause remains unknown. This review will discuss neurochemical aspects of the pathophysiology of autism. Three primary areas will be highlighted, including monoamines (serotonin [5-hydroxytryptamine, 5-HT], dopamine [DA], norepinephrine [NE]), glutamate/ γ -aminobutyric acid (GABA) systems, and neuropeptides. Where data are available, peripheral and central neurochemistry, genetics, neuroimaging, and postmortem findings will be presented.

MONOAMINES

Serotonin

Serotonin neurons are widely distributed throughout the mammalian brain. This neuronal system is one of the earliest to develop, and the turnover rate of 5-HT is higher in the immature mammalian brain than at any other time in life. Serotonin plays a critical role as a growth factor in the immature brain, directing both proliferation and maturation.¹

Initial studies on the pathophysiology of autism focused on the 5-HT system. A recent chapter provided a detailed review of the results from those investigations, including peripheral and central neurochemistry, behavioral/neuroendocrine challenges, genetics, and neuroimaging.² We will provide a brief summary of those findings.

Schain and Freedman³ were the first to study whole blood serotonin (WBS) in autism. Their sample included 3 groups: mildly retarded children, autistic children who were severely retarded, and severely retarded children without autism. Consistent unusual elevations of WBS were found only in the autistic children, although the mean WBS level of the other severely retarded group was higher than that of the mildly retarded group. No differences were found in presenting clinical symptoms between the 6 autistic children with the highest WBS levels and those who had normal levels. These results were largely replicated by Ritvo and colleagues.⁴ In 1987, Anderson and others from Yale published results from their laboratory and reviewed and summarized data on WBS levels in autism to that date.⁵ WBS concentrations were significantly higher in drug-free autistic subjects than in normal controls, with 38% of the subjects determined to be "hyperserotonemic." Results from a subsequent study by McBride et al.⁶ led the investigators to conclude that the prevalence of hy-

From the Department of Psychiatry, Section of Child & Adolescent Psychiatry, Indiana University School of Medicine and the Christian Sarkine Autism Treatment Center, James Whitcomb Riley Hospital for Children, Indianapolis, Ind.

This article is derived from the teleconference series "The Management of Autism and Its Related Disorders," which was held February 10–April 21, 2005, and supported by an educational grant from Janssen Medical Affairs, L.L.C.

This work was supported in part by a Department of Housing and Urban Development grant (B-01-SP-IN-0200, Dr. McDougle), a Research Unit on Pediatric Psychopharmacology–Psychosocial Intervention grant (U10 MH066766) from the National Institute of Mental Health (NIMH), Rockville, Md., to Indiana University (Drs. McDougle, Stigler, and Posey), a Young Investigator Award from the National Alliance for Research on Schizophrenia and Depression (Dr. Posey), a Daniel X. Freedman Psychiatric Research Fellowship Award (Dr. Posey), and a Career Development Award (K23 MH068627) from the NIMH (Dr. Posey).

The section on serotonin in this article is an adaptation of an earlier chapter by the authors and has been substantially revised.

Corresponding author and reprints: Christopher J. McDougle, M.D., Department of Psychiatry, Indiana University School of Medicine, 1111 W. 10th Street, PB 2nd Floor, Indianapolis, IN 46202-4800 (e-mail: cmcdougl@iupui.edu).

perserotonemia in autism may have been previously overestimated because of failure to control for race and pubertal status.

In a large study by Leboyer et al.,⁷ WBS levels were determined in 62 subjects with autism aged 3 to 23 years, 91 healthy controls aged 2 to 16 years, and 118 healthy subjects over 16 years of age. Twenty-nine (48%) of the 60 autistic subjects for whom there was a sample met criteria for hyperserotonemia. Among controls, WBS values diminished with age, whereas WBS levels among autistic subjects appeared to be age-independent. In this same study, 51% of mothers, 45% of fathers, and 87% of siblings (older than 16 years) of autistic subjects had hyperserotonemia.

In summarizing results from studies of WBS in autism, many but not all investigations have found elevated WBS levels in younger autistic subjects that tend to remain higher than those of normal controls across the age range. In contrast, most studies of normal subjects have demonstrated an age-related decline in WBS levels with increasing age. Some investigators have suggested that these results could be explained, in part, by abnormal maturational processes of the 5-HT system in autistic subjects.^{5,7} Additional factors that may affect WBS levels include race, pubertal status, and treatment with psychotropic medication. Whether WBS levels will prove to be a useful quantitative measure in the search for genetic susceptibility to autism remains to be determined.

In general, studies of urinary excretion of 5-hydroxyindoleacetic acid (5-HIAA),⁸ the primary metabolite of 5-HT, and whole blood tryptophan concentrations⁵ have not found significant differences between autistic subjects and controls.

Studies of baseline levels of a measure of central 5-HT function, cerebrospinal fluid (CSF) 5-HIAA, have found no difference between children with autism and controls.⁹⁻¹¹ Two studies that utilized probenecid to block the transport of 5-HIAA out of the CSF found similar¹² or slightly lower¹³ levels in autistic children compared with nonautistic children with psychosis.

Behavioral/neuroendocrine challenge studies have been conducted in autistic subjects. The immediate precursor of 5-HT, 5-hydroxytryptophan (5-HTP), was administered to children with autism and adult normal control subjects.^{14,15} Prolactin response to 5-HTP was reduced in the children with autism, suggesting diminished central 5-HT responsiveness. Blunted prolactin release was also found in response to fenfluramine 60 mg given orally, in an investigation of 7 male young adults with autism and matched healthy controls.¹⁶ Utilizing a different strategy, the acute tryptophan depletion (ATD) paradigm was administered to 17 drug-free adults with autism by McDougle and colleagues.¹⁷ The ATD resulted in a significant reduction in plasma free and total tryptophan, whereas administration of sham depletion (containing tryptophan) led to a significant in-

crease in these plasma measures. Eleven of the 17 subjects showed a worsening of symptoms, including a significant increase in whirling, flapping, pacing, banging, hitting self, rocking, and toe walking, with ATD compared to sham depletion. Another set of challenge studies involved the 5-HT_{1D} receptor agonist sumatriptan, which has been shown to increase growth hormone release. Eleven adults with autism or Asperger's disorder and 9 controls were given subcutaneous sumatriptan and placebo separated by 1 week.¹⁸ The research subjects had a significantly greater growth hormone response than controls, suggesting that a hypersensitivity of the 5-HT_{1D} receptor may exist. In a related study, Hollander et al.¹⁹ reported that the severity of repetitive behavior at baseline in these subjects was positively correlated with the growth hormone response to sumatriptan. The same investigators recently found that the oral administration of *m*-chlorophenylpiperazine (*m*-CPP) resulted in a significant increase in repetitive behaviors and prolactin in adults with autism or Asperger's disorder in comparison with controls.²⁰

A number of investigations of genes involved in the 5-HT system have been conducted in autism. The 5-HT transporter (5-HTT), the site of action of serotonin reuptake inhibitors, has been considered a candidate gene for autism. Cook et al.²¹ were the first to report the presence of an association between the short variant of a functional insertion-deletion polymorphism in the promoter region of 5-HTT (HTTLPR) and autism. In contrast, Klauck et al.²² identified preferential transmission of the long variant of HTTLPR in their sample of autistic subjects. A number of subsequent studies involving subjects from various countries have reported similar results or have been unable to replicate either finding.²

Results from studies involving other genes contributing to the 5-HT system, including the genes encoding the 5-HT₇ receptor²³ and the 5-HT_{2A} receptor,²⁴ respectively, have not identified a significant association with autism. Tryptophan 2,3 dioxygenase (TDO2) is the rate-limiting enzyme in the catabolism of tryptophan, the precursor of 5-HT. A study by Nabi et al.²⁵ demonstrated a significant difference in the transmission of TDO2 haplotypes to autistic subjects, suggesting the presence of a susceptibility mutation in the TDO2 or a nearby gene. Recent investigations have sought a relationship between a subset of autistic subjects with prominent rigid-compulsive behaviors and 5-HTT with some preliminary encouraging results.^{26,27}

Neuroimaging studies of the 5-HT system have also been completed in autism. The first investigation utilized positron emission tomography (PET) to assess the tracer α -[¹¹C]methyl-L-tryptophan (AMT) as an indicator of 5-HT synthesis in 8 autistic children and 5 of their siblings.²⁸ Gross asymmetries of 5-HT synthesis in frontal cortex, thalamus, and cerebellum were found in all 7 of the autistic boys but not in the only female autistic subject. Such asymmetries were not identified in the frontal cortex

or thalamus of the siblings, although 1 sibling showed increased [^{11}C]AMT accumulation in the right dentate gyrus. This boy had a history of calendar calculation, and he ritualistically lined up his toys. The investigators concluded that the focal abnormalities in [^{11}C]AMT accumulation may represent either aberrant innervation by 5-HT terminals or altered function in anatomically normal pathways. A subsequent study by the same investigators,²⁹ again using PET and [^{11}C]AMT, found that for nonautistic children, 5-HT synthesis capacity was more than 200% of adult values until the age of 5 years and then declined toward adult values. In autistic children, 5-HT synthesis capacity increased gradually between the ages of 2 years and 15 years to values 1.5 times adult normal values. It was concluded that humans undergo a period of high brain 5-HT synthesis capacity during childhood, and that this developmental process is disrupted in autistic children.

Dopamine

The monoamine DA is integral to motor and cognitive functioning, as well as hormone release.³⁰ A role for DA in autism has been postulated, in part, based upon the beneficial effects observed with the use of DA D₂ receptor antagonists in treating this population. This class of drugs has been shown to effectively target symptoms commonly exhibited by individuals with autism, such as aggression, self-injurious behavior, and hyperactivity.³¹

To a large extent, neurochemical research in this area has centered on the measurement of the major DA metabolite, homovanillic acid (HVA), in urine and plasma, as well as CSF. When considering this research, it is important to understand that only approximately 25% of urine and plasma HVA appears to result from central DA turnover, and that peripheral measures are primarily able to identify only significant alterations in central DA metabolism.³²

In a study of catecholamine metabolism in 22 youths with autism aged 5 to 16 years and controls matched for age and sex, no significant difference in urinary DA was found between groups.³³ Minderaa and colleagues³⁴ investigated plasma HVA and prolactin levels, as well as urine HVA and DA excretion, in medicated and unmedicated autistic subjects and unmedicated controls. The authors found no significant differences between the autistic and control groups, suggesting normal peripheral indices of DA functioning. Martineau et al.³⁵ measured urine levels of DA and its derivatives, including HVA, 3-methoxytyramine (3MT), and NE + epinephrine (EPI), in 156 children with autism, compared with age-matched mentally retarded and normal controls. The levels were found to decrease significantly with age in all 3 groups. Significantly decreased levels of DA and HVA were found in medicated versus unmedicated autistic youth. The authors hypothesized that the results may be secondary to a defect in maturation of monoaminergic systems in autism.

Several studies that measured CSF HVA levels have been published. Gillberg and Svennerholm¹⁰ reported that group mean levels of CSF HVA were elevated by approximately 50% in autistic children, compared to an age- and sex-matched control group with neurologic disorders. However, similar to previous findings by Cohen and colleagues,^{12,13} a controlled study of CSF monoamine effects with fenfluramine treatment in 9 youths with autism reported normal levels of CSF HVA.³⁶ In addition, Narayan and colleagues¹¹ also found normal levels of CSF HVA in their controlled study of 17 children with autism. The authors reported that the results were consistent with the majority of earlier studies that did not find a group difference in this metabolite in CSF.

Some genetic studies of DA involvement in autism have been completed. Comings and colleagues³⁷ suggested that the A1 allele of the DA D₂ receptor gene may be associated with a number of behavioral disorders in which it may act as a modifying gene. In their case-control study, the authors examined a variety of neuropsychiatric disorders, including autism, which are believed to involve defects in DA neurotransmission. The prevalence of the A1 allele was noted to be significantly increased in the group with autism.

Another study examined the DA D₁ and D₅ receptor genes in autism via restriction endonuclease fingerprinting.³⁸ One novel missense change (L88F) occurred in transmembrane domain II at a highly conserved amino acid in all DA receptors, as well as in α_1 - and β -adrenergic receptors. The mutation was identified in a Caucasian male patient with autism.

Robinson and colleagues³⁹ examined the DA- β -hydroxylase (D β H) gene as a candidate locus in 37 families with 2 or more children with pervasive developmental disorders (PDDs) using the affected sib-pair method. D β H is an enzyme that catalyzes DA to NE. There was no increased concordance for D β H alleles in affected siblings, but the mothers had a higher frequency of alleles containing a 19-base pair deletion. The authors hypothesized that lowered maternal serum D β H activity may produce a suboptimal uterine environment, which, in combination with a genetic susceptibility, could result in PDDs in some families.

Dopaminergic activity has also been investigated via neuroimaging techniques in autism. Using the PET tracer [^{18}F]fluorodopa (FDOPA), Ernst and colleagues⁴⁰ studied 14 children with autism (8 males; age, 10–17 years) and 10 controls (7 males; age, 12–17 years). In the autistic group, regional FDOPA accumulation in the anterior medial prefrontal cortex was significantly reduced by 39%.

In another study employing PET, 6 children aged 3 to 5 years with autism were treated with 6R-L-erythro-5,6,7,8-tetrahydrobiopterin (R-BH₄), a cofactor for tyrosine hydroxylase in the biosynthetic pathway of catecholamines.⁴¹ Study subjects were included only if the investigators

noted a relatively low level of R-BH₄ in the CSF. Prior to treatment, PET revealed increased DA D₂ receptor binding in the caudate and putamen as a whole. After treatment, a 10% decrease in DA D₂ receptor binding was observed. In addition, CSF levels of R-BH₄ were found to be significantly increased.

Norepinephrine

The neurotransmitter NE is associated with arousal, memory, anxiety, and autonomic activity.³⁰ Produced from DA, NE is metabolized to vanillylmandelic acid (VMA) in the periphery, and to 3-methoxy-4-hydroxyphenylglycol (MHPG) in the central nervous system. Plasma and urine levels of NE and its metabolites have been considered to be generally well correlated with central functioning.⁴² However, research has also shown that estimates of the proportion of MHPG in blood and urine originating in the central nervous system, relative to that from the periphery, have been uncertain, ranging from 10% to 60%.⁴³

Regarding studies of blood measures of NE and its metabolites, Lake et al.⁴⁴ investigated levels of NE and DβH in autistic and normal control subjects. The authors recorded a significantly higher level of blood NE in the group with autism. In contrast to this finding, lower levels of DβH were found in the autistic group, perhaps due to the enzyme's longer half-life.

A study of plasma MHPG in youth with autism and normal controls recorded similar group means of 3.7 ng/mL and 3.2 ng/mL, respectively.⁴⁵ Similarly, Minderaa and colleagues⁴⁶ found no significant difference in mean ± SD plasma MHPG levels in unmedicated autistic (3.1 ± 0.6 ng/mL; N = 17), medicated autistic (3.3 ± 1.0 ng/mL; N = 23), and normal control (3.2 ± 1.2 ng/mL; N = 20) groups. In addition, no significant mean differences in levels of MHPG, NE, and EPI were recorded between subjects with autism and normal controls when evening and overnight urines were examined. The authors suggested that notable abnormalities in basal NE measures did not appear to be present in autism.

As a whole, studies of CSF MHPG in autism have found no significant differences compared to controls. Young and colleagues⁴⁷ reported a mean CSF MHPG concentration of 9 ng/mL in subjects with autism, a level comparable to that recorded in normal subjects. Another larger study of 25 youths with autism and age- and sex-matched controls also found similar values between groups.¹⁰

GLUTAMATE AND GABA

Glutamate, the primary excitatory amino acid neurotransmitter, is found in high concentrations throughout the brain. It is thought to be crucial in neuronal plasticity and higher cognitive functioning.⁴⁸ Glutamate receptors are divided into metabotropic and ionotropic types. The ionotropic receptors are further classified into the follow-

ing 3 families: N-methyl-D-aspartate (NMDA), α-amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA), and kainate. Several researchers have postulated that glutamate dysfunction may play a role in autism.^{49,50}

GABA, another amino acid neurotransmitter, is the primary inhibitory neurotransmitter in the brain. It is synthesized from glutamate by glutamic acid decarboxylase (GAD). Investigators have also hypothesized that GABA may have an important role in the pathophysiology of autism.⁵¹

Despite the important roles that these neurotransmitters may play, there has been a relative paucity of literature directly examining glutamatergic or GABA function in autism. This section will review neurochemical, genetic, and postmortem studies concerning the function of these neurotransmitters in autism.

Several reports have suggested that peripheral levels of glutamate are elevated in the plasma of subjects with autism and other PDDs. Aldred et al.⁵² collected blood from 23 subjects aged 4 to 29 years with autism or Asperger's disorder and 55 of their family members (32 parents, 23 siblings) and measured amino acid concentrations. They found that concentrations of glutamate, phenylalanine, lysine, and asparagine were significantly higher in both subjects and family members compared to age-matched controls. Glutamine levels were significantly lower. Moreno-Fuenmayor et al.⁵³ also measured amino acid levels in 14 children with autism (all under 10 years) and age- and sex-matched controls. They found that aspartate was higher and glutamine and asparagine were lower in subjects than in controls. However, another analysis is in disagreement with these findings. Rolf et al.⁵⁴ measured amino acid content and GABA in platelet-rich plasma. They found that aspartate and glutamate were decreased in 18 drug-free children aged 8 to 14 years with autism compared to 14 age-matched healthy controls. GABA and glutamine levels were also significantly lower. In contrast to this, Dhossche et al.⁵¹ reported elevated plasma GABA levels (measured by gas chromatography/mass spectrometry) in a small, heterogeneous sample of 9 subjects aged 5 to 15 years with autism compared to 9 control subjects with attention-deficit/hyperactivity disorder (ADHD). Most of the autistic subjects were taking prescribed psychotropic or anticonvulsant drugs, and all of the ADHD controls were taking psychostimulants. In summary, studies reporting on peripheral amino acid levels in autism present mixed results. Interpretation of these results is also difficult given the small sample sizes, possible medication effects, and different methodologies used.

A number of genetic studies of the glutamate and GABA systems have been conducted in autism. Jamain et al.⁵⁵ showed that the glutamate receptor ionotropic kainate 2 (GRIK2) or glutamate receptor 6 (GluR6) gene is in disequilibrium with autism, and that an excess of maternal transmission of the GRIK2 haplotype exists. Interestingly,

maternal transmission disequilibrium for GRIK2 has also been found by the same group in schizophrenia.⁵⁶ More importantly, this finding was recently replicated by Shuang et al.⁵⁷ in 174 Chinese Han parent-offspring trios. GRIK2 is located in the chromosome 6q21 region, which has been identified as a potential autism susceptibility region by at least 1 genome-wide scan study.⁵⁸

GAD1 encodes glutamic acid decarboxylase 67kDa protein (GAD67), an enzyme important in the conversion of glutamate to GABA. As a decarboxylase, it requires vitamin B₆ as a cofactor, which some believe may have efficacy in autism.⁵⁹ It also occurs on chromosome 2q, which shows evidence for linkage in several genome-wide scans. Rabionet et al.⁶⁰ recently performed association studies on several candidate genes in this region including GAD1. They found no evidence for significant association between these genes and autism.

Ramoz et al.⁶¹ did find linkage for 2 single nucleotide polymorphisms (SNPs) on another chromosome 2q gene, SLC25A12. This gene encodes for the mitochondrial aspartate/glutamate carrier. However, the report by Rabionet et al.⁶⁰ referenced their own unpublished data, which failed to replicate this finding in their sample.

Serajee et al.⁶² found suggestive evidence for linkage disequilibrium between autism and the metabotropic glutamate receptor 8 (GRM8) gene, which occurs on chromosome 7q, another region implicated in genome-wide scans.

Several lines of evidence implicate the 15q11-q13 chromosome region as potentially harboring autism susceptibility genes. This includes numerous reports suggesting that duplications and other abnormalities in this region may occur in as many as 3% of autistic individuals.⁶³ This genetic region has also been implicated in Prader-Willi and Angelman syndromes, which share clinical features with autism. Finally, this region contains several GABA type A receptor subunit genes, which are candidates as autism susceptibility genes.

Cook et al.⁶⁴ tested several loci in this region for linkage disequilibrium and were the first to report an association between a marker (155CA-2) within the GABA receptor subunit β -3 gene (GABRB3) and autism in a sample of 140 trios. Linkage disequilibrium for this marker has been found in one other sample,⁶⁵ but not others.⁶⁶⁻⁶⁹ However, Martin and colleagues⁶⁸ did report linkage disequilibrium with another nearby marker (GABRB3) in this same region.

Menold et al.⁷⁰ examined 16 SNPs located within GABRB3, GABRA5, and GABRG3 for linkage disequilibrium using the Pedigree Disequilibrium Test. Two SNPs located within the GABRG3 gene were in disequilibrium, suggesting that the GABRG3 gene or a nearby gene may contribute to genetic risk for autism. McCauley et al.⁷¹ performed linkage disequilibrium mapping across a region containing a cluster of GABA receptor subunit genes on chromosome 15q12. Six markers individually,

across GABRB3 and GABRA5, and several haplotypes inclusive of those markers, demonstrated nominally significant association.

In summary, there is emerging evidence that the GRIK2 gene may be involved in autism. There is conflicting evidence as to the role of other genes that encode GABA receptor subunits and the mitochondrial aspartate/glutamate carrier. Findings regarding GAD and other glutamate receptors await replication.

The glutamate and GABA systems have also been evaluated in postmortem studies in autism. Purcell et al.⁷² used complementary DNA (cDNA) microarray technology, additional measurements of messenger RNA (mRNA) and protein levels, as well as receptor autoradiography to study the cerebellum and hippocampus in a total of 10 persons with autism and 23 matched controls. They found several genes to be up-regulated in autism, most notably the excitatory amino acid transporter 1 and the glutamate receptor AMPA 1 (GluR1) genes. In addition, higher levels of the corresponding proteins were found by Western blotting. Finally, AMPA receptor density was decreased in both the granule cell layer and molecular cell layer of the cerebellum. Other notable findings were no significant differences in GAD 1/2 protein levels (by Western blotting) or NMDA receptor density (by autoradiography) in the cerebellum.

Both GAD65 and GAD67 (2 isoforms of GAD) levels were measured in postmortem cerebellar (N = 5) and parietal cortices (N = 5) of persons with autism compared to controls.⁷³ In this study, GAD65 (but not GAD67) was significantly lower in cerebellar cortices and GAD67 (but not GAD65) was significantly reduced in parietal cortices. In an autoradiographic study, Blatt et al.⁷⁴ reported decreased density of GABA receptors in hippocampal sections of brain in cases with autism (N = 4) compared to controls (N = 3). The density of 6 other receptors, 5-HT_{1A}, 5-HT₂, cholinergic M₁, high affinity choline uptake site, NMDA, and kainate, did not differ significantly.

In summary, postmortem studies suggest that genes and proteins involved with glutamate and GABA functioning may be abnormal in autism. However, definitive conclusions are difficult to make given the limited number of studies and small sample sizes.

NEUROPEPTIDES

Oxytocin and Vasopressin

The 9 amino acid peptides oxytocin (OT) and vasopressin (AVP) have been implicated in the social behavior of mammals.⁷⁵ These neuropeptides are synthesized in the hypothalamus and secreted from the posterior pituitary gland, exist solely in mammals, and differ at only 2 amino acids.⁷⁶ Receptors for these peptides have been found throughout the limbic system, in the forebrain, and in brain stem autonomic centers.⁷⁷

Animal models of OT and AVP involvement in regulating behavior have laid the groundwork for hypotheses that these peptides may be involved in the pathophysiology associated with PDDs.⁷⁸⁻⁸⁰ OT knockout mice have been found to have impaired social memory in the presence of intact olfactory and general cognitive abilities.⁸¹ OT has been found to promote pair bonding,⁷⁵ and OT antagonist infusion into the nucleus accumbens has inhibited the formation of partner preference in female prairie voles.⁸² In rats, AVP has been shown to facilitate social memory,⁷⁶ and in male prairie voles, AVP has been found to promote pair bonding.⁷⁵

The possible role of OT in the etiology of the social impairment of PDDs has been evaluated at epidemiologic, neurochemical, and therapeutic levels. The hypothesis that human neonatal exposure to OT (pitocin) during labor induction may lead to long-term OT receptor down regulation has been evaluated in 2 studies comparing affected children and controls.⁸³ A retrospective review of the birth histories of 41 children with autism compared with the records from 25 age- and IQ-matched controls found no increased incidence of pitocin exposure in the children with PDDs.⁸⁴ A comparison of the birth records of 633 preschool children with language disorders, autism, or generalized low IQ found similar rates of labor induction among all groups evaluated.⁸⁵

Plasma, but not central, OT levels have been evaluated in a cohort of 30 autistic male children.^{86,87} This investigation compared the OT levels in patients with those of age-matched but not IQ-matched control children. The first report on this cohort noted a decreased level of plasma OT in the autistic children.⁸⁶ The second report using the same cohort again reported lower OT levels in the children with autism but also increased levels of OT extended form, a precursor molecule with a C-terminal 3 amino acid extension.⁸⁷ The authors hypothesized that a deficit in an unspecified prohormone convertase may be responsible for the low OT levels found in this sample of autistic children. The finding of abnormal plasma OT levels in male autistic children is interesting in light of the animal model of OT impacting social attachment, which found a predominant female predisposition to OT susceptibility.⁷⁵

A single study has evaluated the intravenous administration of OT in subjects with PDDs. Using a repeated infusion model with each subject as his own control, 6 adults with autism and 9 adults with Asperger's disorder were monitored for 6 different types of repetitive behavior in 60-minute intervals following infusion of OT or placebo for up to 4 hours.⁸⁸ This investigation reported no significant main effect for drug and a significant reduction in combined repetitive behaviors over time, but no significant difference for any single repetitive behavior was reported. No difference in response was noted between the subjects with autism or Asperger's disorder, and no social measures were obtained during the challenge paradigm.

While OT has been evaluated on several levels in autistic individuals, AVP has been primarily investigated at the genetic level looking for polymorphisms in the gene coding for arginine vasopressin receptor 1A (AVPR1A).^{89,90} One analysis looking at 125 independent autistic probands and 65 autism-affected sibling pair families concluded that differences at the amino acid level in the AVPR1A gene are not likely to confer genetic vulnerability to autism, but that there may be some significant transmission disequilibrium outside of the gene itself in the 5' flanking region.⁸⁹ An analysis of 115 autism trios genotyped 2 polymorphisms from the 5' flanking region of the AVPR1A gene and found nonsignificant transmission patterns.⁹⁰ Interestingly, both studies found different 5' flanking region polymorphisms. This difference was hypothesized to represent varying genetic backgrounds in the 2 samples, or spurious false positive findings.⁹⁰

Investigations of the social impairment associated with PDDs have not yet found significant epidemiologic, neurochemical, therapeutic, or genetic evidence to support a primarily OT- and/or AVP-mediated etiology. Future work may need to focus on central levels of OT and AVP, and/or possibly focus more on how the sexually dimorphic effects of OT and AVP seen in lower mammals may translate into differential effects between genders in individuals with PDDs.

Opioids

Several observations of autistic children led to the early evaluation of opioid dysregulation as a possible etiologic explanation of PDDs. These have included elevated pain threshold, little interest in social interactions, and episodes of motor hyperactivity.⁹¹ These findings appeared to match those seen in infant animals administered opiates.⁹² In one way, the human opioid system relates to the hypothalamic-pituitary-adrenal (HPA) axis and other neuropeptides because β -endorphin is excreted at the same time as corticotropin (ACTH) from the anterior pituitary. This occurs because these peptide hormones are initially part of the same preprohormone.⁹³

Evaluations of β -endorphin (and β -endorphin metabolites in some cases) levels in the serum, CSF, and urine of patients with PDDs compared to controls have yielded conflicting results (see Tordjman et al.,⁹⁴ for review). Ten studies enrolling a total of 142 patients with PDDs have evaluated serum β -endorphin levels, and the results represent a relatively equal mix of increased, decreased, or similar β -endorphin levels found in the patients compared to control subjects. In 2 investigations of CSF β -endorphin levels in patients with PDDs, 1 reported increased and 1 reported decreased levels in affected patients.

Investigation of urinary opioid peptides has attempted to look at whether inadequate processing of exogenous opioids by the gastrointestinal tract may result in over-absorption and finally urinary excretion of the peptides.

Such an evaluation analyzed the urine of 10 children with autism compared to 11 adult controls and found no difference between groups.⁹⁵ This study also specifically evaluated dipeptidyl-peptidase in the serum of patients and found no difference compared with controls. This enzyme is present at the intestinal brush border, and is expected to be involved in the cleaving of exogenous dietary opioids. These findings have been challenged by those who have presented earlier data pointing toward abnormal urinary opioid levels in patients with PDDs.⁹⁶ Disagreement over the preparation of samples and the limits of detection methods emerged among the different groups reporting contradictory results with regard to urinary opioids in patients with PDDs.

The use of naltrexone, an opioid receptor antagonist, has been evaluated in several open-label and placebo-controlled trials in patients with PDDs having unknown, elevated, or normal levels of serum β -endorphin.⁹⁷ Again, as in the studies looking at serum and CSF β -endorphin levels, conflicting results on the efficacy of naltrexone exist, but most controlled studies suggest that the core symptoms of autism and associated maladaptive behavior are not significantly affected by naltrexone.

Overall, the importance of endogenous or exogenous opioid systems in patients with PDDs is subject to much debate driven by contradictory evidence. This uncertainty has led to the hypothesis that it may not be opioid levels per se that may contribute to symptoms commonly seen in PDDs, but the manner in which the endogenous opioid system interacts with other neurotransmitter systems such as the 5-HT, DA, NE, glutamate, or GABA systems.⁹¹ To date, these potential multisystem interactions have not been extensively explored. A final manner in which opioid activity may impact theories of the pathophysiology of PDDs could be derived from the knowledge that β -endorphin has been found to inhibit oxytocin activity in rats.⁹⁸ It is clear that more research is needed to better understand how the endogenous opioid system interacts both with traditional neurotransmitters and with other neuropeptides with respect to the pathobiology and treatment of PDDs.

Cortisol/ACTH

Levels of the anterior pituitary hormone ACTH and the adrenal product cortisol have been evaluated in patients with PDDs as a means to evaluate the HPA axis in these subjects. As with opioids, conflicting results have been the rule rather than the exception.^{94,99} In patients with PDDs, plasma ACTH and cortisol, and cortisol levels in response to the dexamethasone suppression test, have each been found to be similar, lower, or elevated compared to control subjects in different studies.^{94,99} Recently, investigators reported elevated ACTH and low cortisol levels in 36 autistic individuals, commenting that their results were "difficult to interpret" in light of conflicting data previously

reported.⁹⁹ One report on the circadian rhythm of cortisol, as measured by urinary secretion in 19 autistic patients compared to control subjects, found no significant difference in the daily rhythm of cortisol secretion.¹⁰⁰

While it is clear that a simple excess or deficit of ACTH or cortisol is not present in the majority of patients with PDDs, it is less clear how abnormally elevated or decreased levels can be further interpreted at an individual patient level to ascribe any causal relationship between these peptides and common PDD symptomatology.

Melatonin

It has been hypothesized that hypersecretion of melatonin from the pineal gland may be responsible for a cascade of events impacting the HPA axis, thus possibly leading to an "autistic-like" phenotype.¹⁰¹ This theory is based on evidence (primarily from animal studies) that increased melatonin can lead to decreased corticotropin-releasing hormone (CRH) secretion from the hypothalamus, leading to decreased pituitary ACTH and β -endorphin excretion while at the same time causing, by an unknown mechanism, an elevation in whole brain 5-HT.¹⁰¹ One study to date¹⁰² has systematically evaluated levels of melatonin in patients with PDDs. Ten patients with autism aged 16 to 30 years had melatonin levels drawn over a 24-hour period, and these levels were compared to findings from control subjects. The authors reported no difference in mean daily melatonin concentration, with only a trend toward a lower amplitude of melatonin peak at night in autistic patients. The exact role of melatonin in patients with PDDs remains to be fully characterized and understood. Little, if any, evidence to date exists pointing to a primary melatonin dysfunction.

Secretin

While the role of secretin as a classical hormone in the gastrointestinal system is well known, its role as a neuropeptide is continuing to be defined.¹⁰³ This potential neuroactivity has been further investigated, in part, because of preliminary reports of effective secretin treatment of social and communication impairment in patients with PDDs.¹⁰⁴ While controlled studies have failed to demonstrate the efficacy of secretin in treating autistic symptomatology,^{104,105} recent evidence of the neuroactive properties of secretin has become clear.^{103,106-108} In human and rat samples¹⁰⁸ and rat samples alone,¹⁰⁶ secretin immunoreactivity has been shown in Purkinje cells of the cerebellum,^{106,108} central cerebellar nuclei,¹⁰⁸ pyramidal cells of the motor cortex,¹⁰⁸ primary sensory neurons,^{106,108} and the brainstem.¹⁰⁶ Additionally, in separate work in rats, mRNA coding for the secretin receptor has also been found in cerebellar GABA interneurons.¹⁰⁷ This work postulated that in the cerebellum, secretin may be secreted by Purkinje cells, then act as a retrograde messenger modulating GABA activity. It is clear that secretin likely has neuroac-

tive effects. Its potential contributions to the pathophysiology of PDDs remains poorly understood. Based upon the results of numerous placebo-controlled studies, however, secretin has been determined to be an ineffective treatment for autism.^{104,105}

Thyroid Hormone

Thyroid-stimulating hormone (TSH), released by the pituitary gland, has been evaluated in several patients with autism. In 2 separate analyses, each with 10 affected children compared with a similar number of adult controls, no difference in TSH levels was found.^{102,109}

Other Anterior Pituitary Hormones

Plasma growth hormone,⁹⁹ prolactin,^{99,109} luteinizing hormone,¹⁰⁹ and follicle-stimulating hormone¹⁰⁹ levels have been shown to be no different between autistic individuals and controls in small studies.

Other Neuropeptides and Neurotrophins

A novel analysis of neuropeptides and neurotrophins from frozen blood samples of neonates subsequently diagnosed with a PDD (N = 69), mental retardation without autism (N = 54), cerebral palsy (N = 63), and normal control patients (N = 54) found significantly elevated levels of several measured substances in the PDD and mental retardation groups.¹¹⁰ Concentrations of neonatal vasoactive intestinal peptide (VIP), calcitonin gene-related peptide (CGRP), brain derived neurotrophic factor (BDNF), and neurotrophin 4/5 were significantly higher in both the PDD and mental retardation groups. No significant differences between the PDD and mental retardation groups were noted. Concentrations of substance P, pituitary adenylate cyclase-activating polypeptide (PACAP), nerve growth factor (NGF), and neurotrophin 3 were all similar among all groups tested. In a different study, CSF levels of the neurotrophic factor insulin-like growth factor-I were found to be similar in an analysis of 11 autistic patients and 11 age-matched “disabled” controls.¹¹¹ While the results from the neonatal samples may not be specific to PDD, their significance lies in pointing demonstrably to how neuropeptide/neurotrophin dysregulation early in development may set the brain on a course toward disordered development, including, in some cases, a course toward PDDs.¹¹²

CONCLUSION

This review has explored the available literature on neurochemical contributions to the pathophysiology of autism, with a focus on monoamines (5-HT, DA, NE), glutamate/GABA systems, and neuropeptides. With respect to monoamines, the majority of studies that have focused on basal measures of plasma, urine, and CSF have been negative. The 1 exception is that of elevated WBS or

“hyperserotonemia,” which has been replicated in multiple investigations. Its underlying mechanism, however, remains unclear. Behavioral challenges of monoaminergic systems have primarily involved 5-HT. A number of significant differences have been found between autistic subjects (primarily adults) and controls, although most results have not yet been replicated. Furthermore, results from behavioral challenges are largely based on peripheral responses, with central effects often being inferred. Preliminary PET studies involving 5-HT and DA systems have yielded potentially important findings. Concerns about radiation exposure in youth may limit further studies utilizing currently available technology, although investigations in adults may be possible. Postmortem assessment of monoaminergic involvement needs to be completed. Encouraging results from preliminary genetic studies of the glutamate and GABA systems are emerging. The use of magnetic resonance spectroscopy (MRS) will allow for an indirect assessment of these systems in the living brain. Finally, results from animal studies indicate that the OT and AVP systems are important, if not critical, for affiliative behavior. It will be important to continue to incorporate genetic studies into ongoing neuroimaging and postmortem investigations in each of the 3 areas of neurochemistry discussed in this review.

Drug names: naltrexone (Revia and others), probenecid (Probalan and others), sumatriptan (Imitrex).

Disclosure of off-label usage: The authors have determined that, to the best of their knowledge, naltrexone is not approved by the U.S. Food and Drug Administration for the treatment of aggression in autism; probenecid is not approved for use in cerebrospinal fluid studies in autism; and sumatriptan, fenfluramine, and *m*-chlorophenylpiperazine are not approved for use in behavioral challenge studies in autism.

REFERENCES

- Whitaker-Azmitia PM. The role of serotonin and serotonin receptors in development of the mammalian nervous system. In: Zagon IS, McLaughlin PJ, eds. *Receptors in the Developing Nervous System*, vol 2. Neurotransmitters. London, UK: Chapman & Hall; 1993:43-53
- McDougle CJ, Posey DJ, Potenza MN. Neurobiology of serotonin function in autism. In: Hollander E, ed. *Autism Spectrum Disorders*. New York, NY: Marcel Dekker; 2003:199-220
- Schain RJ, Freedman DX. Studies on 5-hydroxyindole metabolism in autistic and other mentally retarded children. *J Pediatr* 1961;58:315-320
- Ritvo ER, Yuwiler A, Geller E, et al. Increased blood serotonin and platelets in early infantile autism. *Arch Gen Psychiatry* 1970;23:566-572
- Anderson GM, Freedman DX, Cohen DJ, et al. Whole blood serotonin in autistic and normal subjects. *J Child Psychol Psychiatry* 1987;28:885-900
- McBride PA, Anderson GM, Hertzog ME, et al. Effects of diagnosis, race, and puberty on platelet serotonin levels in autism and mental retardation. *J Am Acad Child Adolesc Psychiatry* 1998;37:767-776
- Leboyer M, Philippe A, Bouvard M, et al. Whole blood serotonin and plasma beta-endorphin in autistic probands and their first-degree relatives. *Biol Psychiatry* 1999;45:158-163
- Minderaa RB, Anderson GM, Volkmar FR, et al. Urinary 5-hydroxyindoleacetic acid and whole blood serotonin and tryptophan in autistic and normal subjects. *Biol Psychiatry* 1987;22:933-940
- Gillberg C, Svennerholm L, Hamilton-Hellberg C. Childhood psychosis and monoamine metabolites in spinal fluid. *J Autism Dev Disord* 1983; 13:383-396
- Gillberg C, Svennerholm L. CSF monoamines in autistic syndromes and

- other pervasive developmental disorders of early childhood. *Br J Psychiatry* 1987;151:89–94
11. Narayan M, Srinath S, Anderson GM, et al. Cerebrospinal fluid levels of homovanillic acid and 5-hydroxyindoleacetic acid in autism. *Biol Psychiatry* 1993;33:630–635
 12. Cohen DJ, Shaywitz BA, Johnson WT, et al. Biogenic amines in autistic and atypical children: cerebrospinal fluid measures of homovanillic acid and 5-hydroxyindoleacetic acid. *Arch Gen Psychiatry* 1974;31:845–853
 13. Cohen DJ, Caparulo BK, Shaywitz BA, et al. Dopamine and serotonin metabolism in neuropsychiatrically disturbed children: CSF homovanillic acid and 5-hydroxyindoleacetic acid. *Arch Gen Psychiatry* 1977;34:545–550
 14. Hoshino Y, Watanabe M, Tachibana R, et al. A study of the hypothalamus-pituitary function in autistic children by the loading test of 5HTP, TRH, and LH-RH. *Jpn J Brain Res* 1983;9:94–95
 15. Hoshino Y, Tachibana JR, Watanabe M, et al. Serotonin metabolism and hypothalamic-pituitary function in children with infantile autism and minimal brain dysfunction. *Jpn J Psychiatry Neurol* 1984;26:937–945
 16. McBride PA, Anderson GM, Hertzog ME, et al. Serotonergic responsiveness in male young adults with autistic disorder: results of a pilot study. *Arch Gen Psychiatry* 1989;46:213–221
 17. McDougle CJ, Naylor ST, Cohen DJ, et al. Effects of tryptophan depletion in drug-free adults with autistic disorder. *Arch Gen Psychiatry* 1996;53:993–1000
 18. Novotny S, Hollander E, Allen A, et al. Increased growth hormone response to sumatriptan challenge in adult autistic disorders. *Psychiatry Res* 2000;94:173–177
 19. Hollander E, Novotny S, Allen A, et al. The relationship between repetitive behaviors and growth hormone response to sumatriptan challenge in autistic disorder. *Neuropsychopharmacology* 2000;22:163–167
 20. Novotny S, Hollander E, Phillips A, et al. Increased repetitive behaviours and prolactin responsivity to oral *m*-chlorophenylpiperazine in adults with autism spectrum disorders. *Int J Neuropsychopharmacol* 2004;7:249–254
 21. Cook EH, Courchesne R, Lord C, et al. Evidence of linkage between the serotonin transporter and autistic disorder. *Mol Psychiatry* 1997;2:247–250
 22. Klauck SM, Poustka F, Benner A, et al. Serotonin transporter (5-HTT) gene variants associated with autism? *Hum Mol Genet* 1997;6:2233–2238
 23. Lassig JP, Vachirasomtoon K, Hartzell K, et al. Physical mapping of the serotonin 5-HT7 receptor gene (HTR7) to chromosome 10 and pseudogene (HTR7P) to chromosome 12, and testing of linkage disequilibrium between HTR7 and autistic disorder. *Am J Med Genet* 1999;88:472–475
 24. Veenstra-VanderWeele J, Kim SJ, Lord C, et al. Transmission disequilibrium studies of the serotonin 5-HT2A receptor gene (HTR2A) in autism. *Am J Med Genet* 2002;114:277–283
 25. Nabi R, Serajee FJ, Chugani DC, et al. Association of tryptophan 2,3 dioxigenase gene polymorphism with autism. *Am J Med Genet B Neuropsychiatr Genet* 2004;125:63–68
 26. McCauley JL, Olson LM, Dowd M, et al. Linkage and association analysis at the serotonin transporter (SLC6A4) locus in a rigid-compulsive subset of autism. *Am J Med Genet B Neuropsychiatr Genet* 2004;127:104–112
 27. Mulder EJ, Anderson GM, Kema IP, et al. Serotonin transporter intron 2 polymorphism associated with rigid-compulsive behaviors in Dutch individuals with pervasive developmental disorder. *Am J Med Genet B Neuropsychiatr Genet* 2005;133:93–96
 28. Chugani DC, Muzik O, Rothermel R, et al. Altered serotonin synthesis in the dentatohalamocortical pathway in autistic boys. *Ann Neurol* 1997;42:666–669
 29. Chugani DC, Muzik O, Behen M, et al. Developmental changes in brain serotonin synthesis capacity in autistic and nonautistic children. *Ann Neurol* 1999;45:287–295
 30. Moore RY, Bloom FE. Central catecholamine neuron systems: anatomy and physiology of the norepinephrine and epinephrine systems. *Ann Rev Neurosci* 1979;2:113–168
 31. Anderson GM, Hoshino Y. Neurochemical studies in autism. In: Cohen D, Volkmar F, eds. *Handbook of Autism and Pervasive Developmental Disorders*. 2nd ed. New York, NY: John Wiley; 1997:325–343
 32. Maas JW, Hattox SE, Greene NM, et al. Estimates of dopamine and serotonin synthesis by the awake human brain. *J Neurochem* 1980;34:1547–1549
 33. Launay JM, Bursztejn C, Ferrari P, et al. Catecholamine metabolism in infantile autism: a controlled study of 22 autistic children. *J Autism Dev Disord* 1987;17:333–347
 34. Minderaa RB, Anderson GM, Volkmar FR, et al. Neurochemical study of dopamine functioning in autistic and normal subjects. *J Am Acad Child Adolesc Psychiatry* 1989;28:200–206
 35. Martineau J, Barthelemy C, Jouve J, et al. Monoamines (serotonin and catecholamines) and their derivatives in infantile autism: age-related changes and drug effects. *Dev Med Child Neurol* 1992;34:593–603
 36. Ross DL, Klyklo WM, Anderson GM. Cerebrospinal fluid indoleamine and monoamine effects in fenfluramine treatment of autism. *Ann Neurol* 1985;18:394
 37. Comings DE, Comings BG, Muhleman D, et al. The dopamine D2 receptor locus as a modifying gene in neuropsychiatric disorders. *JAMA* 1991;266:1793–1800
 38. Feng J, Sobell JL, Heston LL, et al. Scanning of the dopamine D1 and D5 receptor genes by REF in neuropsychiatric patients reveals a novel missense change at a highly conserved amino acid. *Am J Med Genet B Neuropsychiatr Genet* 1998;81:172–178
 39. Robinson PD, Schutz CK, Macciardi F, et al. Genetically determined low maternal serum dopamine β -hydroxylase levels and the etiology of autism spectrum disorders. *Am J Med Genet* 2001;100:30–36
 40. Ernst M, Zametkin AJ, Matochik JA, et al. Low medial prefrontal dopaminergic activity in autistic children [letter] [Erratum in *Lancet* 1998;351:454]. *Lancet* 1997;350:638
 41. Fernell E, Watanabe Y, Adolfson I, et al. Possible effects of tetrahydrobiopterin treatment in six children with autism—clinical and positron emission tomography data: a pilot study. *Dev Med Child Neurol* 1997;39:313–318
 42. Roy A, Pickar D, DeJong J, et al. Norepinephrine and its metabolites in cerebrospinal fluid, plasma, and urine: relationship to hypothalamic-pituitary-adrenal axis function in depression. *Arch Gen Psychiatry* 1988;45:849–857
 43. Blomberry PA, Kopin IJ, Gordon EK, et al. Conversion of MHPG to vanillylmandelic acid: implication for the importance of urinary MHPG. *Arch Gen Psychiatry* 1980;37:195–198
 44. Lake R, Zeigler MG, Murphy DL. Increased norepinephrine levels and decreased DBH activity in primary autism. *Arch Gen Psychiatry* 1977;35:553–556
 45. Young JG, Cohen DJ, Hattox SE, et al. Plasma free MHPG and neuroendocrine responses to challenge doses of clonidine in Tourette's Syndrome: preliminary report. *Life Sci* 1981;29:1467–1475
 46. Minderaa RB, Anderson GM, Volkmar FR, et al. Noradrenergic and adrenergic functioning in autism. *Biol Psychiatry* 1994;36:237–241
 47. Young JG, Cohen DJ, Kavanaugh ME, et al. Cerebrospinal fluid, plasma, and urinary MHPG in children. *Life Sci* 1981;28:2837–2845
 48. Cotman CW, Kahle JS, Miller SE, et al. Excitatory amino acid neurotransmission. In: Bloom FE, Kupfer DJ, eds. *Psychopharmacology: The Fourth Generation of Progress*. New York, NY: Raven Press; 1995
 49. Carlsson ML. Hypothesis: is infantile autism a hypoglutamatergic disorder? relevance of glutamate-serotonin interactions for pharmacotherapy. *J Neural Transm* 1998;105:525–535
 50. McDougle CJ. Current and emerging therapeutics of autistic disorder and related pervasive developmental disorders. In: Davis KL, Charney D, Coyle JT, et al, eds. *Psychopharmacology: The Fifth Generation of Progress*. Philadelphia, Pa: Lippincott Williams & Wilkins; 2002
 51. Dhossche D, Applegate H, Abraham A, et al. Elevated plasma γ -aminobutyric acid (GABA) levels in autistic youngsters: stimulus for a GABA hypothesis of autism. *Med Sci Monit* 2002;8:PR1–6
 52. Aldred S, Moore KM, Fitzgerald M, et al. Plasma amino acid levels in children with autism and their families. *J Autism Dev Disord* 2003;33:93–97
 53. Moreno-Fuenmayor H, Borjas L, Arrieta A, et al. Plasma excitatory amino acids in autism. *Invest Clin* 1996;37:113–128
 54. Rolf LH, Haarmann FY, Grotemeyer KH, et al. Serotonin and amino acid content in platelets of autistic children. *Acta Psychiatr Scand* 1993;87:312–316
 55. Jamain S, Betancur C, Quach H, et al. Linkage and association of the glutamate receptor 6 gene with autism. *Mol Psychiatry* 2002;7:302–310
 56. Bah J, Quach H, Ebstein RP, et al. Maternal transmission disequilibrium of the glutamate receptor GRIK2 in schizophrenia. *Neuroreport* 2004;15:1987–1991
 57. Shuang M, Liu J, Jia MX, et al. Family-based association study between autism and glutamate receptor 6 gene in Chinese Han trios. *Am J Med*

- Genet B Neuropsychiatr Genet 2004;131:48–50
58. Philippe A, Martinez M, Guilloud-Bataille M, et al. Genome-wide scan for autism susceptibility genes. Paris Autism Research International Sibpair Study [Erratum in Hum Mol Genet 1999;8:1353]. Hum Mol Genet 1999;8:805–812
 59. Pfeiffer SI, Norton J, Nelson L, et al. Efficacy of vitamin B6 and magnesium in the treatment of autism: a methodology review and summary of outcomes. J Autism Dev Disord 1995;25:481–493
 60. Rabionet R, Jaworski JM, Ashley-Koch AE, et al. Analysis of the autism chromosome 2 linkage region: GAD1 and other candidate genes. Neurosci Lett 2004;372:209–214
 61. Ramoz N, Reichert JG, Smith CJ, et al. Linkage and association of the mitochondrial aspartate/glutamate carrier SLC25A12 gene with autism. Am J Psychiatry 2004;161:662–669
 62. Serajee FJ, Zhong H, Nabi R, et al. The metabotropic glutamate receptor 8 gene at 7q31: partial duplication and possible association with autism. J Med Genet 2003;40:e42
 63. Sutcliffe JS, Nurmi EL, Lombroso PJ. Genetics of childhood disorders: XLVII. Autism, pt 6: duplication and inherited susceptibility of chromosome 15q11-q13 genes in autism. J Am Acad Child Adolesc Psychiatry 2003;42:253–256
 64. Cook EH Jr, Courchesne RY, Cox NJ, et al. Linkage-disequilibrium mapping of autistic disorder, with 15q11-13 markers. Am J Hum Genet 1998;62:1077–1083
 65. Buxbaum JD, Silverman JM, Smith CJ, et al. Association between a GABRB3 polymorphism and autism. Mol Psychiatry 2002;7:311–316
 66. Maestrini E, Lai C, Marlow A, et al. Serotonin transporter (5-HTT) and γ -aminobutyric acid receptor subunit beta 3 (GABRB3) gene polymorphisms are not associated with autism in the IMGSA families. Am J Med Genet 1999;88:492–496
 67. Salmon B, Hallmayer J, Rogers T, et al. Absence of linkage and linkage disequilibrium to chromosome 15q11-q13 markers in 139 multiplex families with autism. Am J Med Genet 1999;88:551–556
 68. Martin ER, Menold MM, Wolpert CM, et al. Analysis of linkage disequilibrium in γ -aminobutyric acid receptor subunit genes in autistic disorder. Am J Med Genet 2000;96:43–48
 69. Nurmi EL, Bradford Y, Chen Y, et al. Linkage disequilibrium at the Angelman syndrome gene UBE3A in autism families. Genomics 2001;77:105–113
 70. Menold MM, Shao Y, Wolpert CM, et al. Association analysis of chromosome 15 GABA_A receptor subunit genes in autistic disorder. J Neurogenet 2001;15:245–259
 71. McCauley JL, Olsen LM, Delahanty R, et al. A linkage disequilibrium map of the 1-Mb 15q12 GABA(A) receptor subunit cluster and association to autism. Am J Med Genet B Neuropsychiatr Genet 2004;131:51–59
 72. Purcell AE, Jeon OH, Zimmerman AW, et al. Postmortem brain abnormalities of the glutamate neurotransmitter system in autism. Neurology 2001;57:1618–1628
 73. Fatemi SH, Halt AR, Stary JM, et al. Glutamic acid decarboxylase 65 and 67 kDa proteins are reduced in the autistic parietal and cerebellar cortices. Biol Psychiatry 2002;52:805–810
 74. Blatt GJ, Fitzgerald CM, Guptill JT, et al. Density and distribution of hippocampal neurotransmitter receptors in autism: an autoradiographic study. J Autism Dev Disord 2001;31:537–543
 75. Insel TR. A neurobiological basis of social attachment. Am J Psychiatry 1997;154:726–735
 76. Insel TR, O'Brien DJ, Leckman JF. Oxytocin, vasopressin, and autism: is there a connection? Biol Psychiatry 1999;45:145–157
 77. Barberis C, Tribollet E. Vasopressin and oxytocin receptors in the central nervous system. Crit Rev Neurobiol 1996;10:119–154
 78. Panksepp J. Commentary on the possible role of oxytocin in autism. J Autism Dev Disord 1993;23:567–568
 79. Modahl C, Fein D, Waterhouse L, et al. Does oxytocin deficiency mediate social deficits in autism? J Autism Dev Disord 1992;22:449–451
 80. Young LJ, Pitkow LJ, Ferguson JN. Neuropeptides and social behavior: animal models relevant to autism. Mol Psychiatry 2002;7:S38–S39
 81. Ferguson JN, Young LJ, Hearn EF, et al. Social amnesia in mice lacking the oxytocin gene. Nat Genet 2000;25:284–288
 82. Young LJ, Lim MM, Gingrich B, et al. Cellular mechanisms of social attachment. Horm Behav 2001;40:133–138
 83. Wahl RU. Could oxytocin administration during labor contribute to autism and related behavioral disorders? a look at the literature. Med Hypotheses 2004;63:456–460
 84. Gale S, Ozonoff S, Lainhart J. Brief report: pitocin induction in autistic and nonautistic individuals. J Autism Dev Disord 2003;33:205–208
 85. Fein D, Allen D, Dunn M, et al. Pitocin induction and autism [letter]. Am J Psychiatry 1997;154:438–439
 86. Modahl C, Green L, Fein D, et al. Plasma oxytocin levels in autistic children. Biol Psychiatry 1998;43:270–277
 87. Green L, Fein D, Modahl C, et al. Oxytocin and autistic disorder: alterations in peptide forms. Biol Psychiatry 2001;50:609–613
 88. Hollander E, Novotny S, Hanratty M, et al. Oxytocin infusion reduces repetitive behaviors in adults with autistic and Asperger's disorders. Neuropsychopharmacology 2003;28:193–198
 89. Kim SJ, Young LJ, Gonen D, et al. Transmission disequilibrium testing of arginine vasopressin receptor 1A (AVPR1A) polymorphisms in autism. Mol Psychiatry 2002;7:503–507
 90. Wassink TH, Piven J, Vieland VJ, et al. Examination of AVPR1A as an autism susceptibility gene. Mol Psychiatry 2004;9:968–972
 91. Sher L. Autistic disorder and the endogenous opioid system. Med Hypotheses 1997;48:413–414
 92. Panksepp J, Siviy SM, Normansell LA. Brain opioids and social emotions. In: Reite M, Field T, eds. The Psychobiology of Attachment and Emotion. New York, NY: Academic Press; 1985:3–49
 93. Guyton AC, Hall JE. Textbook of Medical Physiology. 9th ed. Philadelphia, Pa: WB Saunders Company; 1996
 94. Tordjman S, Anderson GM, McBride PA, et al. Plasma beta-endorphin, adrenocorticotropin hormone, and cortisol in autism. J Child Psychol Psychiatry 1997;38:705–715
 95. Hunter LC, O'Hare A, Herron WJ, et al. Opioid peptides and dipeptidyl-peptidase in autism. Dev Med Child Neurol 2003;45:121–128
 96. Shattock P, Hooper M, Waring R. Opioid peptides and dipeptidyl-peptidase in autism [letter]. Dev Med Child Neurol 2004;46:357
 97. Erickson CA, Stigler KA, Posey DJ, et al. Psychopharmacology. In: Volkmar FR, ed. Autism and Pervasive Developmental Disorders. 2nd ed. Cambridge, UK: Cambridge University Press; 2005. In press
 98. Bicknell RJ, Leng G, Lincoln DW, et al. Nalazone excites oxytocin neurons in the supraoptic nucleus of lactating rats after chronic morphine treatment. J Physiol 1988;396:297–317
 99. Curin JM, Terzic J, Petkovic ZB, et al. Lower cortisol and higher ACTH levels in individuals with autism. J Autism Dev Disord 2003;33:443–448
 100. Richdale AL, Prior MR. Urinary cortisol circadian rhythm in a group of high-functioning children with autism. J Autism Dev Disord 1992;22:433–447
 101. Chamberlain RS, Herman BH. A novel biochemical model linking dysfunctions in brain melatonin, proopiomelanocortin peptides, and serotonin in autism. Biol Psychiatry 1990;28:773–793
 102. Nir I, Meir D, Zilber N, et al. Brief report: circadian melatonin, thyroid-stimulating hormone, prolactin, and cortisol levels in serum of young adults with autism. J Autism Dev Disord 1995;25:641–654
 103. Ng SSM, Yung WH, Chow BKC. Secretin as a neuropeptide. Mol Neurobiol 2002;26:97–107
 104. Esch BE, Carr JE. Secretin as a treatment for autism: a review of the evidence. J Autism Dev Disord 2004;34:543–556
 105. Patel NC, Yeh JY, Shepherd MD, et al. Secretin treatment for autistic disorder: a critical analysis. Pharmacotherapy 2002;22:905–914
 106. Koves K, Kausz M, Reser D, et al. What may be the anatomical basis that secretin can improve the mental functions in autism? Regul Pept 2002;109:167–172
 107. Yung WH, Leung PS, Ng SS, et al. Secretin facilitates GABA transmission in the cerebellum. J Neurosci 2001;21:7063–7068
 108. Koves K, Kausz M, Reser D, et al. Secretin and autism: a basic morphological study about the distribution of secretin in the nervous system. Regul Pept 2004;123:209–215
 109. Hoshino Y, Watanabe M, Tachibana R, et al. The TRH and LH-RH loading test in autistic children. Fukushima J Med Sci 1985;31:55–61
 110. Nelson KB, Grether JK, Croen LA, et al. Neuropeptides and neurotrophins in neonatal blood of children with autism or mental retardation. Ann Neurol 2001;49:597–606
 111. Vanhala R, Turpeinen U, Riikonen R. Low levels of insulin-like growth factor-I in cerebrospinal fluid in children with autism. Dev Med Child Neurol 2001;43:614–616
 112. Minshew N. What are neuropeptides and neurotrophins and why are they important in autism? [editorial] J Autism Dev Disord 2001;31:517