

It is illegal to post this copyrighted PDF on any website.

Functional Impairment in Youth With ADHD: Normative Data and Norm-Referenced Cutoff Points for the Before School Functioning Questionnaire and the Parent Rating of Evening and Morning Behavior Scale, Revised

Stephen V. Faraone, PhD^{a,*}; Norberto J. DeSousa, MA^b; Marina Komolova, PhD^c;
F. Randy Sallee, MD, PhD^d; Bev Inledon, PhD^b; and Timothy E. Wilens, MD^e

ABSTRACT

Objective: Children with attention-deficit/hyperactivity disorder (ADHD) frequently manifest behavioral difficulties in the morning prior to school and in the afternoons and evenings. We sought to establish norms for 2 time-specific measures of functioning: the Before School Functioning Questionnaire (BSFQ) and the Parent Rating of Evening and Morning Behavior Scale, Revised (PREMB-R), which includes Morning (AM) and Evening (PM) subscales.

Methods: The normative online survey of a representative US sample of 1,200 primary caregivers of children and adolescents aged between 6 and 17 years was conducted in June 2016. A quota system was used whereby caregivers of 50 male and 50 female children or adolescents were recruited in each age group, ie, 100 parents for each of the 12 age groups. Diagnosis of ADHD relied on a caregiver's report that his or her child was so diagnosed by a health professional.

Results: Across all items of the BSFQ, youth with current untreated ADHD or a history of ADHD were rated as more severely ill than those without ADHD (all unadjusted *P* values < .001), even after adjustment for psychiatric comorbidity (all adjusted *P* values < .001). A similar pattern was observed for the PREMB-R AM (all unadjusted *P* values < .001; all adjusted *P* values < .001, except for item 1 [*P* = .01]) and PREMB-R PM (all unadjusted *P* values < .001; all adjusted *P* values < .001). The use of a large population sample allowed for computation of age-stratified norms for 4 thresholds of risk: screening risk (80th percentile), mild functional impairment (90th percentile), moderate functional impairment (93rd percentile), and severe functional impairment (98th percentile).

Conclusions: The norms generated by this study can guide clinicians in the use of the BSFQ and PREMB-R for identifying those ADHD youth who may be experiencing difficulties in the early morning and late afternoon/evening. Such tools are needed given the availability of treatments that can target ADHD symptoms and impairments at these extremes of the daily routine.

J Clin Psychiatry 2020;81(1):19m12956

To cite: Faraone SV, DeSousa NJ, Komolova M, et al. Functional impairment in youth with ADHD: normative data and norm-referenced cutoff points for the Before School Functioning Questionnaire and the Parent Rating of Evening and Morning Behavior Scale, Revised. *J Clin Psychiatry*. 2020;81(1):19m12956.

To share: <https://doi.org/10.4088/JCP.19m12956>

© Copyright 2019 Physicians Postgraduate Press, Inc.

^aDepartments of Psychiatry and of Neuroscience and Physiology, SUNY Upstate Medical University, Syracuse, New York

^bIronshore Pharmaceuticals & Development, Inc, Grand Cayman, Cayman Islands

^cHighland Therapeutics Inc, Toronto, Ontario, Canada

^dIronshore Pharmaceuticals Inc, Durham, North Carolina

^ePediatric Psychopharmacology Unit of the Psychiatry Department, Massachusetts General Hospital, and Harvard Medical School, Department of Psychiatry, Boston, Massachusetts

*Corresponding author: Stephen V. Faraone, PhD, SUNY Upstate Medical University, Institute for Human Performance, Room 3707, 505 Irving Ave, Syracuse, NY 13210 (sfaraone@childpsychresearch.org).

The early morning is a source of distress and disability for children with ADHD. Barkley and Cunningham¹ showed that ADHD impaired early morning organization, self-care, preparation for the school day, and transportation to school. Failing to complete their morning routine puts children at risk for tardiness, forgetting to take homework to school, and family conflict. Parents of stimulant-treated ADHD children report stress while getting their child ready for school and may be late to work when their child is late to school.² Whalen et al³ showed that children's before-school ADHD symptoms reduced parenting effectiveness. Faraone et al⁴ surveyed 300 caregivers of stimulant-treated children and reported high levels of early morning functional impairments in the child. The child's impairments reduced the emotional well-being of parents, worsened the parent's functioning during the early morning, and increased conflict with siblings.

ADHD symptoms also impair patients in the late afternoon and evening. For children, symptom control into the evening hours is important for socialization, family interactions, and completing chores and homework.³ Many adults with ADHD have work schedules that require attention to work or driving during the evening. At home, ADHD symptoms in adults interfere with parenting and marital interactions. A survey of 201 families with ADHD children² found that symptoms were high during dinner and evening homework time and at bedtime.

For children, the standard for assessing time-specific symptoms is the laboratory school study, which brings ADHD children together for a day in a simulated classroom during which they are assessed for behaviors relevant to ADHD.⁵ Likewise, workplace simulations have been developed to assess the duration of ADHD symptoms in adults with ADHD.⁶ Although simulation methods provide valuable data about the duration of action of medications,⁷ they place patients in artificial environments. A classroom that includes only ADHD children may not

You are prohibited from making this PDF publicly available.

Clinical Points

- The early morning and the late afternoon/evening are times of distress and disability for youth with attention-deficit/hyperactivity disorder.
- The present work provides clinicians with normative data on validated assessment tools that will allow them to classify their patients' risks for early morning and late afternoon/evening functional impairments.
- These norms provide a rational approach for selecting treatments to target such time-specific functional impairments.

generalize to real classrooms. Simulated environments do not assess behaviors that are important in the naturalistic environment such as getting to school on time, driving home from work, staying seated for dinner, and settling down at bedtime. For assessing naturalistic behaviors, researchers and clinicians need reliable and valid measures to test the time-specific efficacy of treatments.

We have been evaluating the psychometric features of 2 time-sensitive measures for youth with ADHD: the Before School Functioning Questionnaire (BSFQ)⁸ and the Parent Rating of Evening and Morning Behaviors Scale, Revised (PREMB-R).^{9,10} The BSFQ comprises 20 items that assess ADHD-associated symptoms and functional impairments prior to attending school or other activities. The BSFQ investigator-rated scale has very good internal homogeneity (Cronbach $\alpha = .91$), good test-retest reliability (correlation = 0.60), good concurrent validity (correlations = 0.42–0.86), and a strong treatment effect in a clinical trial of the methylphenidate transdermal system (effect size = -0.93).¹¹ The BSFQ separated drug from placebo in trials of guanfacine extended release (parent-reported scale)¹² and delayed-release and extended-release methylphenidate (DR/ER-MPH, formerly HLD200) (investigator-rated scale).¹³

The PREMB-R Morning (AM) subscale measures early morning functional (EMF) impairments; the Evening (PM) subscale assesses functional impairment in the late afternoon and evening. Michelson et al¹⁴ reported that atomoxetine was significantly better than placebo for reducing ADHD symptoms but was not effective for reducing early morning or late afternoon/evening functional impairments. In contrast, the early morning and late afternoon/evening scores were improved with atomoxetine in subsequent studies.^{15–17} The PREMB-R AM has significant internal homogeneity ($\alpha = .65$) and test-retest reliability (correlations = 0.52–0.45) and good concurrent validity (correlations = 0.50–0.71).⁹ The PREMB-R PM also has high internal homogeneity ($\alpha = .81$), good test-retest reliability (correlations = 0.74–0.83), and good concurrent validity (correlations = 0.47–0.80).¹⁰

Although prior work with the PREMB-R AM and PM and the BSFQ shows them to be reliable and valid assessments of early morning and late afternoon/evening behaviors, no normative data are available to help clinicians determine when a child's level of impairment is clinically significant.

Such norms would also be useful in clinical trials to address the degree to which treatments return ADHD patients to the normative range of functioning. To address that gap in the literature, we conducted a population survey of 1,200 families with the goal of establishing cut-points for each scale for use in clinical practice and research.

METHODS

Research Participants

The normative online survey was conducted on a sample of 1,200 primary caregivers of children and adolescents (6–17 years old). They were recruited through a consumer research panel (N = 275,000) representative of the US population in terms of age, sex, education, employment, ethnicity, and geography. Participants completed a screening questionnaire to assess the following inclusion criteria: (1) they self-identified as a primary caregiver of a child aged 6 to 17 years; (2) their child either never had ADHD, had a history of ADHD, or currently had ADHD without treatment (medical, psychological, or holistic) during the past 3 months; and (3) their child had not been diagnosed with birth defects of the brain or spinal cord injury, traumatic brain injury, severe concussions, cerebral palsy, seizure disorders, tic disorders or Tourette disorder, bipolar disorder, or psychosis. To determine if the child had any ADHD, other psychiatric disorders, or exclusionary disorders, we asked the parent if the child had ever been diagnosed by a health professional, a method that has been used, for example, by the Centers for Disease Control and Prevention (CDC).¹⁸ We did not, however, query for specific DSM-5 or other structured diagnostic criteria. If there was more than one eligible child, respondents were instructed to select and answer the subsequent questions based on the child whose birthday was next. We recruited parents of 50 male and 50 female youth in each of the 12 age groups. We did not preferentially recruit families that had a child with ADHD.

Eligible caregivers rated their children on the BSFQ and the PREMB-R. The survey was fielded and completed in June 2016 by REPASS, Inc (Newtown Square, Pennsylvania), a market research firm. Surveys were completed voluntarily and anonymously, and respondents were blinded to the research sponsor (Ironshore Pharmaceuticals & Development, Inc; Grand Cayman, Cayman Islands). The survey was conducted in adherence to the Marketing Research Association's Code of Marketing Research Standards (<https://www.insightsassociation.org/issues-policies/mra-code-marketing-research-standards>). The survey contained no option for any of the 18 standard types of protected health information (PHI) data to be collected and therefore did not require Institutional Review Board (IRB) approval.

Assessments

The BSFQ comprises 20 items that cover commonly reported areas of dysfunction in the early morning,

It is illegal to post this copyrighted PDF on any website.

Table 1. Demographic and Clinical Data^a

Characteristic	Overall (N = 1,200)	Non-ADHD (n = 1,079)	ADHD (n = 121)	P Value ^b
Age of child, mean (SD), y	11.50 (3.45)	11.43 (3.45)	12.09 (3.44)	.047
Male	600 (50.0)	524 (48.6)	76 (62.8)	.004
Respondent relationship				.799
Mother	787 (65.6)	711 (65.9)	76 (62.8)	
Father	342 (28.5)	304 (28.2)	38 (31.4)	
Stepmother	39 (3.3)	36 (3.3)	3 (2.5)	
Stepfather	32 (2.7)	28 (2.6)	4 (3.3)	
Comorbid conditions				
Birth defects of brain/spinal cord/deformity	0 (0)	0 (0)	0 (0)	NA
Traumatic brain injury/severe concussions	0 (0)	0 (0)	0 (0)	NA
Cerebral palsy	0 (0)	0 (0)	0 (0)	NA
Epilepsy or seizure disorders	0 (0)	0 (0)	0 (0)	NA
Tic disorders/Tourette disorder	0 (0)	0 (0)	0 (0)	NA
Psychosis	0 (0)	0 (0)	0 (0)	NA
Bipolar disorder	0 (0)	0 (0)	0 (0)	NA
Cigarette smoking	8 (0.7)	6 (0.6)	2 (1.7)	.414
Substance use disorder	7 (0.6)	4 (0.4)	3 (2.5)	.024
Oppositional defiant disorder	16 (1.3)	5 (0.5)	11 (9.1)	<.001
Conduct disorder	8 (0.7)	4 (0.4)	4 (3.3)	.002
Depression	43 (3.6)	36 (3.3)	7 (5.8)	.264
Insomnia/sleep disorders	26 (2.2)	21 (1.9)	5 (4.1)	.216
Anxiety	61 (5.1)	47 (4.4)	14 (11.6)	.001
Obsessive-compulsive disorder	13 (1.1)	6 (0.6)	7 (5.8)	<.001
Learning disability	57 (4.8)	36 (3.3)	21 (17.4)	<.001
Developmental disorder	17 (1.4)	9 (0.8)	8 (6.6)	<.001
Other	96 (8.0)	87 (8.1)	9 (7.4)	.949
None of those listed	939 (78.3)	876 (81.2)	63 (52.1)	<.001

^aValues shown as n (%) unless otherwise noted.

^bStatistical differences between non-ADHD and ADHD samples were determined by *t* test or χ^2 test, as appropriate.

Abbreviations: ADHD = attention-deficit/hyperactivity disorder, NA = not applicable, SD = standard deviation.

before-school activities (eg, breakfast, hygiene, time awareness, getting to school), associated with ADHD.⁸ Using the BSFQ, primary caregivers rated their child's EMF impairment between the time of awakening and before the school day or other morning activities (ie, 6:00 AM and 9:00 AM) on a severity scale of 0 to 3. The PREMB-R AM comprises 3 items that assess the difficulty of functioning during the early morning, and the PREMB-R PM comprises 8 items that assess the difficulty of functioning during the late afternoon and evening. Using the PREMB-R AM and PREMB-R PM, primary caregivers rated their child's temporal functional impairment on a severity scale of 0 to 3.

Statistical Analyses

Because the rating scale data have only integer values that are right truncated by the number of items and are markedly non-normally distributed, statistical analyses relied on truncated Poisson regression with scale scores as dependent variables. To predict ADHD diagnosis from each individual item score and total scores, we used logistic regression. We used multinomial logistic regression for trinary ADHD diagnoses (no history of ADHD vs history of ADHD vs current untreated ADHD). Each model was adjusted for the number of the following psychiatric comorbidities: cigarette smoking, substance use disorder, oppositional defiant disorder, conduct disorder, depression, insomnia/sleep disorders, anxiety disorder, obsessive-compulsive disorder, learning disability, and developmental disorder.

We determined if scale scores were significantly affected by age and sex after adjusting for the presence of ADHD and other comorbidities using a Bonferroni-adjusted α level of .008. We then computed normative cutoff scores using the same percentile cut-points defined by DuPaul et al¹⁹ for the ADHD Rating Scale:

- Screening risk: 80th percentile
- Mild functional impairment: 90th percentile
- Moderate functional impairment: 93rd percentile
- Severe functional impairment: 98th percentile

Because the prevalence of ADHD in this sample (10.1%) is about twice the population expectation of 5%,²⁰ we computed percentiles after randomly deleting half of the ADHD participants with the goal of estimating cut-points that would generalize to the overall population.

RESULTS

Of the 1,200 families surveyed, 121 index youth had been diagnosed with ADHD. Among these, 80 had current ADHD but had not been treated within the 3 months prior to participation and 41 did not currently have ADHD and were not receiving treatment. This prevalence of 10% is consistent with that reported by the CDC using similar methodology¹⁸ but larger than the population expectation of 5% from meta-analyses.²⁰ The ADHD group was half a year older and

Table 2. Effects of Sex, Age, and Psychiatric Disorders on the Total BSFQ Score

Characteristic	n	Mean	SD	P Value ^a
Age group				<.001
Children (aged 6–12 y)	700	15.48	11.89	
Adolescents (aged 13–17 y)	500	12.20	12.18	
Sex				.554
Male	600	14.47	12.52	
Female	600	13.77	11.70	
Current ADHD				<.001/<.001 ^b
Non-ADHD	1,079	12.70	11.00	
ADHD	121	26.74	14.16	
History of ADHD				<.001
Non-ADHD	1,079	12.70	11.00	
History of ADHD	41	21.17	14.58	
Current untreated ADHD	80	29.60	13.15	
Other psychiatric disorders				<.001
None	939	12.14	11.06	
One or more	261	21.22	13.08	

^aA *t* test was used for comparisons of children versus adolescents, males versus females, no comorbidities versus ≥ 1 comorbidity, and non-ADHD versus ADHD (untreated); a nonparametric Kruskal-Wallis 1-way analysis of variance was used for comparisons between no history of ADHD (non-ADHD), history of ADHD, and current untreated ADHD.

^bP value adjusted for the following comorbid disorders/conditions with a logistic regression model: cigarette smoking, substance use disorder, oppositional defiant disorder, conduct disorder, depression, insomnia/sleep disorders, anxiety disorder, obsessive-compulsive disorder, learning disability, and developmental disorder.

Abbreviations: ADHD = attention-deficit/hyperactivity disorder, BSFQ = Before School Functioning Questionnaire, SD = standard deviation.

was more likely to be male (Table 1). There were large and significant differences for most psychiatric disorders.

Adolescents had significantly lower BSFQ scores than young children; the effect of sex was not significant (Table 2). Those with a history of ADHD or other psychiatric disorders had higher BSFQ scores than other youth. Table 3 shows the same pattern of results for the PREMB-R AM and PM. Both ADHD status and number of other disorders were significant, independent predictors of BSFQ and PREMB-R scores. The BSFQ also asks respondents “How many minutes did it take your child to wake up and get out of bed?” and “How many minutes did it take your child to complete their early morning routine?” For the first question, both ADHD groups were significantly more likely to take more time getting out of bed than the non-ADHD group even after adjustment for significant effects of age, sex, and psychiatric comorbidity ($P < .001$). The mean \pm SD times were 16.7 ± 12.2 minutes for those with a history of ADHD and 10.9 ± 10.2 minutes for others. Responses to the second question did not differ significantly by ADHD status. The mean \pm SD times were 33.2 ± 18.9 minutes for those with a history of ADHD and 30.9 ± 18.3 minutes for those without such a history.

For all 3 scales, the distribution of total scores was heavily skewed. Many children had low or no impairment, and decreasing numbers of children showed higher levels of impairment. The distributions of these scores are shown in Supplementary Figure 1. The distributions of severity for each item of the BSFQ are given in Supplementary Figure 2. Across all 20 items of the BSFQ, youth with current untreated ADHD or a history of ADHD were rated as more severely ill than those without ADHD (all unadjusted P values $< .001$),

even after adjusting for psychiatric comorbidity (all adjusted P values $< .001$). Supplementary Figure 3 shows a similar pattern of results for the PREMB-R. Across the 3 items of PREMB-R AM, youth with current untreated ADHD or a history of ADHD had more severe ratings than those without ADHD, even after adjusting for psychiatric comorbidity (all unadjusted P values $< .001$; all adjusted P values $< .001$, except for item 1 [$P = .01$]). Across the 8 items of the PREMB-R PM, youth with current untreated ADHD or a history of ADHD had more severe ratings than those without ADHD, even after adjusting for psychiatric comorbidity (all unadjusted P values $< .001$; all adjusted P values $< .001$).

Because each of the total scores was associated with age but not sex, we defined 4 cut-points based on age alone: screening risk (80th percentile), mild functional impairment (90th percentile), moderate functional impairment (93rd percentile), and severe functional impairment (98th percentile). We used 4 age bins: 6 to 8, 9 to 11, 12 to 14, and 15 to 17 years. These cut-points are given in Table 4. We also computed cut-points for a data set that was modified to have a prevalence of ADHD, which is more consistent with the population expectation (5%) from meta-analyses.¹⁹ These data are also included in Table 4. These cut-points are slightly, albeit not dramatically, lower. As a third step, we smoothed these cut-points to be consistent with our finding that all 3 scores decrease with age. Despite that highly significant effect, 4 cut-points violate that pattern due to natural variability. We smoothed these by replacing them and the cut-point for the prior age group with the mean of those 2 cut-points. This method pools data across adjacent age groups to compute a common cut-point. The modified cut-points are bolded in Table 4.

Figure 1 shows the fraction of subjects exceeding each cut-point stratified by ADHD and comorbidity status. For example, Figure 1A shows that 78% of children having both ADHD and another disorder screen positive at the 80th percentile; 53% of those having only ADHD but not another psychiatric disorder also screen positive at this cut-point. For nearly all scores and cut-points, children with ADHD and another disorder are most likely to exceed the cut-point, those with only ADHD are second most likely, those without ADHD but with another disorder are third most likely, and those with no disorder are least likely.

For the BSFQ questions asking about the amount of time to get out of bed, the 80th percentile was 15 minutes, the 90th percentile was 25 minutes, the 93rd percentile was 30 minutes, and the 98th percentile was 43 minutes. For time to complete the early morning routine, these percentiles were 45 minutes, 60 minutes, 60 minutes, and 80 minutes, respectively (the data were too sparse to separate the 90th and 93rd percentiles).

DISCUSSION

The prevalence of ADHD in this sample was similar to a prior report from the CDC¹⁸ that used a telephone interview to query about prior diagnoses of ADHD but higher than

It is illegal to post this copyrighted PDF on any website.

Table 3. Effects of Sex, Age, and Psychiatric Disorders on the Total PREMB-R AM and PREMB-R PM Scores

Characteristic	n	PREMB-R AM			PREMB-R PM		
		Mean	SD	P Value ^a	Mean	SD	P Value ^a
Age group				.042			<.001
Children (aged 6–12 y)	700	2.56	2.24		6.22	5.33	
Adolescents (aged 13–17 y)	500	2.33	2.24		5.03	5.30	
Sex				.155			.823
Male	600	2.39	2.26		5.85	5.53	
Female	600	2.53	2.22		5.60	5.17	
Current ADHD				<.001/<.001 ^b			<.001/<.001 ^b
Non-ADHD	1,079	2.27	2.13		5.05	4.80	
ADHD	121	4.15	2.49		11.76	6.16	
History of ADHD				<.001			<.001
Non-ADHD	1,079	2.27	2.13		5.05	4.80	
History of ADHD	41	4.07	2.69		10.27	6.70	
Current untreated ADHD	80	4.19	2.39		12.53	5.77	
Other psychiatric disorders				<.001			<.001
None	939	2.15	2.08		4.81	4.88	
One or more	261	3.59	2.43		9.04	5.66	

^aA *t* test was used for comparisons in between children versus adolescents, males versus females, no comorbidities versus ≥ 1 comorbidity, and non-ADHD versus ADHD (untreated); a non-parametric Kruskal-Wallis 1-way analysis of variance was used for comparisons between no history of ADHD (non-ADHD), history of ADHD, and current untreated ADHD.

^bP value adjusted for the following comorbid disorders/conditions with a logistic regression model: cigarette smoking, substance use disorder, oppositional defiant disorder, conduct disorder, depression, insomnia/sleep disorders, anxiety disorder, obsessive-compulsive disorder, learning disability, and developmental disorder.

Abbreviations: ADHD = attention-deficit/hyperactivity disorder; PREMB-R AM = Parent Rating of Morning and Evening Behavior-Revised, Morning subscale; PREMB-R PM = PREMB-R Evening subscale, SD = standard deviation.

Table 4. Norm-Referenced Percentile Cutoff Points for the BSFQ, PREMB-R AM, and PREMB-R PM

Age, y	Cutoff Points for Original Data Set (10% ADHD Prevalence), by Percentile				Cutoff Points for Modified Data Set (5% ADHD Prevalence), by Percentile				Cutoff Points for Smoothed Modified Data Set (5% ADHD Prevalence), ^a by Percentile			
	80th	90th	93rd	98th	80th	90th	93rd	98th	80th	90th	93rd	98th
BSFQ												
6–8	26	33.5	37.5	42.5	25	33	35	41	25	33	36	43
9–11	24	33.5	38	45	22	30	36	45	22	30	36	43
12–14	24	32	36	41.5	22	30	34	41	22	30	34	42
15–17	22	30	33	45	21	28	31	42	21	28	31	42
PREMB-R AM												
6–8	5	6	6	8	5	6	6	7	5	6	6	8
9–11	5	6	6	8	4	6	6	8	4	6	6	8
12–14	4	6	6	7	4	5	6	7	4	6	6	7
15–17	5	6	6	8	4	6	6	7	4	6	6	7
PREMB-R PM												
6–8	11	15	16	19	11	15	16	18	11	15	16	19
9–11	9.5	14.5	16	19.5	9	13	16	19	9	13	16	19
12–14	10	13	15.5	18.5	9	12	14	17	9	13	15	18
15–17	10	14	15	18.5	9	13.5	15	18	9	13	15	18

^aSmoothed data are rounded up for fractional results (values in bold).
 Abbreviations: ADHD = attention-deficit/hyperactivity disorder; BSFQ = Before School Functioning Questionnaire; PREMB-R AM = Parent Rating of Evening and Morning Behavior, Revised, Morning subscale; PREMB-R PM = PREMB-R Evening subscale.

in data derived from a meta-analysis²⁰ of studies that interviewed respondents about symptoms and impairment. We found a higher male versus female prevalence of ADHD and higher rates of psychiatric disorders in ADHD compared with non-ADHD participants—findings highly consistent with prior literature.²¹


The total scores from the BSFQ, PREMB-R AM, and PREMB-R PM were significantly associated with ADHD and with the presence of other psychiatric disorders. The

effects of ADHD on scale scores was not accounted for by comorbid psychiatric disorders, which suggests that ADHD has an independent effect on the early morning and late afternoon/evening functioning of children and adolescents. Respondents rated youth with ADHD as showing greater severity for each item of the BSFQ, PREMB-R AM, and PREMB-R PM. For most items, about half of youth with current ADHD or a history of ADHD were rated as showing moderate or severe functional impairment.

You are prohibited from making this PDF publicly available.

It is illegal to post this copyrighted PDF on any website.

Figure 1. Proportion Exceeding Each Percentile Cutoff Point for the (A) BSFQ, (B) PREMB-R AM, and (C) PREMB-R PM by Presence of ADHD and Other Psychiatric Disorders



You are prohibited from making this PDF publicly available.

Abbreviations: ADHD = attention-deficit/hyperactivity disorder; BSFQ = Before School Functioning Questionnaire; PREMB-R AM = Parent Rating of Evening and Morning Behavior Scale, Revised, Morning subscale; PREMB-R PM = PREMB-R Evening subscale.

It is illegal to post this copyrighted PDF on any website.

For all scale scores, we found a significant effect for age but not for sex. The decrease in symptoms with age is consistent with other studies of ADHD,²² which show that symptoms of hyperactivity and impulsivity tend to decrease in adolescence whereas symptoms of inattention persist. The lack of sex effects raises the possibility that temporally specific measures of ADHD-associated features may be more sensitive to the disorder in females than diagnostic items. If so, that could have implications for future revisions of the diagnostic criteria for ADHD.

We presented the first norms for these 3 scores, stratifying by age. Inspection of Figure 1 motivates several recommendations. Because the PREMB-R AM has only 3 items, it cannot distinguish mild from moderate functional impairments, which suggests that the BSFQ is better for identifying EMF impairment. The 4 cut-points selected few children without ADHD and without other psychiatric disorders. This finding indicates that if these scores are applied to population samples, there would not be many false-positive identifications, especially for the more extreme cut-points. For the ADHD sample, the screening criterion (80th percentile) would select 50% to 80% of patients. The 90th percentile identifies 25% to 50% of ADHD youth as having early morning or late afternoon or evening impairments. Given that so many ADHD youth show extreme deviations from the normative level of functioning, clinicians should consider screening for such impairments, even among youth currently receiving treatment, as such youth have been documented to have EMF impairments.^{2,4}

In clinical practice, clinicians should use the normative cut-point that fits the context of the clinical setting and the population to be assessed. For example, in a pediatrics

clinic, where it is feasible to follow up many positive screens with a clinical interview of the parent, one could use a highly sensitive threshold (the 80th or 90th percentiles). When follow-up interviews are less feasible, the 93rd or 98th percentiles allow fewer false-positive findings at the price of not identifying all youth with ADHD who have impairments. When screening youth with ADHD using the BSFQ and PREMB-R PM, the mild functional impairment criterion identifies between 30% and 50% of ADHD youth as impaired (Figure 1) and should be considered as a useful threshold for optimizing early morning and late afternoon/evening functioning.

Our work is limited by using online surveys rather than in-person interviews and not collecting information about diagnostic criteria. Instead, we relied on the respondent's report about prior psychiatric diagnoses of their child. Although not ideal, this method has been used by others¹⁸ and is unlikely to have caused false-positive associations between ADHD and time-specific functional impairments. Nevertheless, future work should attempt to replicate these findings using structured interview diagnoses of ADHD that document age at onset, impairment, and the cross-situational expression of symptoms and impairment.

In summary, functional impairments in the early morning and late afternoon/evening hours are associated with ADHD and cannot be accounted for by other disorders. We have presented norms, stratified by age, which can guide clinicians in the use of these scores for identifying those ADHD youth who may be experiencing difficulties in these time periods. Such tools are needed given the availability of treatments that can target ADHD symptoms and impairments at these extremes of the daily routine.

Submitted: June 14, 2019; accepted September 24, 2019.

Published online: December 10, 2019.

Author contributions: All authors listed in the article have seen and approved the submission of this version of the article and take full responsibility for the content.

Potential conflicts of interest: In the past year, **Dr Faraone** received income, potential income, travel expenses continuing education support, and/or research support from Tris; Otsuka; Arbor; Ironshore Pharmaceuticals & Development, Inc; Shire; Akili Interactive; Enzymotec; Sunovion; Supernus; and Genomind. With his institution, he has US patent US20130217707 A1 for the use of sodium-hydrogen exchange inhibitors in the treatment of attention-deficit/hyperactivity disorder (ADHD). He also receives royalties from books published by Guilford Press: *Straight Talk about Your Child's Mental Health*; Oxford University Press: *Schizophrenia: The Facts*; and Elsevier: *ADHD: Non-Pharmacologic Interventions*. He is principal investigator of www.adhdinadults.com. **Dr Wilens** receives or has received grant support from the National Institutes of Health (NIH) National Institute on Drug Abuse (NIDA) and Lloyd Foundation. He is or has been a consultant for NIH NIDA; Ironshore Pharmaceuticals & Development, Inc; KemPharm; Otsuka/Neurovance; the National Football League Program for Substances of

Abuse (ERM Associates); and Minor/Major League Baseball. He has a published book with Guilford Press: *Straight Talk About Psychiatric Medications for Kids* and co-edited *ADHD Across the Lifespan* (Cambridge Univ Press) and *Comprehensive Clinical Psychiatry* (Elsevier). **Mr DeSousa** and **Dr Incledon** are employees/consultants of Ironshore Pharmaceuticals & Development, Inc. **Dr Sallee** is an employee of Ironshore Pharmaceuticals Inc and is a member of the board of directors for P2D Bioscience. **Dr Komolova** is an employee of Highland Therapeutics Inc.

Funding/support: This work was funded by Ironshore Pharmaceuticals & Development, Inc.

Role of the sponsor: Ironshore Pharmaceuticals & Development, Inc, paid for the collection and analysis of data. Employees of Ironshore Pharmaceuticals & Development, Inc, and its affiliated companies were involved in the design of the study and interpretation of results and, hence, are included as authors.

Acknowledgments: The authors acknowledge Rick Nullmeier, BA, of ADVANCE Life Sciences Consulting (Hebron, Kentucky), who provided project management support; and Helen Doll, DPhil, and Hannah Lewis, PhD, of ICON (Gaithersburg, Maryland), who performed initial data analysis. These individuals were paid by the study sponsor for services provided.

Supplementary material: Available at PSYCHIATRIST.COM.

REFERENCES

1. Barkley RA, Cunningham CE. The effects of methylphenidate on the mother-child interactions of hyperactive children. *Arch Gen Psychiatry*. 1979;36(2):201-208.
2. Sallee FR. Early morning functioning in stimulant-treated children and adolescents with attention-deficit/hyperactivity disorder, and its impact on caregivers. *J Child Adolesc Psychopharmacol*. 2015;25(7):558-565.
3. Whalen CK, Henker B, Jamner LD, et al. Toward mapping daily challenges of living with ADHD: maternal and child perspectives using electronic diaries. *J Abnorm Child Psychol*. 2006;34(1):115-130.
4. Faraone SV, Schachar RJ, Barkley RA, et al. Early morning functional impairments in stimulant-treated children with attention-deficit/hyperactivity disorder versus controls: impact on the family. *J Child Adolesc Psychopharmacol*. 2017;27(8):715-722.
5. Wigal SB, Wigal TL. The laboratory school protocol: its origin, use, and new applications. *J Atten Disord*. 2006;10(1):92-111.
6. Fried R, Surman C, Hammerness P, et al. A controlled study of a simulated workplace laboratory for adults with attention deficit hyperactivity disorder. *Psychiatry Res*. 2012;200(2-3):949-956.
7. Gomeni R, Fang LL, Bressolle-Gomeni F, et al. A general framework for assessing In vitro/In vivo

It is illegal to post this copyrighted PDF on any website.

- as a tool for maximizing the benefit-risk ratio of a treatment using a convolution-based modeling approach. *CPT Pharmacometrics Syst Pharmacol*. 2019;8(2):97–106.
8. Faraone SV, Hammerness PG, Wilens TE. Reliability and validity of the Before-School Functioning Scale in children with ADHD. *J Atten Disord*. 2018;22(11):1040–1048.
 9. Faraone SV, Childress A, Wigal SB, et al. Reliability and validity of the Daily Parent Rating of Evening and Morning Behavior Scale, Revised. *J Atten Disord*. 2018;22(11):1066–1073.
 10. Faraone SV, DeSousa NJ, Sallee FR, et al. Psychometric validation of the Before School Functioning Questionnaire and Parent Rating of Evening and Morning Behavior Scale, Revised in children with attention-deficit/hyperactivity disorder. *J Am Acad Child Psychiatry*. 2017;56(10):S212.
 11. Wilens TE, Hammerness P, Martelon M, et al. A controlled trial of the methylphenidate transdermal system on before-school functioning in children with attention-deficit/hyperactivity disorder. *J Clin Psychiatry*. 2010;71(5):548–556.
 12. Wilens TE, McBurnett K, Turnbow J, et al. Morning and evening effects of guanfacine extended release adjunctive to psychostimulants in pediatric ADHD. *J Atten Disord*. 2017;21(2):110–119.
 13. Pliszka SR, Wilens TE, Bostrom S, et al. Efficacy and safety of HLD200, delayed-release and extended-release methylphenidate, in children with attention-deficit/hyperactivity disorder. *J Child Adolesc Psychopharmacol*. 2017;27(6):474–482.
 14. Michelson D, Allen AJ, Busner J, et al. Once-daily atomoxetine treatment for children and adolescents with attention deficit hyperactivity disorder: a randomized, placebo-controlled study. *Am J Psychiatry*. 2002;159(11):1896–1901.
 15. Sutton V, Sumner C, Allen AJ, et al. Validity, Reliability, and Responsiveness of the DPREMB-R Scale for ADHD. Annual Meeting of the American Academy of Child and Adolescent Psychiatry. 2003; Miami Beach, Florida.
 16. Sangal RB, Owens J, Allen AJ, et al. Effects of atomoxetine and methylphenidate on sleep in children with ADHD. *Sleep*. 2006;29(12):1573–1585.
 17. Kelsey DK, Sumner CR, Casat CD, et al. Once-daily atomoxetine treatment for children with attention-deficit/hyperactivity disorder, including an assessment of evening and morning behavior: a double-blind, placebo-controlled trial. *Pediatrics*. 2004;114(1):e1–e8.
 18. Visser SN, Danielson ML, Bitsko RH, et al. Trends in the parent-report of health care provider-diagnosed and medicated attention-deficit/hyperactivity disorder: United States, 2003–2011. *J Am Acad Child Adolesc Psychiatry*. 2014;53(1):34–46.e2.
 19. DuPaul GJ, Reid R, Anastopoulos AD, et al. Parent and teacher ratings of attention-deficit/hyperactivity disorder symptoms: factor structure and normative data. *Psychol Assess*. 2016;28(2):214–225.
 20. Polanczyk GV, Willcutt EG, Salum GA, et al. ADHD prevalence estimates across three decades: an updated systematic review and meta-regression analysis. *Int J Epidemiol*. 2014;43(2):434–442.
 21. Faraone SV, Asherson P, Banaschewski T, et al. Attention-deficit/hyperactivity disorder. *Nat Rev Dis Primers*. 2015;1(1):15020.
 22. Faraone SV, Biederman J, Mick E. The age-dependent decline of attention deficit hyperactivity disorder: a meta-analysis of follow-up studies. *Psychol Med*. 2006;36(2):159–165.

See supplementary material for this article at PSYCHIATRIST.COM.

You are prohibited from making this PDF publicly available.



THE JOURNAL OF
CLINICAL PSYCHIATRY

THE OFFICIAL JOURNAL OF THE AMERICAN SOCIETY OF CLINICAL PSYCHOPHARMACOLOGY

Supplementary Material

Article Title: Functional Impairment in Youth With ADHD: Normative Data and Norm-Referenced Cutoff Points for the Before School Functioning Questionnaire and Parent Rating of Evening and Morning Behavior Scale-Revised

Author(s): Stephen V. Faraone, PhD; Norberto J. DeSousa, MA; Marina Komolova, PhD; F. Randy Sallee, MD, PhD; Bev Incledon, PhD; and Timothy E. Wilens, MD

DOI Number: <https://doi.org/10.4088/JCP.19m12956>

List of Supplementary Material for the article




1. [Figure 1](#) Histograms of the (A) BSFQ, (B) PREMB-R AM and (C) PREMB-R PM Total Scores
2. [Figure 2](#) Distribution of Severity Ratings on the 20 Items of the BSFQ by ADHD Status
3. [Figure 3](#) Distribution of Severity Ratings on the 3 Items of the PREMB-R AM and the 8 Items of PREMB-R PM by ADHD Status

Disclaimer

This Supplementary Material has been provided by the author(s) as an enhancement to the published article. It has been approved by peer review; however, it has undergone neither editing nor formatting by in-house editorial staff. The material is presented in the manner supplied by the author.


SUPPLEMENTARY MATERIAL

Supplementary Figure 1: Histograms of the (A) BSFQ, (B) PREMB-R AM and (C) PREMB-R PM Total Scores

A BSFQ**B PREMB-R-AM****C PREMB-R-PM**

BSFQ, Before School Functioning Questionnaire; PREMB-R AM, Parent Rating of Morning and Evening Behavior-Revised, Morning subscale; PREMB-R PM, Parent Rating of Morning and Evening Behavior-Revised, Evening subscale.


Supplementary Figure 2. Distribution of Severity Ratings on the 20 Items of the BSFQ by ADHD Status^a




^aStatistical differences across the three groups (ie, youth without ADHD [n = 1079], those with a history of ADHD [n = 41], and those with current untreated ADHD [n = 80]) were determined by Chi-square test and adjusted for the following comorbid disorders/conditions with a logistic regression model: cigarette smoking, substance use disorder, oppositional defiant disorder, conduct disorder, depression, insomnia/sleep disorders, anxiety disorder, obsessive compulsive disorder, learning disability, and developmental disorder; ^bUnadjusted P value; ^cAdjusted P value. ADHD, attention-deficit/hyperactivity disorder; BSFQ, Before School Functioning Questionnaire.

Supplementary Figure 3. Distribution of Severity Ratings on the 3 Items of the PREMB-R AM and the 8 Items of PREMB-R PM by ADHD Status^a

PREMB-R AM



PREMB-R PM



^a Statistical differences across the three groups (ie, youth without ADHD [n = 1079], those with a history of ADHD [n = 41], and those with current untreated ADHD [n = 80]) were determined by Chi-square test and adjusted for the following comorbid disorders/conditions with a logistic regression model: cigarette smoking, substance use disorder, oppositional defiant disorder, conduct disorder, depression, insomnia/sleep disorders, anxiety disorder, obsessive compulsive disorder, learning disability, and developmental disorder; ^b Unadjusted P value; ^c Adjusted P value.

ADHD, attention-deficit/hyperactivity disorder; PREMB-R AM, Parent Rating of Morning and Evening Behavior-Revised, Morning subscale; PREMB-R PM, Parent Rating of Morning and Evening Behavior-Revised, Evening subscale.