

### A Crossover Study of Prolactin Changes Associated With Risperidone and Olanzapine

**Sir:** We read with interest the recent article by Melkersson<sup>1</sup> that described the differential effect on prolactin elevation in schizophrenic patients treated with clozapine (none of the patients had elevated prolactin), olanzapine (24% of the patients had elevated prolactin), and risperidone (89% of the patients had elevated prolactin). Previous head-to-head clinical trials have revealed similar results.<sup>2-5</sup> However, these comparison studies might be limited by the individual variation in endocrine profiles, demographic variables such as age and gender, and concomitant medication use.<sup>6,7</sup> We designed a crossover study to compare the prolactin level changes in patients switching from olanzapine to risperidone or vice versa.

**Method.** This observational crossover study compared the prolactin levels associated with olanzapine and risperidone treatment in patients with DSM-IV schizophrenia. The subjects were treated in the routine outpatient setting and had poor or partial response to risperidone or olanzapine for at least 3 months, defined by a Clinical Global Impressions-Severity of Illness scale<sup>8</sup> score of more than 4 (moderately ill). They were willing to change to the other medication (olanzapine or risperidone) and gave full informed consent. Patients with specific medical diseases, such as diabetes mellitus, dyslipidemia, cardiovascular diseases, and hypertension, and patients who required hospitalization for an acute exacerbation of psychotic illness were not included in this study. The institutional ethics review board approved this research, and the study was conducted from January 2004 to June 2005.

After the new agent was started, the patients' previous antipsychotic agents were completely discontinued in 2 weeks. The clinicians were free to titrate the dose of the second drug according to the patient's clinical condition. The concomitant agents, including anticholinergics (biperiden in 2 patients),  $\beta$ -blocker (propranolol in 2 patients), and benzodiazepines (lorazepam in 8 patients, estazolam in 1 patient, and fludiazepam in 1 patient), were kept at the same dose in the study period. No other antipsychotics or hormonal treatments were coadministered during the preswitch treatment of at least 3 months or for the 3 months after the switch. Prolactin levels were assessed following overnight (> 12 hours) fasting before medication switch and 3 months after crossover. The paired t test was used to compare prolactin levels for baseline and post-baseline differences with the Statistical Package for the Social Sciences, Version 9.0 (SPSS Inc., Chicago, Ill.). The difference in prolactin level was considered statistically significant if a p value was equal to or less than .05.

**Results.** Seventeen patients (8 women and 9 men) with schizophrenia completed this study. Their mean  $\pm$  SD age was  $34.9 \pm 7.5$  (range, 22–46) years. In the 9 patients taking risperidone at the time of inclusion (risperidone-first group), there was a significant ( $p = .004$ ) decrease in prolactin level after the switch to olanzapine ( $59.9 \pm 38.5$   $\mu\text{g/L}$  before switch;  $13.6 \pm 9.7$   $\mu\text{g/L}$  after switch). In the other 8 patients (olanzapine-first group), prolactin level was increased after the switch to risperidone ( $26.1 \pm 19.7$   $\mu\text{g/L}$  before switch;  $40.0 \pm 16.2$   $\mu\text{g/L}$  after switch), but the difference did not reach a significant level ( $p = .106$ ). When all patients ( $N = 17$ ) were compared after olanzapine treatment and after risperidone treatment (irrespective of the order), the mean prolactin level was significantly higher ( $p = .001$ ) in patients receiving risperidone ( $50.3 \pm 31.1$   $\mu\text{g/L}$ ) than in those receiving olanzapine ( $19.5 \pm 16.1$   $\mu\text{g/L}$ ).

There were no significant differences in the mean doses of olanzapine ( $10.0 \pm 4.3$  mg/day for risperidone-first group vs.  $8.1 \pm 2.6$  mg/day for olanzapine-first group) or risperidone ( $3.7 \pm 1.8$  mg/day for risperidone-first group vs.  $2.6 \pm 0.7$  mg/day for olanzapine-first group) taken as first or second drugs. The mean  $\pm$  SD durations of pharmacotherapy prior to the switch of risperidone and olanzapine were  $17.3 \pm 6.9$  weeks and  $18.0 \pm 9.1$  weeks, respectively. However, there was an imbalanced male-to-female distribution between the risperidone-first group (3 to 6) and olanzapine-first group (6 to 2).

To our knowledge, this is first crossover comparison study of prolactin changes in the same patients, switching from olanzapine to risperidone or vice versa. This study supports an association of elevated levels of serum prolactin with the use of risperidone as compared with olanzapine, which is consistent with the results of most previous head-to-head comparison studies.<sup>2-5</sup> When patients switched from risperidone to olanzapine, serum prolactin level decreased significantly. This is consistent with the results of Kim and colleagues' study.<sup>9</sup> However, their study included only female subjects switched from risperidone to olanzapine (not crossover), and the study duration was only 2 weeks.

Our study has some limitations. First, probably due to the small sample size, the increase of prolactin level in patients switching from olanzapine to risperidone did not reach a significant level. Second, due to an observational study design, the patients were not free from antipsychotic drugs at enrollment, the male-to-female distribution between the risperidone-first group (3 to 6) and the olanzapine-first group (6 to 2) was not balanced, the durations of risperidone and olanzapine treatments were not the same, and the assignment to risperidone or olanzapine as first treatment was not randomized. Finally, the dosage of both drugs was not controlled, although there were no significant differences between groups.

Nevertheless, our study clearly showed that in the average clinical dosage, elevated levels of prolactin are associated with the use of risperidone and decrease with olanzapine. On the basis of our findings, we recommend that prolactin should be closely monitored in patients during treatment with atypical antipsychotic drugs, especially with risperidone.

*The work was supported by grants DOH 94F044, DMR-92-017, DOH94-DC-1111, and CMU93-M-24 from the Department of Health, China Medical University and Hospital, Taichung, Taiwan. This study received no financial support from the pharmaceutical industry.*

*The authors report no additional financial or other relationship relevant to the subject matter of this letter.*

#### REFERENCES

- Melkersson K. Differences in prolactin elevation and related symptoms of atypical antipsychotics in schizophrenic patients. *J Clin Psychiatry* 2005;66:761–767
- Volavka J, Czobor P, Cooper TB, et al. Prolactin levels in schizophrenia and schizoaffective disorder patients treated with clozapine, olanzapine, risperidone, or haloperidol. *J Clin Psychiatry* 2004;65:57–61
- Kinon BJ, Gilmore JA, Liu H, et al. Hyperprolactinemia in response to antipsychotic drugs: characterization across comparative clinical trials. *Psychoneuroendocrinology* 2003;28(suppl 2):69–82
- Ahl J, Kinon BJ, Liu-Seifert H. Sexual dysfunction associated with neuroleptic-induced hyperprolactinemia improves with reduction in prolactin levels. *Ann N Y Acad Sci* 2004;1032:289–290
- Tran PV, Hamilton SH, Kuntz AJ, et al. Double-blind comparison of olanzapine versus risperidone in the treatment of schizophrenia and

- other psychotic disorders. *J Clin Psychopharmacol* 1997;17:407–418
6. Su KP, Wu PL, Pariante CM. A crossover study on lipid and weight changes associated with olanzapine and risperidone. *Psychopharmacology (Berl)* 2005;183:383–386
  7. Su KP, Shen WW, Chuang CL, et al. A pilot cross-over design study on QTc interval prolongation associated with sulpiride and haloperidol [letter]. *Schizophr Res* 2003;59:93–94
  8. Guy W. ECDEU Assessment Manual for Psychopharmacology. US Dept Health, Education, and Welfare publication (ADM) 76-338. Rockville, Md: National Institute of Mental Health; 1976:218–222
  9. Kim KS, Pae CU, Chae JH, et al. Effects of olanzapine on prolactin levels of female patients with schizophrenia treated with risperidone. *J Clin Psychiatry* 2002;63:408–413

**Cher-Yeang Lin, M.D.**

**Po-Lun Wu, M.D.**

Department of Psychiatry  
China Medical University Hospital  
Taichung, Taiwan

**Carmine M. Pariante, M.D., Ph.D.**

Institute of Psychiatry  
King's College London  
London, United Kingdom

**Kuan-Pin Su, M.D.**

Department of Psychiatry  
China Medical University Hospital  
Taichung, Taiwan

## Recurrent Priapism Associated With Use of Aripiprazole

**Sir:** Priapism is one of the most serious adverse effects of psychotropic medications. It may be defined simply as abnormally prolonged erection of the penis (or clitoris) that is usually but not invariably painful. Though it is relatively rare as an adverse effect of psychotropic medication, as many as 25% to 40% of priapism cases are attributable to psychotropics.<sup>1,2</sup> Treatment of priapism within 4 to 6 hours is important because the condition may cause permanent erectile dysfunction and even penile gangrene. That several of the first-generation antipsychotics may be associated with priapism has been recognized for decades, but a few instances have been reported with all of the second-generation antipsychotics except aripiprazole. In their review, Compton and Miller<sup>1</sup> noted priapism to be associated with clozapine, olanzapine, and risperidone. Subsequently, priapism has been attributed to ziprasidone<sup>3,4</sup> and quetiapine,<sup>5,6</sup> as well. Herein we report what we believe is the first published case of priapism associated with aripiprazole.

**Case report.** Mr. A, a 47-year-old African American man diagnosed with chronic paranoid schizophrenia 15 years ago, had diabetes mellitus, hypertension, and hyperlipidemia and had been off all psychotropic medication for several years. His outpatient medications were atorvastatin and metformin. He was then prescribed aripiprazole in 2005 by his outpatient psychiatrist, and the first episode of prolonged and painful penile erection started about 6 hours after the first dose of aripiprazole. He was treated at another hospital with intracavernosal phenylephrine, which alleviated the priapism. However, the priapism soon recurred and he presented to our hospital with a painful erection of about 7 hours' duration. Cavernal irrigation and injection of phenylephrine resulted in prompt detumescence. This treatment was repeated when the priapism recurred 2 days later. The patient refused a shunt procedure for recurrent priapism. He then proceeded to have 3 more episodes over a total of 7 days, even

though no additional doses of aripiprazole or another anti-psychotic were taken, but responded to cavernosal irrigation and phenylephrine each time. In addition, he was put on pseudoephedrine treatment, which has been reported as being helpful for priapism.<sup>7</sup>

During the hospital stay, the psychosis was quiescent and he was treated with diazepam 15 mg/day only. The patient then remained free of prolonged erections for the next 2 days and was discharged. He was asked to continue pseudoephedrine, apply ice packs to the penis, and abstain from sexual activity. Further monitoring and treatment of his psychiatric condition were deferred to the outpatient psychiatrist. After discharge he was lost to follow-up by us.

No prior incident of priapism had occurred in the patient's life. He had used cocaine and cannabis in the past, but there was no substance use in the 8 years prior to this admission. The urine drug screen was negative. Diabetes mellitus, hypertension, and hyperlipidemia and the medications atorvastatin and metformin are not known to be associated with priapism. Atorvastatin and metformin have no pharmacodynamic ( $\alpha$ -blockade) or pharmacokinetic interaction with aripiprazole. There was no history of sickle-cell disease, perineal trauma, blood-cell dyscrasia, neoplasm, chronic prostatitis, or spider bite, all of which have been associated with priapism.

We used the strategy recommended by Mago<sup>8</sup> to be followed prior to publishing case reports. A search of MEDLINE (PubMed) and EMBASE was conducted on January 23, 2006, using the terms *aripiprazole* and *priapism*, both as text words and as medical subject heading (MeSH) terms. The only case found in the literature search that was potentially related to the present one was that of priapism developing in a 16-year-old boy upon addition of oxcarbazepine to the patient's existing regimen of aripiprazole (5 mg/day) and lithium.<sup>9</sup> The authors attributed the priapism to oxcarbazepine, and after this was discontinued, the patient continued taking lithium and aripiprazole without further prolonged erections.

The common feature of many medications capable of producing priapism appears to be  $\alpha$ -adrenergic blockade, though the fact that medications with no significant  $\alpha$ -adrenergic blockade can also be associated with priapism suggests that several different mechanisms may operate. Aripiprazole has significant  $\alpha_1$ -adrenergic receptor blocking activity.<sup>10</sup> This consideration and the onset of the condition within a few hours after the first dose of aripiprazole suggest that this drug was indeed the cause of priapism in our patient. The recurrence of priapism several times over a week in this patient can be explained by the long half-life of aripiprazole. The mean elimination half-life for aripiprazole and its active metabolite are about 75 and 94 hours, respectively.<sup>11</sup>

A variety of psychoactive medications have been associated with priapism. These include antidepressants (selective serotonin reuptake inhibitors, venlafaxine, trazodone, nefazodone, bupropion), anxiolytics (buspirone, hydroxyzine), and antipsychotics (both first-generation and second-generation). Of the second-generation antipsychotics, clozapine, olanzapine, risperidone, ziprasidone, and quetiapine have been reported to be associated with priapism. Our case indicates that aripiprazole should be added to this list. In addition, the case illustrates that when adverse effects due to aripiprazole occur, they may persist for several days due to the long half-life of the drug and its active metabolite, and the patient should be managed and monitored accordingly.

*Dr. Mago is a consultant for Bristol-Myers Squibb; has received grant/research support from Bristol-Myers Squibb, AstraZeneca, and*

GlaxoSmithKline; and has served on the speakers or advisory boards for Bristol-Myers Squibb and Pfizer. Dr. Kunkel has served on the speakers or advisory boards for Wyeth, Forest, Pfizer, and Intramed. Mr. Anolik and Dr. Johnson report no financial or other relationship relevant to the subject of this letter.

## REFERENCES

1. Compton MT, Miller AH. Priapism associated with conventional and atypical antipsychotic medications: a review. *J Clin Psychiatry* 2001;62:362–366
2. Thompson JW Jr, Ware MR, Blashfield RK. Psychotropic medication and priapism: a comprehensive review. *J Clin Psychiatry* 1990;51:430–433
3. Reeves RR, Kimble R. Prolonged erections associated with ziprasidone treatment: a case report [letter]. *J Clin Psychiatry* 2003;64:97–98
4. Kaufman KR, Stern L, Mohebbati A, et al. Ziprasidone-induced priapism requiring surgical treatment. *Eur Psychiatry* 2006;21:48–50
5. Davol P, Rukstalis D. Priapism associated with routine use of quetiapine: case report and review of the literature. *Urology* 2005;66:880
6. Harrison G, Dilley JW, Loeb L, et al. Priapism and quetiapine in an HIV-positive male [letter]. *J Clin Psychopharmacol* 2006;26:100–101
7. Lowe FC, Jarow JP. Placebo-controlled study of oral terbutaline and pseudoephedrine in management of prostaglandin E1-induced prolonged erections. *Urology* 1993;42:51–54
8. Mago R. Searching the medical literature for case reports [letter with reply]. *J Clin Psychiatry* 2004;65:444–445
9. Negin B, Murphy TK. Priapism associated with oxcarbazepine, aripiprazole, and lithium [letter]. *J Am Acad Child Adolesc Psychiatry* 2005;44:1223–1224
10. Shapiro DA, Renock S, Arrington E, et al. Aripiprazole, a novel atypical antipsychotic drug with a unique and robust pharmacology. *Neuropsychopharmacology* 2003;28:1400–1411
11. Winans E. Aripiprazole. *Am J Health Syst Pharm* 2003;60:2437–2445

**Rajnish Mago, M.D.**  
**Robert Anolik, A.B.**  
**Robert A. Johnson, M.D.**  
**Elisabeth J. S. Kunkel, M.D.**  
 Department of Psychiatry  
 Thomas Jefferson University  
 Philadelphia, Pennsylvania

### Intimate Partner Violence Victimization Among Adults With Severe Mental Illness: Results of a Cross-Sectional Study

**Sir:** There is strong evidence that adults with severe mental illness evidence high rates of both lifetime trauma exposure (particularly physical and sexual abuse) and posttraumatic stress disorder (PTSD) relative to the general population.<sup>1</sup> However, little attention has yet been paid in this population to intimate partner violence (IPV), a form of trauma that is often recurrent and linked with long-term mental and physical health consequences.

**Method.** Data were obtained (but not reported) from a cross-sectional study<sup>2</sup> of a random sample of public-sector psychiatric patients recruited from a day-hospital program from 2002 to 2004. A total of 156 randomly identified patients were approached for study participation; 142 provided informed consent, and 141 (79 men, 62 women) completed relevant study measures. We examined rates of adult sexual and physical abuse perpetrated by a romantic partner by using the Trauma Assessment for Adults<sup>3</sup> (TAA; for review of psychometrics, see reference 3), as well as characteristics of these assaults (e.g.,

self-reported fear of being killed or seriously injured at the time of the assault, physical injury, and whether the assault occurred more than once). Due to the nature of the TAA, we did not have the data to assess incidents of IPV perpetration, and thus, we focus exclusively on IPV victimization in our results. We also examined PTSD Checklist<sup>4</sup> scores among those with a history of IPV.

**Results.** Of the final sample, 21 patients (14.9%) reported at least one incident of IPV victimization. Only 1 (1.3%) of 79 men and 20 (32.3%) of 62 women reported a history of IPV victimization. Although not mutually exclusive, 5 (25.0%) of 20 patients (Ns for IPV victimization variables ranged from 19 to 21 due to missing data) also reported physical force or threat of force for unwanted sexual contact, 9 (42.9%) of 21 patients reported a physical assault with a weapon, and 15 (71.4%) of 21 patients reported a physical assault without a weapon by a partner. Seventeen (89.5%) of 19 patients thought they would be killed or seriously injured at the time of the assault, 14 (73.7%) of 19 patients suffered a physical injury, and 14 (73.7%) of 19 patients reported multiple incidents of IPV victimization. The mean (SD) score for those endorsing IPV victimization on the PTSD checklist was 43.57 (19.03), with 8 participants (38.1%) endorsing a score of 50 or above (indicative of PTSD) and 4 (19.0%) endorsing a score between 40 and 49 (clinically meaningful subthreshold symptoms).

A startlingly high number (32.3%) of female public-sector patients with severe mental illness have histories of IPV victimization, and IPV victimization experiences are both recurrent and harmful. These data are notable in that the TAA does not specifically inquire about IPV-related assaults (i.e., the interviewer asks about the perpetrator only after the patient reports a physical or sexual assault), and thus, these rates are likely an underestimate of IPV victimization in this population. Our results are all the more troublesome in light of recent findings by Heru and colleagues<sup>5</sup> indicating that over 90% of suicidal inpatients reported a past year history of IPV victimization and perpetration. These findings clearly suggest that the presence of severe and recurrent forms of IPV place individuals with severe mental illness, who already constitute a vulnerable population, at extreme risk. Thus, as public-sector clinicians increase their efforts to screen for trauma and PTSD in this population, particular attention should be paid to assessing for the presence and severity of both IPV victimization and perpetration using more comprehensive IPV measures such as the Revised Conflict Tactics Scales.<sup>6</sup>

*This study was supported by National Institute of Mental Health grant #R01 MH65517.*

*Drs. Grubaugh and Frueh report no additional financial affiliation or other relationship relevant to the subject matter of this letter.*

## REFERENCES

1. Mueser KT, Salyers MP, Rosenberg SD, et al. Interpersonal trauma and posttraumatic stress disorder in patients with severe mental illness: demographic, clinical, and health correlates. *Schizophr Bull* 2004;30:45–57
2. Frueh BC, Knapp RG, Cusack KJ, et al. Patients' reports of traumatic or harmful experiences within the psychiatric setting. *Psychiatr Serv* 2005;56:1123–1133
3. Resnick HS, Best CL, Kilpatrick DG, et al. Review of Trauma Assessment for Adults (TAA). In: Stamm BH, ed. *Measurement of Stress, Trauma, and Adaptation*. Lutherville, Md: Sidran Press; 1996:362–365
4. Blanchard EB, Jones-Alexander J, Buckley TC, et al. Psychometric properties of the PTSD Checklist (PCL). *Behav Res Ther* 1996;34:669–673

5. Heru AM, Stuart GL, Rainey S, et al. Prevalence and severity of intimate partner violence and associations with family functioning and alcohol abuse in psychiatric inpatients with suicidal intent. *J Clin Psychiatry* 2006;67:23–29
6. Straus MA, Hamby SL, Boney-McCoy S, et al. The Revised Conflict Tactics Scales (CTS-2). *J Fam Issues* 1996;17:283–316

**Anouk L. Grubaugh, Ph.D.**  
**B. Christopher Frueh, Ph.D.**

Department of Psychiatry and Behavioral Sciences  
Medical University of South Carolina  
Charleston, South Carolina

### Internet Assessment of Obsessive-Compulsive Disorder

**Sir:** The Internet may provide an important and low-cost means to improve communication between patients and clinicians through e-mail and medical information Web sites. In the past year, several reports have also appeared of the successful use of Internet Web sites to help reduce symptoms of anxiety, depression, and phobic avoidance (see, for example, Botella et al.<sup>1</sup> and Schneider et al.<sup>2</sup>).

Here, we report data on Web site usability ratings, as well as age, gender, and OCD severity, from individuals who completed an Internet OCD behavioral assessment.

**Method.** We developed an Internet Web site for anonymous self-assessment and treatment planning by sufferers of obsessive-compulsive disorder (OCD) that included components from a computerized interactive telephone system called BT-STEPS, which was found in a randomized controlled trial to effectively guide behavioral self-assessment and treatment.<sup>3</sup>

Print advertisements were placed in a Boston newspaper and the Obsessive-Compulsive Foundation newsletter, and of 111 OCD sufferers who visited the Web site in response to the advertisements, 83 (75%) successfully completed a brief 1-session behavioral assessment (including identification of primary rituals and situations that trigger them, family involvement, and interference with life functioning) and initial behavior therapy treatment plan that they could print out and share with their therapist, if desired.

**Results.** Although we expected that most visitors to the Web site would be technologically savvy users younger than 30 years, the mean age of completers was 40.4 years (SD = 12.9), and of the 77 users who gave their age, 22 (29%) were aged 40 to 49 years, another 17 (22%) were aged 50 to 64 years, and 3 (4%) were 65 years or older. Of 80 subjects answering the gender question, 45 (56%) were female.

All users rated the Web site as either “very easy” or “somewhat easy” to use, and, significantly, ease of use was unrelated to age ( $r = 0.13$ ,  $p < .25$ ). Those with more severe OCD (as measured by the Yale-Brown Obsessive Compulsive Scale) were more likely to rate the Web site as “very easy” to use.

Given the shortage of clinicians skilled in effective behavioral treatments for OCD, Internet self-assessment and treatment planning appear to be feasible. Although there remains a substantial “digital divide” in Internet use based on income and education,<sup>4</sup> among our assessable respondents, age, gender, and OCD severity were not roadblocks to ease of use of Internet self-assessment. This bodes well for the future acceptance of Web sites containing reliable medical information, since according to a survey, among the next generation of senior

citizens—those currently 50 to 64 years old—the Internet has surpassed television, books, magazines, newspapers, and radio as a source of “a lot” of health information.<sup>4</sup>

*Dr. Baer shares intellectual property rights to the BT-STEPS program mentioned in the letter. Dr. Minichiello reports no financial affiliation or other relationship relevant to the subject matter of this letter.*

### REFERENCES

1. Botella C, Hofmann SG, Moscovitch DA. A self-applied, internet-based intervention for fear of public speaking. *J Clin Psychol* 2004;60:881–886
2. Schneider AJ, Mataix-Cols D, Marks IM, et al. Randomized controlled trial in phobia/panic disorder of two forms of net-guided self-help, each with brief live phone support. *Psychother Psychosom* 2005;74:154–164
3. Greist JH, Marks IM, Baer L, et al. Behavior therapy for obsessive-compulsive disorder guided by a computer or by a clinician compared with relaxation as a control. *J Clin Psychiatry* 2002;63:138–145
4. Kaiser Family Foundation. e-Health and the Elderly: How Seniors Use the Internet for Health Information. Key Findings From a National Survey of Older Americans. January 2005. Available at: <http://www.kff.org/entmedia/upload/e-Health-and-the-Elderly-How-Seniors-Use-the-Internet-for-Health-Information-Key-Findings-From-a-National-Survey-of-Older-Americans-Survey-Report.pdf>. Accessibility verified June 7, 2006

**Lee Baer, Ph.D.**  
**William E. Minichiello, Ed.D.**  
Department of Psychiatry  
Harvard Medical School  
Boston, Massachusetts

### Hypertriglyceridemia Associated With Direct Effects of Olanzapine Rather Than With Weight Gain: A Case Report

**Sir:** Hypertriglyceridemia and weight gain are well-documented side effects associated with olanzapine therapy.<sup>1</sup> Thus, it has become important to monitor weight and lipid profile including triglycerides in patients taking olanzapine or any other atypical antipsychotic. A total cholesterol level is made up of very-low-density lipoprotein (VLDL), low-density lipoprotein (LDL), and high-density lipoprotein (HDL). Triglycerides are a component of the VLDL portion of the total cholesterol.<sup>2</sup> It has been proposed that the lipid changes mentioned above with olanzapine are related to weight gain<sup>3</sup>; however, weight gain often does not seem to correlate to the severity of the hypertriglyceridemia observed. The relevance of weight gain to lipid changes remains unclear.<sup>3</sup> With both weight gain and hypertriglyceridemia occurring in many cases, it is difficult to delineate the direct effects of olanzapine versus the indirect effects associated with olanzapine-induced obesity. We report a case that demonstrated improvement of hyperlipidemia after olanzapine discontinuation even though the patient continued to be overweight.

**Case report.** Mr. A was a 22-year-old white man diagnosed with DSM-IV schizophrenia. He had a long history of hospitalizations with a very long index admission, allowing us to view lipid changes over time. Prior to initiation of olanzapine therapy in January 1998, Mr. A had a total cholesterol level within normal range (Table 1), but there were no triglyceride measurements. Olanzapine was initiated, and his total cholesterol level

Table 1. Medication Doses, Plasma Olanzapine and Lipid Levels, and Weight Changes in Temporal Sequence<sup>a</sup>

Day	Olanzapine		Triglycerides (mg/dL) <sup>c</sup>	Cholesterol (mg/dL) <sup>d</sup>	Gemfibrozil Dose (mg/d) <sup>e</sup>	Weight (kg)	Body Mass Index
	Dose (mg/d)	Plasma Level (ng/mL) <sup>b</sup>					
-547	0	...	...	179	0	...	...
1 <sup>f-h</sup>	10	...	...	...	0	64.4	24.4
258 <sup>f,g,i</sup>	35	50	802	293	0	77.5	29.3
351 <sup>f,g</sup>	35	...	1410	305	0	81.3	30.8
367 <sup>f,g</sup>	35	...	...	...	1200	81.1	30.7
378 <sup>f,g</sup>	35	...	775	258	1200	80.4	30.5
408 <sup>g</sup>	35	...	348	219	1200	79.5	30.1
436 <sup>g</sup>	35	67	152	121	1200	80.8	30.6
449	0	...	AWOL	AWOL	0	AWOL	AWOL
472 <sup>j</sup>	10	0 <sup>k</sup>	155 <sup>k</sup>	216 <sup>k</sup>	0	79.5	30.1
497	20	32	311	265	1200	78.5	29.7
532 <sup>i</sup>	35	...	729	248	0	79.0	29.9
679	35	...	537	267	0	80.8	30.6
708	35	...	579	278	0	80.4	30.5

<sup>a</sup>Some lipid profiles are not described.

<sup>b</sup>Plasma olanzapine levels measured within 3 weeks of lipid profile.

<sup>c</sup>Reference range, 40–160 mg/dL.

<sup>d</sup>Reference range, 120–200 mg/dL.

<sup>e</sup>Patient refused gemfibrozil during some periods.

<sup>f</sup>Patient was taking fluphenazine 25 mg/day, which was later decreased and discontinued.

<sup>g</sup>Patient was taking bupropion 4 mg/day, which was later decreased and discontinued.

<sup>h</sup>Day of first olanzapine dose.

<sup>i</sup>During short periods of time, the patient agreed to be on a low-calorie diet; most of the time, he refused.

<sup>j</sup>The patient returned to the hospital after 23 days (day 472). During his absence, the patient took no medications.

<sup>k</sup>Blood was collected before the patient was given the first 10-mg dose after his return to the hospital.

Abbreviation: AWOL = absent without leave. Symbol: ... = not available.

increased. One year after olanzapine initiation, his triglyceride level was 1410 mg/dL, and his cholesterol level was 305 mg/dL (weight = 81.3 kg). Gemfibrozil treatment lowered the patient's triglyceride level when he was compliant with medication (Table 1).

After 14 months, the patient went away without leave (escaped from the hospital). Upon his return after 23 days with no medication, his olanzapine level was not detectable. His lipids were measured, revealing a triglyceride level of 155 mg/dL and total cholesterol level of 216 mg/dL (weight = 79.5 kg). These lipid levels in the absence of gemfibrozil treatment were some of the lowest lipid levels recorded in the patient since initiation of olanzapine. In the year after Mr. A's return to the hospital and reinitiation of olanzapine at 35 mg/day, limited changes were seen in his weight (range, 78.5–80.8 kg), but his triglyceride level, in the absence of gemfibrozil, increased to a range of 537 to 729 mg/dL (Table 1).

To our knowledge, this is the first case demonstrating changes in olanzapine's triglyceridemic effects that were unrelated to weight gain and that were seen following the temporal pattern of presence and absence of olanzapine in the patient's system. The pattern of olanzapine treatment was on-off-on, while the pattern of the olanzapine-induced weight gain was on-on-on. The patient described in this letter consumed no alcohol during his hospitalization in a closely supervised locked unit, although he reported consuming some beers during his absence from the hospital. He had no signs or symptoms of cirrhosis, nephrotic syndrome, pancreatitis, or other causes of increased triglycerides; thus, in this patient, olanzapine appeared to have

a direct effect of increasing triglycerides independent of weight gain.

Hopefully, future research will discern mechanisms of action related to the lipid changes seen with olanzapine and other atypical antipsychotics, allowing for the creation of novel molecules or modifications of current molecules without such deleterious metabolic changes.

*This case was published without external support.*

*In the past 3 years, Dr. de Leon has been on the advisory boards of Bristol-Myers Squibb and Roche Molecular Systems, Inc.; received researcher-initiated grants from Eli Lilly and Roche Molecular Systems, Inc.; and lectured with support from Eli Lilly (once), Roche Molecular Systems, Inc. (5 times), and Bristol-Myers Squibb (once). Dr. Markham-Abedi reports no financial or other relationship relevant to the subject of this letter.*

## REFERENCES

1. Osser DN, Najaren DM, Dufresne RL. Olanzapine increases weight gain and serum triglyceride levels. *J Clin Psychiatry* 1999;60:767–770
2. Zipes DP. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 7th ed. Philadelphia, Pa: Saunders; 2005
3. Meyer JM. Novel antipsychotics and severe hyperlipidemia. *J Clin Psychopharmacol* 2001;21:369–373

**Courtney Markham-Abedi, M.D.**

**Jose de Leon, M.D.**

University of Kentucky Mental Health Research Center  
Eastern State Hospital  
Lexington, Kentucky