

The Role of Dopamine in Obsessive-Compulsive Disorder: Preclinical and Clinical Evidence

Damiaan Denys, M.D., Ph.D.; Joseph Zohar, M.D.;
and Herman G. M. Westenberg, Ph.D.

Obsessive-compulsive disorder (OCD) is a frequent and chronic psychiatric disorder that has been linked closely to the serotonin system mainly because of the antiobsessional efficacy of selective serotonin reuptake inhibitors (SSRIs). A limitation of the serotonin hypothesis of OCD is that a substantial number of the patients with OCD show no significant improvement after an adequate trial with SSRIs. There is substantial evidence that these patients may benefit from addition of antipsychotics to their ongoing SSRI treatment, suggesting that dopamine also might play a role in the pathophysiology of OCD. In this review, the preclinical and clinical evidence on the role of dopamine in OCD is summarized.
(J Clin Psychiatry 2004;65[suppl 14]:11-17)

Obsessive-compulsive disorder (OCD) is a chronic psychiatric disorder characterized by recurrent persistent thoughts (obsessions) and/or repetitive compulsory behaviors (compulsions). Over the past 2 decades, it has been suggested that OCD might be related to the functioning of brain serotonin systems. This hypothesis is based largely on the notion that selective serotonin reuptake inhibitors (SSRIs) possess antiobsessional efficacy.¹⁻³ There is now growing evidence that the dopamine system may be involved in OCD as well.⁴⁻⁶ In this article, the preclinical and clinical evidence supporting the role for dopamine in the pathophysiology of OCD will be reviewed. Evidence for the involvement of dopamine in OCD is derived from studies using animal models of OCD and from clinical studies using neurochemical, pharmacologic, genetic, and neuroimaging strategies.

ANIMAL MODELS

Campbell and colleagues⁷ have investigated the behavioral consequences of transgenic stimulation of a regional

subpopulation of the dopamine neurons that express the dopamine-1 (D₁) receptor in the cortex and amygdala by generating mice that express an intracellular form of cholera toxin. The study suggests that chronic stimulation of these D₁-expressing neurons induces complex compulsive behavior that resembles symptoms of OCD in humans.⁸ Although these mice were resistant to behavioral inhibition by a D₁ receptor antagonist and supersensitive to the D₂ receptor antagonist sulpiride,⁹ Campbell and colleagues^{10,11} suggested that chronic potentiation of cortical and limbic D₁-expressing neurons may cause obsessive-compulsive behaviors.

In another animal model, in which rats are chronically treated with the selective D_{2/3} receptor agonist quinpirole, a ritual-like set of behavioral acts resembling OCD checking behavior was observed.¹²⁻¹⁵ Postmortem analyses in these animals revealed increased dopamine tissue levels in the nucleus accumbens and right prefrontal cortex.¹⁶ Joel and Avisar¹⁷ developed a rat model of OCD based on the hypothesis that a deficient response feedback mechanism underlies obsessions and compulsions. Rats undergoing extinction of lever-pressing for food after the attenuation of an external feedback for this behavior exhibit excessive lever-pressing unaccompanied by an attempt to collect a reward, which may be analogous to the excessive and unreasonable behavior seen in OCD. Administration of the D₁ receptor antagonist SCH 23390 reduced the number of compulsive lever-presses without affecting the number of lever-presses followed by an attempt to collect a reward.¹⁸ On the basis of electrophysiologic data, Joel and Doljansky¹⁸ suggested that compulsive lever-pressing depends on a phasic decrease in stimulation of D₁ receptors.¹⁸ These data, using putative animal models of OCD, suggest a role of dopamine,

From the Rudolf Magnus Institute of Neuroscience, Department of Psychiatry, University Medical Center Utrecht, Utrecht, The Netherlands (Drs. Denys and Westenberg), and the Division of Psychiatry, Chaim Sheba Medical Center, Tel Hashomer, Israel (Dr. Zohar).

This article is derived from the proceedings of the 6th International Obsessive-Compulsive Disorder Conference "Beyond Refractory Obsessions and Anxiety States: Toward Remission," which was held November 13-15, 2003, in Lanzarote, Spain, and supported by an unrestricted educational grant from Solvay Pharmaceuticals.

Corresponding author and reprints: Damiaan Denys, M.D., Ph.D., UMC, Dept. of Psychiatry (B.01.206), P.O. Box 85500, 3508 GA Utrecht, the Netherlands (e-mail: D.A.J.P.denys@azu.nl).

in particular of the D₁ and D₂ receptors, in the mechanism underlying compulsive behavior.

DOPAMINE AND METABOLITE CONCENTRATIONS IN HUMANS

Baseline measures of dopamine and its metabolite homovanillic acid (HVA) in psychotropic-naïve patients may provide direct evidence for a possible role of dopamine in OCD. Thoren and colleagues¹⁹ assessed HVA in cerebrospinal fluid (CSF) levels before and after 3 weeks of treatment with clomipramine but found no change. Benkelfat and coworkers²⁰ found no differences between the mean plasma HVA level of 13 medication-free patients with OCD and 29 normal controls. Swedo et al.,²¹ examining CSF levels of HVA in 43 children with primary OCD, reported that CSF HVA levels were not significantly related to OCD symptoms and did not correlate with improvement following 5 weeks of treatment with clomipramine. Hollander et al.,²² on the other hand, observed a significant decrease in plasma HVA in 12 OCD patients relative to 10 controls following fenfluramine treatment, and Zahn et al.²³ showed that CSF metabolites of serotonin and dopamine, but not of norepinephrine, were positively correlated with electrodermal responsivity in a reaction time task in 43 adolescents and children with OCD.

Marazziti et al.²⁴ measured platelet sulfotransferase activity in 17 drug-free OCD patients and an equal number of healthy controls. Sulfotransferase is an enzyme involved in the catabolism of catecholamines such as dopamine and has similar kinetic characteristics in brain and platelets. Their results showed a higher level of sulfotransferase activity in OCD patients compared with control subjects, suggesting an increased dopaminergic neurotransmission in OCD.

In summary, the neurochemical studies on the role of dopamine metabolites in OCD patients have, by and large, yielded no evidence for an abnormal dopamine function in OCD.

PHARMACOLOGIC CHALLENGE TESTS

Another approach to assessing the functioning of the dopaminergic system is the evaluation of the behavioral and neuroendocrine effects response to administration of indirect (cocaine and amphetamine) or direct (apomorphine [APO] and bromocriptine) dopamine receptor agonists. Cocaine, a dopamine transporter blocker, elevates synaptic dopamine levels and increases the dopamine transporter density.²⁵ It has been reported that chronic use of cocaine may be associated with stereotyped examining, searching and sorting behaviors, and an exacerbation of obsessive-compulsive symptoms.^{26–30} Moreover, cocaine-abusing patients are at increased risk for the later development of OCD.³¹ Methylphenidate and amphetamine have

been reported to exacerbate or induce^{30,32–36} as well as to improve^{37–39} OCD symptoms. A limitation of the use of cocaine, methylphenidate, and amphetamine as dopaminergic probes is that they also release serotonin and norepinephrine.

Pitchot et al.⁴⁰ assessed the growth hormone (GH) response to 0.5 mg of APO in 8 drug-free OCD patients and 8 healthy male volunteers. No difference in mean GH peak response was found, suggesting that the dopaminergic function was not altered in OCD. In contrast, Brambilla et al.,⁴¹ studying 15 patients with OCD and 15 age/sex-matched controls, found a blunted GH response to APO in OCD patients, hinting at a postsynaptic dopamine receptor subsensitivity. In the same sample, however, cortisol (CORT) responses to stimulation with APO were not different between the 2 groups.⁴² Longhurst et al.⁴³ examined the effects of catecholamine depletion in 6 drug-free patients with the tyrosine hydroxylase inhibitor alpha-methyl-*para*-tyrosine (AMPT) and found no significant changes in obsessive-compulsive symptom severity as compared with placebo. Although bromocriptine, a selective dopamine receptor agonist, has been shown to induce stereotypies in animals, one report describes an improvement of obsessive-compulsive symptoms in OCD patients.⁴⁴ Results from pharmacologic challenge studies with dopamine receptor agonists in OCD are as yet inconsistent, but there are indications that obsessive-compulsive symptoms may be related to increased dopamine neurotransmission.

PHARMACOTHERAPY

Indirect evidence for a role of dopamine in OCD comes from treatment studies using pharmacologic agents that modulate the dopaminergic activity in the brain.

Antipsychotics in Monotherapy

Three years after its introduction, chlorpromazine was tested in 75 outpatients with obsessional neurosis and allied disorders in a placebo-controlled trial.⁴⁵ A significant response to chlorpromazine as compared with placebo was observed in 27 patients (36%), but it was judged to be disappointing in relieving compulsive symptoms. Ever since, no placebo-controlled trial with a typical antipsychotic drug has been conducted in OCD. Except for some case reports,^{46–50} typical antipsychotics in monotherapy are considered to be ineffective in OCD, mainly on grounds of individual clinical experience. McDougle et al.⁵¹ assessed the efficacy of clozapine monotherapy in 12 adults with refractory OCD in a 10-week, open-label trial with clozapine but found no significant change. There have been no placebo-controlled trials with atypical antipsychotics, such as risperidone, olanzapine, quetiapine, or ziprasidone, as monotherapy for OCD.

Table 1. Overview of Addition Trials With Atypical Antipsychotics in Obsessive-Compulsive Disorder

Year of Publication	Author	Design	N	Dose, mg/day	Trial Length, wk	YBOCS Score Decrease, %	Responders, N (%)
Risperidone^a							
1995	McDougle et al ⁹⁰	Open	3	1	1–4	55	3 (100)
1996	Saxena et al ⁹¹	Open	21	0.5–8.0	3	NA	14 (67)
1997	Stein et al ⁹²	Open	8	1–2	4	22	3 (37)
2000	Pfanner et al ⁹³	Open	20	1–3	8	31	15 (75)
2000	McDougle et al ⁹⁴	Double-blind placebo	20	1–7	6	29	11 (55)
2002	Baxter et al ⁹⁵	Double-blind placebo	10	1	9	40	NA
2003	Hollander et al ⁹⁶	Double-blind placebo	10	0.5–3.0	8	...	4 (40)
Olanzapine^b							
1999	Weiss et al ⁹⁷	Open	10	1.25–20.00	8	40	7 (70)
2000	Koran et al ⁹⁸	Open	10	2.5–10.0	8	16	3 (30)
2000	Bogetto et al ⁹⁹	Open	23	5	12	30	NA
2001	Francobandiera ¹⁰⁰	Open	9	2.5–5.0	6	38	6 (66)
2002	Crocq et al ¹⁰¹	Open	8	5–15	4–74	...	7 (88)
2003	D'Amico et al ¹⁰²	Open	21	10	12	26	7 (33)
2004	Bystritsky et al ¹⁰³	Double-blind placebo	13	5–20	6	19	6 (46)
2004	Shapira et al ⁵⁴	Double-blind placebo	22	5–10	6	± 25	9 (41)
Quetiapine^c							
2002	Mohr et al ¹⁰⁴	Open	8	50–300	6	21	4 (50)
2002	Atmaca et al ¹⁰⁵	Single-blind placebo	14	50–200	8	56	10 (71)
2002	Denys et al ¹⁰⁶	Open	10	200	8	35	7 (70)
2003	Sevincok and Topuz ⁵⁵	Open	8	150	10	27	2 (25)
2004	Denys et al ¹⁰⁹	Double-blind placebo	20	200–300	8	34	8 (40)

^aFor all risperidone studies, total N = 92 and mean YBOCS score decrease = 35%.

^bFor all olanzapine studies, total N = 116 and mean YBOCS score decrease = 27%.

^cFor all quetiapine studies, total N = 60 and mean YBOCS score decrease = 34%.

Abbreviations: NA = not available, YBOCS = Yale-Brown Obsessive-Compulsive Scale.

It is of note that de novo emergence or exacerbation of OCD symptoms during treatment with antipsychotics has been described extensively in patients with psychotic disorders. Lykouras et al.⁵² have recently reviewed the reported cases of OCD symptoms induced by atypical antipsychotics.

Antipsychotics in Addition to Selective Serotonin Reuptake Inhibitors

Addition of pimozide and haloperidol to SSRIs proved effective for patients who were refractory to treatment, in particular for patients with comorbid chronic tic disorders or schizotypal personality disorders.⁵³ Following the successful combination of typical antipsychotics with SSRIs for treatment-refractory patients, there were a number of studies combining atypical antipsychotics with SSRIs (Table 1). Although 2 negative studies have been published,^{54,55} risperidone, olanzapine, and quetiapine appear to be efficacious in addition to SSRIs for therapy-refractory patients at low doses within 6 to 8 weeks. Recently, in an open trial, Metin et al.⁵⁶ evaluated the efficacy of 325 mg/day of amisulpride, a selective dopamine D_{2/3} antagonist, to augment the effect of SSRI treatment in 20 patients with treatment-resistant OCD and observed a significant improvement in 95% of the patients.

Mechanism of Action?

It is intriguing that antipsychotics in monotherapy lack efficacy in OCD, while they are capable of inducing

de novo OCD symptoms in psychotic disorders and are efficacious in combination with SSRIs in some patients with OCD. It has been proposed that serotonin-2A (5-HT_{2A}) receptor antagonism exacerbates obsessive-compulsive symptoms by increasing the firing rate of the dopamine neurons, whereas D₂ receptor antagonism reduces obsessive-compulsive symptoms through inhibition of the dopamine neurons.⁵⁷ Zhang et al.^{58,59} have shown in rats that the combination of olanzapine and fluoxetine may increase synergistically extracellular dopamine and norepinephrine levels in the prefrontal cortex, and Denys et al.⁶⁰ found that the combination of quetiapine and fluvoxamine may cause a synergistic dopamine increase in the prefrontal cortex and thalamus. Since the combination of antipsychotics and SSRIs does not result in augmented serotonin levels, it is unlikely that an altered serotonergic neurotransmission is underlying the clinical efficacy of this combination. Additional research is warranted to determine whether changes in extracellular dopamine levels may account for the clinical efficacy of the augmentation strategy with atypical antipsychotics in OCD.

NEUROIMAGING

In vivo neuroimaging of dopamine transporters and receptors with positron emission tomography (PET) or single photon emission computer tomography (SPECT) offers another tool with which to probe the dopaminergic function in OCD. Sawle et al.⁶¹ did not find an alteration in

Table 2. Overview of Dopamine-Related Association Studies in Obsessive-Compulsive Disorder (OCD)

Polymorphism	Year of Publication	Author	N	Results
Dopamine Transporter				
40-base-pair repeat in the DAT 1 gene	1998	Billett et al ⁷²	100 OCD patients and controls	Negative
40-base-pair repeat in the DAT 1 gene	2000	Frisch et al ⁶⁶	75 OCD patients and 172 controls	Negative
40-base-pair repeat in the DAT 1 gene	2003	Hemmings et al ⁶⁷	71 OCD patients and 129 controls	Negative
D₂ receptor				
Three exons of the DRD2 gene, 4, 5, 6	1994	Novelli et al ⁷¹	45 OCD patients and 26 controls	Negative
Taq-IA	1996	Nicolini et al ⁶⁹	66 OCD patients and 54 controls	OCD patients + tics had a higher frequency of Taq I A2 allele (p = .014)
Taq-IA and the serine/cysteine variation	1998	Billett et al ⁷²	100 OCD patients and controls	Negative
Taq-IA and the serine/cysteine variation	2004	Denys et al ⁷⁰	150 OCD patients and 150 controls	Male OCD patients had a higher frequency of Taq I A2 allele (p = .020)
D₃ receptor				
Msc I in the first exon	1994	Catalano et al ⁶⁸	97 OCD patients and 97 controls	Negative
Msc I in the first exon	1996	Nicolini et al ⁶⁹	67 OCD patients and 54 controls	Negative
Msc I in the first exon	1998	Billett et al ⁷²	100 OCD patients and controls	Negative
D₄ receptor				
13 base pair deletion in the first exon	1996	Di Bella et al ⁷⁵	157 OCD patients and 162 controls	Negative
48-base-pair repeat	1997	Cruz et al ⁷³	12 OCD patients + tics and 49 patients	OCD patients + tics had a higher prevalence of the 7-fold variant (p = .018)
48-base-pair repeat	1998	Billett et al ⁷²	100 OCD patients and controls	Negative
48-base-pair repeat	2000	Frisch et al ⁶⁶	75 OCD patients and 172 controls	Negative
48-base-pair repeat	2003	Hemmings et al ⁶⁷	71 OCD patients and 129 controls	Negative
48-base-pair repeat	2003	Millet et al ⁷⁴	55 trios	An absence of transmission of the allele 2 (p = .005)
COMT				
Val-158-Met substitution	1997	Karayorgou et al ⁷⁶	73 OCD patients and 148 controls	Male OCD patients had a higher frequency of LL genotype (p = .0002)
Val-158-Met substitution	1998	Ohara et al ⁸¹	17 OCD patients and 135 controls	Negative
Val-158-Met substitution	1999	Karayorgou et al ⁷⁷	110 nuclear OCD families	Preferential transmission of the L allele in male OCD patients (p = .0057)
Val-158-Met substitution	2000	Schindler et al ⁸⁰	72 OCD patients/parent trios	A tendency for association with homozygosity at COMT locus (p = .056)
Val-158-Met substitution	2002	Alsobrook et al ⁷⁸	56 OCD patients and 112 parents	Female OCD patients had a higher frequency of LL genotype (p = .049)
Val-158-Met substitution	2001	Niehaus et al ⁷⁹	54 OCD patients and 54 controls	OCD patients had a higher frequency of H/L genotype (p = .0017)
C → T transition	2001	Kinnear et al ¹⁰⁷	48 OCD patients and 48 controls	Negative
Val-158-Met substitution	2004	Denys et al ⁷⁰	150 OCD patients and 150 controls	Male OCD patients had a higher frequency of L alleles (p = .0035)
Val-158-Met substitution	2003	Erdal et al ⁸²	59 OCD patients and 114 controls	Negative
Val-158-Met substitution	2004	Meira-Lima et al ¹⁰⁸	79 OCD patients and 202 controls	Negative

Abbreviations: COMT = catechol-O-methyl transferase, DAT = dopamine transporter.

[¹⁸F]-6-Fluorodopa uptake into the caudate and putamen in a small (PET) study involving 6 OCD patients with obsessional slowness. A decrease in [¹⁸F]-6-Fluorodopa uptake is believed to reflect a reduction in the number of nigrostriatal dopaminergic neurons. Van der Wee et al.⁶² found higher binding ratios of [¹²³I] Beta-CIT to the dopamine transporter in the left basal ganglia in 15 drug-naïve patients with OCD. Kim et al.⁶³ found an increased dopamine transporter binding ratio in the right basal ganglia and a tendency toward an increased dopamine transporter binding ratio in the left basal ganglia with [¹²³I] IPT SPECT in

15 OCD patients. Pogarell et al.,⁶⁴ on the other hand, did not detect significant differences in striatal dopamine transporter [¹²³I] Beta-CIT binding between 7 patients and 10 controls. Finally, Denys et al.⁶⁵ observed a decreased [¹²³I] IBZM binding in the left caudate in 10 OCD patients, suggesting a down-regulated dopamine D₂ receptor. Altogether, these findings indicate that the dopamine system in the basal ganglia could be involved in OCD. Higher dopamine transporter densities in tandem with a down-regulation of the D₂ receptor suggest higher synaptic concentrations of dopamine in the basal ganglia in OCD.

ASSOCIATION STUDIES

A number of candidate gene studies have been conducted in order to elucidate the contribution of the dopamine system in OCD (Table 2). Investigations of the role of the dopamine transporter and D₃ receptor showed no significant association between a particular allele and OCD.⁶⁶⁻⁶⁹ Except for the TAQ I A2 allele that was found to be associated with OCD patients and tics in male patients,^{69,70} no statistically significant differences in allele frequencies or genotype between OCD patients and controls were found with regard to the D₂ receptor.^{71,72} Cruz et al.⁷³ and Millet et al.⁷⁴ found an association for the D₄ receptor, but other reports were negative.^{66,67,72,75}

The role of catechol-O-methyl transferase (COMT) has been extensively investigated in OCD since COMT is an enzyme that has a crucial role in the elimination of dopamine, and higher dopamine levels may be implicated in OCD. Karayiorgou et al.^{76,77} found evidence for an association between the low-activity COMT allele and OCD in male OCD patients in a case-control study and a family-based study, whereas Alsobrook et al.⁷⁸ found evidence pointing to an association between the low-activity COMT allele and OCD in female OCD patients. Niehaus et al.⁷⁹ reported a preponderance of COMT high/low heterozygotes in an Afrikaner population of 54 OCD patients but did not observe gender differences. Schindler et al.⁸⁰ found no association between any particular allele and OCD but found a tendency for an association with homozygosity at the COMT locus. Ohara et al.⁸¹ did not find any association in a small sample of 24 Japanese patients and neither did Erdal et al.⁸² in a sample of 59 Turkish patients. In line with the study by Karayiorgou et al.,⁷⁷ Denys et al.⁷⁰ found a higher frequency of the low-activity COMT in male OCD patients. On the other hand, a recent meta-analysis⁸³ of the COMT gene in 144 OCD patients and 337 controls showed insufficient evidence to support an association.

A HYPERDOPAMINERGIC STATE?

Despite some inconsistencies, in general, the results from most studies hint at an association between OCD and increased midbrain dopamine neurotransmission. The hypothesis of increased dopamine neurotransmission in the basal ganglia is in agreement with various working hypotheses of the pathophysiology of OCD such as the hyperactive corticostriatal model, the amygdalocentric model, or the model of behavioral addiction in OCD. The corticostriatal working model of OCD suggests an imbalance of the direct versus indirect pathway that produces a hyperactive circuit responsible for the repetitive behaviors seen in OCD.^{84,85} Since D₁ preferentially activates the direct and D₂ the indirect pathway, and the density of D₁ receptors in the basal ganglia is higher than the density of D₂ receptors, increased concentrations of dopamine are most

likely to result in a dominant D₁-regulated direct circuit and consequently in a hyperactive corticostriatal system.

In the amygdalocentric model, the acquisition and expression of conditioned fear mediated by the amygdala are believed to be actively inhibited by feedback mechanisms from the medial prefrontal cortex.⁸⁶ One way to understand OCD is that the normal cortical inhibition of the amygdala is malfunctioning and that the anxiety responses induced by the amygdala therefore become more intrusive and chronic in patients with OCD. Rosenkranz and Grace⁸⁷ demonstrated that the response of the amygdala to cortical inhibition is potentially affected by alterations of the mesolimbic dopaminergic system. When dopamine is increased, the ability of the prefrontal cortex to suppress the affective responses generated in the amygdala is attenuated.

A number of phenomenological characteristics of OCD such as loss of voluntary control, repetitiveness, compulsiveness, reinforcement of behavior, aberrant habit learning, and uncertainty resemble addictive behavior and may be understood within the conceptual framework of "behavioral addiction."⁸⁸ In particular, ritualistic-compulsive actions share similarities with addictive behavior. There is little doubt that midbrain dopamine has positive reinforcing properties, and it is liable that the reinforcing nature of compulsions originates from increased dopamine transmission. Recently, Fiorillo et al.⁸⁹ demonstrated that dopamine neurons show increased firing during prolonged periods of uncertainty. Subjective uncertainty is a main feature of pathologic gambling and plays a major role in OCD.

CONCLUSION

To explore these hypotheses, more direct studies of the dopamine function in OCD are needed, in particular, measurements of dopamine and its metabolite and pharmacologic challenge studies with dopaminergic probes during behavioral stimulation. To date, there is sufficient preclinical and clinical evidence that implicates the dopamine system in OCD, but more studies are warranted to understand the function of dopamine in the pathophysiology of OCD.

Drug names: apomorphine (Apokyn), bromocriptine (Parlodel and others), chlorpromazine (Thorazine, Sonazine, and others), clomipramine (Anafranil and others), clozapine (Fazaclo, Clozaril, and others), fluoxetine (Prozac and others), haloperidol (Haldol and others), methylphenidate (Metadate, Ritalin, and others), olanzapine (Zyprexa), pimozide (Orap), quetiapine (Seroquel), risperidone (Risperdal), ziprasidone (Geodon).

Disclosure of off-label usage: The authors have determined that, to the best of their knowledge, apomorphine, bromocriptine, chlorpromazine, clozapine, haloperidol, methylphenidate, olanzapine, pimozide, quetiapine, risperidone, and ziprasidone are not approved by the U.S. Food and Drug Administration for the treatment of obsessive-compulsive disorder.

REFERENCES

1. Barr LC, Goodman WK, Price LH, et al. The serotonin hypothesis of obsessive compulsive disorder: implications of pharmacologic challenge

- studies. *J Clin Psychiatry* 1992;53(4, suppl):17-28
2. Pigott TA. OCD: where the serotonin selectivity story begins. *J Clin Psychiatry* 1996;57(suppl 6):11-20
 3. Blier P, de Montigny C. Possible serotonergic mechanisms underlying the antidepressant and anti-obsessive-compulsive disorder responses. *Biol Psychiatry* 1998;44:313-323
 4. Goodman WK, McDougle CJ, Price LH. The role of serotonin and dopamine in the pathophysiology of obsessive compulsive disorder. *Int Clin Psychopharmacol* 1992;7(suppl 1): 35-38
 5. McDougle CJ. Update on pharmacologic management of OCD: agents and augmentation. *J Clin Psychiatry* 1997;58(suppl 12):11-17
 6. Goodman WK, McDougle CJ, Price LH, et al. Beyond the serotonin hypothesis: a role for dopamine in some forms of obsessive compulsive disorder? *J Clin Psychiatry* 1990;51(8, suppl):36-43
 7. Campbell KM, de Lecea L, Severynse DM, et al. OCD-like behaviors caused by a neuropotentiating transgene targeted to cortical and limbic D1+ neurons. *J Neurosci* 1999;19:5044-5053
 8. Campbell KM, McGrath MJ, Burton FH. Behavioral effects of cocaine on a transgenic mouse model of cortical-limbic compulsion. *Brain Res* 1999;833:216-224
 9. Campbell KM, McGrath MJ, Burton FH. Differential response of cortical-limbic neuropotentiated compulsive mice to dopamine D1 and D2 receptor antagonists. *Eur J Pharmacol* 1999;371:103-111
 10. McGrath MJ, Campbell KM, Parks CR, et al. Glutamatergic drugs exacerbate symptomatic behavior in a transgenic model of comorbid Tourette's syndrome and obsessive-compulsive disorder. *Brain Res* 2000;877:23-30
 11. Campbell KM, Veldman MB, McGrath MJ, et al. TS+OCD-like neuropotentiated mice are supersensitive to seizure induction. *Neuroreport* 2000;11:2335-2338
 12. Einat H, Szechtman H. Perseveration without hyperlocomotion in a spontaneous alternation task in rats sensitized to the dopamine agonist quinpirole. *Physiol Behav* 1995;57:55-59
 13. Szechtman H, Sulis W, Eilam D. Quinpirole induces compulsive checking behavior in rats: a potential animal model of obsessive-compulsive disorder (OCD). *Behav Neurosci* 1998;112:1475-1485
 14. Ben Pazi A, Szechtman H, Eilam D. The morphogenesis of motor rituals in rats treated chronically with the dopamine agonist quinpirole. *Behav Neurosci* 2001;115:1301-1317
 15. Szechtman H, Eckert MJ, Tse WS, et al. Compulsive checking behavior of quinpirole-sensitized rats as an animal model of obsessive-compulsive disorder (OCD): form and control. *BMC Neurosci* 2001;2:4
 16. Sullivan RM, Talangbayan H, Einat H, et al. Effects of quinpirole on central dopamine systems in sensitized and non-sensitized rats. *Neuroscience* 1998; 83:781-789
 17. Joel D, Avisar A. Excessive lever pressing following post-training signal attenuation in rats: a possible animal model of obsessive compulsive disorder? *Behav Brain Res* 2001;123:77-87
 18. Joel D, Doljansky J. Selective alleviation of compulsive lever-pressing in rats by D(1), but not D(2), blockade: possible implications for the involvement of D(1) receptors in obsessive-compulsive disorder. *Neuropsychopharmacology* 2003;28:77-85
 19. Thoren P, Asberg M, Bertilsson L, et al. Clomipramine treatment of obsessive-compulsive disorder, 2: biochemical aspects. *Arch Gen Psychiatry* 1980;37:1289-1294
 20. Benkelfat C, Mefford IN, Masters CF, et al. Plasma catecholamines and their metabolites in obsessive-compulsive disorder. *Psychiatry Res* 1991;37: 321-331
 21. Swedo SE, Leonard HL, Kruesi MJ, et al. Cerebrospinal fluid neurochemistry in children and adolescents with obsessive-compulsive disorder. *Arch Gen Psychiatry* 1992;49:29-36
 22. Hollander E, Stein DJ, Saoud JB, et al. Effects of fenfluramine on plasma HVA in OCD [letter]. *Psychiatry Res* 1992;42:185-188
 23. Zahn TP, Kruesi MJ, Swedo SE, et al. Autonomic activity in relation to cerebrospinal fluid neurochemistry in obsessive and disruptive children and adolescents. *Psychophysiology* 1996;33:731-739
 24. Marazziti D, Hollander E, Lensi P, et al. Peripheral markers of serotonin and dopamine function in obsessive-compulsive disorder. *Psychiatry Res* 1992;42: 41-51
 25. Little KY, Zhang L, Desmond T, et al. Striatal dopaminergic abnormalities in human cocaine users. *Am J Psychiatry* 1999;156:238-245
 26. McDougle CJ, Goodman WK, Delgado PL, et al. Pathophysiology of obsessive-compulsive disorder [letter]. *Am J Psychiatry* 1989;146:1350-1351
 27. Rosse RB, Fay-McCarthy M, Collins JP Jr, et al. The relationship between cocaine-induced paranoia and compulsive foraging: a preliminary report. *Addiction* 1994;89:1097-1104
 28. Rosse RB, Fay-McCarthy M, Collins JP Jr, et al. Transient compulsive foraging behavior associated with crack cocaine use. *Am J Psychiatry* 1993;150:155-156
 29. Koizumi HM. Obsessive-compulsive symptoms following stimulants. *Biol Psychiatry* 1985;20:1332-1333
 30. Satel SL, McDougle CJ. Obsessions and compulsions associated with cocaine abuse [letter]. *Am J Psychiatry* 1991;148:947
 31. Rosse RB, McCarthy MF, Alim TN, et al. Saccadic distractibility in cocaine dependent patients: a preliminary laboratory exploration of the cocaine-OCD hypothesis. *Drug Alcohol Depend* 1994;35:25-30
 32. Lemus CZ, Robinson DG, Kronig M, et al. Behavioral responses to a dopaminergic challenge in obsessive-compulsive disorder. *J Anxiety Disord* 1991;5: 369-373
 33. Kouris S. Methylphenidate-induced obsessive-compulsiveness [letter]. *J Am Acad Child Adolesc Psychiatry* 1998;37:135
 34. Kotsopoulos S, Spivak M. Obsessive-compulsive symptoms secondary to methylphenidate treatment [letter]. *Can J Psychiatry* 2001;46:89
 35. Frye PE, Arnold LE. Persistent amphetamine-induced compulsive rituals: response to pyridoxine(B6). *Biol Psychiatry* 1981;16:583-587
 36. Iyo M, Sekine Y, Matsunaga T, et al. Methamphetamine-associated obsessional symptoms and effective risperidone treatment: a case report [letter]. *J Clin Psychiatry* 1999;60:337-338
 37. Joffe RT, Swinson RP. Methylphenidate in primary obsessive-compulsive disorder. *J Clin Psychopharmacol* 1987;7:420-422
 38. Joffe RT, Swinson RP, Levitt AJ. Acute psychostimulant challenge in primary obsessive-compulsive disorder. *J Clin Psychopharmacol* 1991;11:237-241
 39. Insel TR, Hamilton JA, Guttmacher LB, et al. D-amphetamine in obsessive-compulsive disorder. *Psychopharmacology (Berl)* 1983;80:231-235
 40. Pitchot W, Hansenne M, Moreno AG, et al. Growth hormone response to apomorphine in obsessive-compulsive disorder. *J Psychiatry Neurosci* 1996;21:343-345
 41. Brambilla F, Bellodi L, Perna G, et al. Dopamine function in obsessive-compulsive disorder: growth hormone response to apomorphine stimulation. *Biol Psychiatry* 1997;42:889-897
 42. Brambilla F, Perna G, Bussi R, et al. Dopamine function in obsessive compulsive disorder: cortisol response to acute apomorphine stimulation. *Psychoneuroendocrinology* 2000;25:301-310
 43. Longhurst JG, Carpenter LL, Epperson CN, et al. Effects of catecholamine depletion with AMPT (alpha-methyl-para-tyrosine) in obsessive-compulsive disorder. *Biol Psychiatry* 1999;46:573-576
 44. Ceccherini-Nelli A, Guazzelli M. Treatment of refractory OCD with the dopamine agonist bromocriptine [letter]. *J Clin Psychiatry* 1994;55:415-416
 45. Trethowan WH, Scott PA. Chlorpromazine in obsessive-compulsive and allied disorders. *Lancet* 1955;268:781-785
 46. Hussain MZ, Ahad A. Treatment of obsessive-compulsive neurosis [letter]. *Can Med Assoc J* 1970;103:648
 47. O'Regan JB. Treatment of obsessive-compulsive neurosis [letter]. *Can Med Assoc J* 1970;103:650-651
 48. O'Regan JB. Treatment of obsessive-compulsive neurosis with haloperidol [letter]. *Can Med Assoc J* 1970;103:167-168
 49. Altschuler M. Massive doses of trifluoperazine in the treatment of compulsive rituals [letter]. *Am J Psychiatry* 1962;119:367-368
 50. Rivers-Bulkeley N, Hollender MH. Successful treatment of obsessive-compulsive disorder with loxapine [letter]. *Am J Psychiatry* 1982;139:1345-1346
 51. McDougle CJ, Barr LC, Goodman WK, et al. Lack of efficacy of clozapine monotherapy in refractory obsessive-compulsive disorder. *Am J Psychiatry* 1995;152:1812-1814
 52. Lykouras L, Alevizos B, Michalopoulou P, et al. Obsessive-compulsive symptoms induced by atypical antipsychotics: a review of the reported cases. *Prog Neuropsychopharmacol Biol Psychiatry* 2003;27:333-346
 53. McDougle CJ, Goodman WK, Price LH, et al. Neuroleptic addition in fluvoxamine-refractory obsessive-compulsive disorder. *Am J Psychiatry* 1990;147: 652-654
 54. Shapira NA, Ward HE, Mandoki M, et al. A double-blind, placebo-controlled trial of olanzapine addition in fluoxetine-refractory obsessive-compulsive disorder. *Biol Psychiatry* 2004;55:553-555
 55. Sevincok L, Topuz A. Lack of efficacy of low doses of quetiapine addition in refractory obsessive-compulsive disorder. *J Clin Psychopharmacol* 2003;23: 448-450
 56. Metin O, Yazici K, Tot S, et al. Amisulpiride augmentation in treatment resistant obsessive-compulsive disorder: an open trial. *Hum Psychopharmacol* 2003;18:463-467
 57. Ramasubbu R. Antiobsessional effect of risperidone add-on treatment in serotonin reuptake inhibitor-refractory obsessive-compulsive disorder may be dose-dependent [letter]. *Arch Gen Psychiatry* 2002;59:472-473
 58. Zhang W, Bymaster FP. The in vivo effects of olanzapine and other antipsychotic agents on receptor occupancy and antagonism of dopamine D1, D2, D3, 5HT2A and muscarinic receptors. *Psychopharmacology (Berl)* 1999;141: 267-278

59. Zhang W, Perry KW, Wong DT, et al. Synergistic effects of olanzapine and other antipsychotic agents in combination with fluoxetine on norepinephrine and dopamine release in rat prefrontal cortex. *Neuropsychopharmacology* 2000;23:250–262
60. Denys D, Klompmakers AA, Westenberg HG. Synergistic dopamine increase in the rat prefrontal cortex with the combination of quetiapine and fluvoxamine. *Psychopharmacology (Berl)*. In press
61. Sawle GV, Hymas NF, Lees AJ, et al. Obsessional slowness: functional studies with positron emission tomography. *Brain* 1991;114(pt 5):2191–2202
62. van der Wee N, Stevens H, Hardeman H, et al. Enhanced densities of dopamine but not of serotonin transporters in psychotropic-naïve patients with obsessive-compulsive disorder [abstract]. *J Nucl Med* 2001;42:238P
63. Kim CH, Koo MS, Cheon KA, et al. Dopamine transporter density of basal ganglia assessed with [(123)I]IPT SPET in obsessive-compulsive disorder. *Eur J Nucl Med Mol Imaging* 2003;30:1637–1643
64. Pogarell O, Hamann C, Popperl G, et al. Elevated brain serotonin transporter availability in patients with obsessive-compulsive disorder. *Biol Psychiatry* 2003;54:1406–1413
65. Denys D, van der Wee N, Janssen J, et al. Low level of dopaminergic D2 receptor binding in obsessive-compulsive disorder. *Biol Psychiatry* 2004;55:1041–1045
66. Frisch A, Michaelovsky E, Rockah R, et al. Association between obsessive-compulsive disorder and polymorphisms of genes encoding components of the serotonergic and dopaminergic pathways. *Eur Neuropsychopharmacol* 2000;10:205–209
67. Hemmings SM, Kinnear CJ, Niehaus DJ, et al. Investigating the role of dopaminergic and serotonergic candidate genes in obsessive-compulsive disorder. *Eur Neuropsychopharmacol* 2003;13:93–98
68. Catalano M, Sciuto G, Di Bella D, et al. Lack of association between obsessive-compulsive disorder and the dopamine D3 receptor gene: some preliminary considerations. *Am J Med Genet* 1994;54:253–255
69. Nicolini H, Cruz C, Camarena B, et al. DRD2, DRD3 and 5HT2A receptor genes polymorphisms in obsessive-compulsive disorder. *Mol Psychiatry* 1996;1:461–465
70. Denys D, van Nieuwerburgh P, Deforce D, et al. An association between dopamine system genes and obsessive-compulsive disorder. In: Denys D. *On Certainty*. Utrecht, The Netherlands: UMC Utrecht; 2004:163–171
71. Novelli E, Nobile M, Diaferia G, et al. A molecular investigation suggests no relationship between obsessive-compulsive disorder and the dopamine D2 receptor. *Neuropsychobiology* 1994;29:61–63
72. Billett EA, Richter MA, Sam F, et al. Investigation of dopamine system genes in obsessive-compulsive disorder. *Psychiatr Genet* 1998;8:163–169
73. Cruz C, Camarena B, King N, et al. Increased prevalence of the seven-repeat variant of the dopamine D4 receptor gene in patients with obsessive-compulsive disorder with tics. *Neurosci Lett* 1997;231:1–4
74. Millet B, Chabane N, Delorme R, et al. Association between the dopamine receptor D4 (DRD4) gene and obsessive-compulsive disorder. *Am J Med Genet* 2003;116:55–59
75. Di Bella D, Catalano M, Cichon S, et al. Association study of a null mutation in the dopamine D4 receptor gene in Italian patients with obsessive-compulsive disorder, bipolar mood disorder and schizophrenia. *Psychiatr Genet* 1996;6:119–121
76. Karayiorgou M, Altemus M, Galke BL, et al. Genotype determining low catechol-O-methyltransferase activity as a risk factor for obsessive-compulsive disorder. *Proc Natl Acad Sci U S A* 1997;94:4572–4575
77. Karayiorgou M, Sobin C, Blundell ML, et al. Family-based association studies support a sexually dimorphic effect of COMT and MAOA on genetic susceptibility to obsessive-compulsive disorder. *Biol Psychiatry* 1999;45:1178–1189
78. Alsobrook JP, Zohar AH, Leboyer M, et al. Association between the COMT locus and obsessive-compulsive disorder in females but not males. *Am J Med Genet* 2002;114:116–120
79. Niehaus DJ, Kinnear CJ, Corfield VA, et al. Association between a catechol-o-methyltransferase polymorphism and obsessive-compulsive disorder in the Afrikaner population. *J Affect Disord* 2001;65:61–65
80. Schindler KM, Richter MA, Kennedy JL, et al. Association between homozygosity at the COMT gene locus and obsessive compulsive disorder. *Am J Med Genet* 2000;96:721–724
81. Ohara K, Nagai M, Suzuki Y, et al. No association between anxiety disorders and catechol-O-methyltransferase polymorphism. *Psychiatry Res* 1998;80:145–148
82. Erdal ME, Tot S, Yazici K, et al. Lack of association of catechol-O-methyltransferase gene polymorphism in obsessive-compulsive disorder. *Depress Anxiety* 2003;18:41–45
83. Azzam A, Mathews C, Reus V. A meta-analysis of the association between the catecholamine-O-methyl-transferase gene and obsessive-compulsive disorder. *Am J Med Genet* 2002;114:834–835
84. Saxena S, Brody AL, Schwartz JM, et al. Neuroimaging and frontal-subcortical circuitry in obsessive-compulsive disorder. *Br J Psychiatry Suppl* 1998:26–37
85. Jenike MA, Baer L, Minichiello WE. *Obsessive-Compulsive Disorders. Practical Management*, 3rd Edition. St. Louis, MO: Mosby; 1998
86. LeDoux JE. *The Emotional Brain*. New York, NY: Simon & Schuster; 1996
87. Rosenkranz JA, Grace AA. Dopamine-mediated modulation of odour-evoked amygdala potentials during pavlovian conditioning. *Nature* 2002;417:282–287
88. Holden C. ‘Behavioral’ addictions: do they exist? *Science* 2001;294:980–982
89. Fiorillo CD, Tobler PN, Schultz W. Discrete coding of reward probability and uncertainty by dopamine neurons. *Science* 2003;299:1898–1902
90. McDougle CJ, Fleischmann RL, Epperson CN, et al. Risperidone addition in fluvoxamine-refractory obsessive-compulsive disorder: three cases. *J Clin Psychiatry* 1995;56:526–528
91. Saxena S, Wang D, Bystritsky A, et al. Risperidone augmentation of SRI treatment for refractory obsessive-compulsive disorder. *J Clin Psychiatry* 1996;57:303–306
92. Stein DJ, Bouwer C, Hawkrige S, et al. Risperidone augmentation of serotonin reuptake inhibitors in obsessive-compulsive and related disorders. *J Clin Psychiatry* 1997;58:119–122
93. Pfanner C, Marazziti D, Dell’Osso L, et al. Risperidone augmentation in refractory obsessive-compulsive disorder: an open-label study. *Int Clin Psychopharmacol* 2000;15:297–301
94. McDougle CJ, Epperson CN, Pelton GH, et al. A double-blind, placebo-controlled study of risperidone addition in serotonin reuptake inhibitor-refractory obsessive-compulsive disorder. *Arch Gen Psychiatry* 2000;57:794–801
95. Baxter LR, Li X, Jackson WT, et al. Adjunctive risperidone in the treatment of SSRI-refractory obsessive compulsive disorder. Presented at the annual meeting of the Society of Biological Psychiatry; May 16–18, 2002; Philadelphia, Pa
96. Hollander E, Rossi NB, Sood E, et al. Risperidone augmentation in treatment-resistant obsessive-compulsive disorder: a double-blind, placebo-controlled study. *Int J Neuropsychopharmacol* 2003;6:397–401
97. Weiss EL, Potenza MN, McDougle CJ, et al. Olanzapine addition in obsessive-compulsive disorder refractory to selective serotonin reuptake inhibitors: an open-label case series. *J Clin Psychiatry* 1999;60:524–527
98. Koran LM, Ringold AL, Elliott MA. Olanzapine augmentation for treatment-resistant obsessive-compulsive disorder. *J Clin Psychiatry* 2000;61:514–517
99. Bogetto F, Bellino S, Vaschetto P, et al. Olanzapine augmentation of fluvoxamine-refractory obsessive-compulsive disorder (OCD): a 12-week open trial. *Psychiatry Res* 2000;96:91–98
100. Francobandiera G. Olanzapine augmentation of serotonin uptake inhibitors in obsessive-compulsive disorder: an open study. *Can J Psychiatry* 2001;46:356–358
101. Crocq MA, Leclercq P, Guillon MS, et al. Open-label olanzapine in obsessive-compulsive disorder refractory to antidepressant treatment. *Eur Psychiatry* 2002;17:296–297
102. D’Amico G, Cedro C, Muscatello MR, et al. Olanzapine augmentation of paroxetine-refractory obsessive-compulsive disorder. *Prog Neuropsychopharmacol Biol Psychiatry* 2003;27:619–623
103. Bystritsky A, Ackerman DL, Rosen RM, et al. Augmentation of serotonin reuptake inhibitors in refractory obsessive-compulsive disorder using adjunctive olanzapine: a placebo-controlled trial. *J Clin Psychiatry* 2004;65:565–568
104. Mohr N, Vythilingum B, Emsley RA, et al. Quetiapine augmentation of serotonin reuptake inhibitors in obsessive-compulsive disorder. *Int Clin Psychopharmacol* 2002;17:37–40
105. Atmaca M, Kuloglu M, Tezcan E, et al. Quetiapine augmentation in patients with treatment resistant obsessive-compulsive disorder: a single-blind, placebo-controlled study. *Int Clin Psychopharmacol* 2002;17:115–119
106. Denys D, van Megan H, Westenberg H. Quetiapine addition to serotonin reuptake inhibitor treatment in patients with treatment-refractory obsessive-compulsive disorder: an open-label study. *J Clin Psychiatry* 2002;63:700–703
107. Kinnear C, Niehaus DJ, Seedat S, et al. Obsessive-compulsive disorder and a novel polymorphism adjacent to the oestrogen response element (ERE 6) upstream from the COMT gene. *Psychiatr Genet* 2001;11:85–87
108. Meira-Lima I, Shavitt RG, Miguita K, et al. Association analysis of the catechol-o-methyltransferase (COMT), serotonin transporter (5-HTT) and serotonin 2A receptor (5HT2A) gene polymorphisms with obsessive-compulsive disorder. *Genes Brain Behav* 2004;3:75–79
109. Denys D, De Geus F, van Megan HJ, et al. A double-blind, randomized, placebo-controlled trial of quetiapine addition in patients with obsessive-compulsive disorder refractory to serotonin reuptake inhibitors. *J Clin Psychiatry* 2004;65:1040–1048