

# Fundamentals of Attention-Deficit/Hyperactivity Disorder: Circuits and Pathways

Amy F. T. Arnsten, Ph.D.

Neuropsychological and imaging studies have shown that attention-deficit/hyperactivity disorder (ADHD) is associated with alterations in prefrontal cortex (PFC) and its connections to striatum and cerebellum. Research in animals, in combination with observations of patients with cortical lesions, has shown that the PFC is critical for the regulation of behavior, attention, and affect using representational knowledge. The PFC is important for sustaining attention over a delay, inhibiting distraction, and dividing attention, while more posterior cortical areas are essential for perception and the allocation of attentional resources. The PFC in the right hemisphere is especially important for behavioral inhibition. Lesions to the PFC produce a profile of distractibility, forgetfulness, impulsivity, poor planning, and locomotor hyperactivity. The PFC is very sensitive to its neurochemical environment, and optimal levels of norepinephrine and dopamine are needed for proper PFC control of behavior and attention. Recent electrophysiologic studies in animals suggest that norepinephrine enhances “signals” through postsynaptic  $\alpha_{2A}$ -adrenoceptors in PFC, while dopamine decreases “noise” through modest levels of  $D_1$ -receptor stimulation. Blockade of  $\alpha_2$ -receptors in the monkey PFC re-creates the symptoms of ADHD, resulting in impaired working memory, increased impulsivity, and locomotor hyperactivity. Genetic alterations in catecholamine pathways may contribute to dysregulation of PFC circuits in this disorder. Stimulant medications may have some of their therapeutic effects by increasing endogenous stimulation of  $\alpha_{2A}$ -adrenoceptors and dopamine  $D_1$ -receptors in the PFC, optimizing PFC regulation of behavior and attention.

(*J Clin Psychiatry* 2006;67[suppl 8]:7–12)

**A**ttention-deficit/hyperactivity disorder (ADHD) is characterized by inattention, impaired behavioral inhibition, and increased motor activity. *Inattention* is a nonprecise term and thus can lead to confusion in diagnosis. For many patients, *inattention* refers to increased distractibility, poor sustained attention, and increased susceptibility to interference, i.e., impaired regulation of attention. However, the term *inattention* can also be applied to symptoms resembling perceptual neglect, i.e., the inability to allocate sufficient attentional resources to stimuli for adequate perception. Recent understanding of the higher cortical circuits underlying attention and behavioral regulation provides important insights to navigate these complexities and helps to elucidate the likely neural bases underlying the symptoms of ADHD. These findings complement those arising from imaging studies of ADHD that have found volumetric differences in prefrontal cortex

(PFC), cerebellum, and possibly striatum (reviewed by Giedd et al.<sup>1</sup>). Furthermore, our increasing understanding of catecholamine actions on these circuits provides insight into the mechanism of action of medications used to treat this disorder. These data also help us to understand how genetic changes in catecholamine pathways could result in ADHD symptomatology. This article briefly reviews the anatomy and function of the higher association cortices relevant to ADHD, the connections of these cortical areas to the basal ganglia and cerebellum, and the powerful modulation of these circuits by dopamine and norepinephrine.

## THE FUNCTIONAL CONTRIBUTIONS OF HIGHER ASSOCIATION CORTICES

The association cortices make distinct contributions to our attentional experience (summarized in Figure 1): the higher-order sensory cortices, such as the inferior temporal cortex, process sensory features and can focus resources on a particular detail, e.g., the color red; the posterior parietal association cortex allocates attentional resources, allowing us to orient attention in time and space; and the PFC regulates attention, inhibiting processing of irrelevant stimuli, sustaining attention over long delays, and dividing and coordinating attention. These cortical areas are intricately interconnected,<sup>2</sup> creating both feedforward and feedback

---


*From the Department of Neurobiology, Yale University School of Medicine, New Haven, Conn.*

*This article is derived from the planning teleconference “New Developments in the Treatment of Attention-Deficit/Hyperactivity Disorder,” which was held May 16, 2005, and supported by an educational grant from Cephalon, Inc.*

*This work was supported by grant R37 AG060636 from the U.S. Public Health Service, Bethesda, Md.*

*Corresponding author and reprints: Amy F. T. Arnsten, Ph.D., Department of Neurobiology, Yale University School of Medicine, New Haven, CT 06510 (e-mail: amy.arnsten@yale.edu).*

**Figure 1. Brain Regions Involved in the Regulation of Attention in the Primate Brain as Shown on the Lateral Surface<sup>a</sup>**



<sup>a</sup>The basal ganglia circuits are not evident from this view. The basal ganglia are thought to contribute to the automatic planning, selection, initiation, and execution of complex movements and thoughts.

loops<sup>3</sup> that work together to provide a unified attentional experience. The PFC plays a similar role in regulating movement, inhibiting inappropriate behavioral responses. The following is a brief review of these cortical systems.

**Inferior Temporal Cortex**

The inferior temporal cortex is specialized for processing visual features, determining “what” things are based on their color and shape. (In contrast, the superior temporal cortex processes auditory information, performing both featural and spatial analysis. This work has proceeded more recently<sup>4</sup> and thus will not be discussed in this review). Processing of a visual stimulus by inferior temporal neurons can be either diminished or enhanced, depending upon sensory conditions and internal directions (reviewed in Desimone<sup>5</sup> and Kastner et al.<sup>6</sup>). The activity of these neurons is captured by salient stimuli (e.g., high contrasts), but repeated experience with the same visual stimulus leads to decreased responding.<sup>5</sup> This decrease may account for the boredom of repetition, e.g., in a school setting. Processing of visual stimuli is also diminished by interference from nearby stimuli in the same visual field. Both of these suppressive effects result from intrinsic properties of inferior temporal neurons and can be prevented by inputs from the PFC or parietal association cortex. These “top-down” projections thus allow for directed selective attention of visual feature processing.

**Posterior Parietal Association Cortex**

The parietal association cortex plays a critical role in “paying attention.”<sup>7</sup> This cortex is specialized for analyzing movement and spatial relationships, for analyzing

quantity and constructing spatial maps, and for orienting attention in time and space.<sup>8</sup> Lesions to the right posterior parietal association cortex can result in contralateral neglect: the loss of perception for the left side of visual space. This cortex is therefore critical to conscious attention. Recordings from parietal neurons in monkeys are consistent with a role in the allocation of attention, including the covert movement of attention.<sup>9</sup> Neurons in area 7a also appear to create world-referenced maps of visual space,<sup>10</sup> and they project this information to the PFC, which uses this information to guide behavior during spatial working memory tasks.

**Prefrontal Cortex**

The PFC has particular relevance to ADHD, as imaging studies indicate that ADHD patients often have smaller PFC volume, particularly on the right side.<sup>11–13</sup> The PFC uses representational knowledge, i.e., working memory, to guide overt responses (movement) as well as covert responses (attention), allowing us to inhibit inappropriate behaviors and to gate the processing of irrelevant stimuli (reviewed in references 14–17). Patients with PFC lesions are easily distracted, have poor concentration and organization, are more vulnerable to disruption from proactive interference, and can be impulsive, especially when the lesions involve the right hemisphere.<sup>18</sup> PFC lesions impair the ability to sustain attention, particularly over a delay, and reduce the ability to gate sensory input.<sup>19</sup> PFC lesions impair divided and focused attention, and these attentional deficits have been associated with lesions in the left, superior PFC.<sup>15</sup> PFC lesions similarly impair attentional function in monkeys and rats, rendering animals more vulnerable to distraction or other types of interference and impairing attentional regulation on set-shifting tasks.<sup>20,21</sup>

Electrophysiologic studies in monkeys have shown that PFC neurons are able to hold modality-specific information “on-line” over a delay and use this information to guide behavior in the absence of environmental information. Importantly, PFC neurons can maintain delay-related firing in the presence of distracting stimuli, protecting performance from interference.<sup>22</sup> Delay-related firing also underlies behavioral inhibition, i.e., the ability to suppress a prepotent response.<sup>23</sup> Therefore, the cellular basis of many executive functions can be studied in monkeys performing higher cognitive tasks.

**CORTICAL PROJECTIONS TO BASAL GANGLIA AND CEREBELLUM**

The association cortices project down to both the basal ganglia and cerebellum in a series of parallel, closed-loop circuits.<sup>24</sup> The PFC and parietal and temporal association cortices all project to the caudate nucleus as part of the “cognitive circuit” through the basal ganglia, which in turn projects back to the PFC and premotor cortices. Simi-

larly, the PFC and parietal association cortices project to the cerebellum by way of the pontine nuclei, and the cerebellum in turn projects back to the association cortices by way of dentate projections to thalamus. Although the basal ganglia and cerebellum have long been known to be important for the regulation of movement, their role in higher cognitive function is just beginning to be researched. If these structures influence cognition in a manner similar to their influences on movement, the basal ganglia may be important for the planning, selection, initiation, and execution of thoughts, while the cerebellum may serve as a "biological gyroscope," correcting cognitive function on a faster timescale. Recent data suggest that the cerebellar circuits may influence basal ganglia circuits via projections through the intralaminar thalamic nuclei (P. Strick, Ph.D.; electronic communication; August 2005). Parts of the cerebellum have been found to be consistently smaller in children with ADHD.<sup>25</sup>

Interestingly, the basal ganglia are powerfully modulated by dopamine, while the cerebellum is heavily innervated by norepinephrine. Dopamine promotes thalamocortical stimulation of movement and thought via D<sub>1</sub>-receptor activation of the direct pathways and D<sub>2</sub> inhibition of the indirect pathways in basal ganglia (for example, see Steiner and Gerfen<sup>26</sup>). Norepinephrine modulates cerebellar processing through  $\beta$ -receptor mechanisms that activate cyclic adenosine monophosphate (cAMP)/protein kinase A (PKA) intracellular signaling (for example, see Cartford et al.<sup>27</sup>). Therefore, genetic alterations that affect catecholamine actions may alter basal ganglia and cerebellar function. In addition to these dense subcortical projections, both dopamine and norepinephrine innervate the association cortices in primates,<sup>28,29</sup> and this will be the focus of the current review.

### MODULATION OF CORTICAL CIRCUITS

There are multiple arousal systems that project to the cortical mantle. Although the present article focuses on dopamine and norepinephrine, the reader should remember that many more molecules perform related functions, e.g., acetylcholine, serotonin, and orexin. Research on the modulatory influences of dopamine and norepinephrine on parietal and temporal cortical function is limited, although existing research suggests that norepinephrine may enhance "signal/noise" processing in these areas via the actions of  $\beta$ - and  $\alpha_1$ -adrenoceptors and impair the cognitive operations of these brain regions via  $\alpha_2$ -adrenoceptors.<sup>30-32</sup> There has been a much greater tradition of studying catecholamine influences on PFC, given the landmark findings of Brozoski et al.,<sup>33</sup> who showed that catecholamine depletion in PFC is as destructive as ablation of the tissue itself. The following is a brief summary of this field (for more detailed reviews, see Arnsten and Robbins<sup>34</sup> and Arnsten and Li<sup>35</sup>).

### Dopamine

Dopamine acts at both the D<sub>1</sub> (D<sub>1</sub> and D<sub>5</sub>) and D<sub>2</sub> (D<sub>2</sub>, D<sub>3</sub>, and D<sub>4</sub>) receptor families. It should be noted that norepinephrine has very high affinity for D<sub>4</sub>-receptors<sup>36</sup> and thus should really be considered a catecholamine receptor rather than a dopamine receptor. Currently, there are no pharmacologic agents that distinguish D<sub>1</sub>- from D<sub>5</sub>-receptors; thus, these receptors are discussed as 1 entity. Finally, as there is little information on D<sub>3</sub>-receptor actions in PFC at this time, this receptor is not reviewed.

**D<sub>1</sub>-receptor actions.** The PFC is rich in D<sub>1</sub>-receptors, with a bilaminar distribution found in both upper and lower layers.<sup>37</sup> The D<sub>1</sub> subtype is especially focused on spines, while the D<sub>5</sub> subtype is found more commonly on shafts of pyramidal neurons.<sup>38</sup> Stimulation of this family of receptors in the PFC produces an inverted "U"-shaped dose-response influence on the working memory and attention regulation processes of the PFC.<sup>39,40</sup> While modest levels of D<sub>1</sub>-receptor stimulation are essential to PFC function, high levels of dopamine release, e.g., during exposure to stress, impair working memory. Therefore, high doses of D<sub>1</sub>-agonist impair working memory performance. This inverted U has been observed in mice,<sup>41</sup> rats,<sup>42</sup> monkeys,<sup>43</sup> and humans.<sup>44</sup> A similar inverted "U" has been described at the cellular level (S. Vijayraghavan, B.S.; M. Wang, Ph.D.; S. G. Birnbaum, Ph.D., et al.; manuscript submitted), where moderate levels of D<sub>1</sub>-receptor stimulation suppress neuronal processing of irrelevant information (i.e., reduce "noise"), sharpening tuning and rendering PFC firing more selective. In contrast, high levels of D<sub>1</sub>-receptor stimulation reduce the relevant signals as well as noise, thus producing a nonspecific inhibition of cell firing. These effects most likely occur through cAMP/PKA intracellular signaling (S. Vijayraghavan, B.S.; M. Wang, Ph.D.; S. G. Birnbaum, Ph.D., et al.; manuscript submitted).

Human experiments also note an inverted "U," with the more D<sub>1</sub>-like compounds being most effective in improving working memory.<sup>45</sup> Interestingly, genetic studies in humans indicate similar results.<sup>44</sup> Some people have a substitution (methionine for valine) in the enzyme catechol O-methyltransferase that breaks down dopamine and norepinephrine. The methionine substitution results in weaker enzyme activity and thus more dopamine. Under basal conditions, subjects with this substitution have better working memory and more efficient PFC activation than those with the native enzyme. However, following amphetamine and/or stress exposure, those with the methionine substitutions become markedly impaired (presumably due to excessive dopamine stimulation), while those with the native enzyme improve (presumably due to more optimal dopamine levels).<sup>44</sup>

**D<sub>2</sub>-receptor actions.** There has been far less research on the influence of D<sub>2</sub>-receptor stimulation on PFC function. D<sub>2</sub>-receptors are concentrated on neurons in layer 5

(the output layer that projects to striatum) and overall show lower levels of binding than the D<sub>1</sub>-receptor family.<sup>37</sup> Initial studies showed that blockade of D<sub>2</sub>-receptors in the PFC of monkeys performing a working memory task had no effect on performance.<sup>46</sup> However, there was no room for improvement in this task, and studies in rats suggest that excessive D<sub>2</sub>-receptor stimulation impairs working memory abilities.<sup>47</sup> Recent electrophysiologic studies have shown that D<sub>2</sub>-receptor stimulation increases the response-related firing of PFC neurons in monkeys performing a working memory task,<sup>48</sup> consistent with the localization of these receptors on cells projecting to areas guiding movement. The response-related firing may be a form of corollary discharge, informing the brain that a response has taken place. Given the potential role of impaired corollary discharge in hallucinations, and the role of D<sub>2</sub> blockade in antipsychotic efficacy, these findings may have special relevance to schizophrenia.

**D<sub>4</sub>-receptor actions.** D<sub>4</sub>-receptors are concentrated on  $\gamma$ -aminobutyric acid (GABA)-ergic interneurons<sup>49</sup> and appear to inhibit GABA transmission via G<sub>i</sub>-mediated reduction of cAMP signaling.<sup>50</sup> Weaker D<sub>4</sub>-receptor actions thus lead to excessive GABA transmission and suppression of pyramidal cell firing (Wang et al.<sup>50</sup> and M. Wang, Ph.D.; A.F.T.A.; manuscript in preparation). ADHD is associated with the increased incidence of the 7-repeat polymorphism of the D<sub>4</sub>-receptor, which weakens D<sub>4</sub>-receptor efficacy.<sup>51,52</sup> Therefore, the basic physiology in animals would suggest that subjects with this polymorphism would have insufficient D<sub>4</sub> inhibition of GABA and thus insufficient PFC pyramidal cell firing. Stimulant medication may increase endogenous dopamine (and norepinephrine) stimulation of D<sub>4</sub>-receptors, thus normalizing GABAergic inhibition. However, there is also some evidence from basic physiologic studies indicating that D<sub>4</sub>-receptors can inhibit pyramidal cells,<sup>50</sup> so the actions of these receptors are not entirely straightforward.

### Norepinephrine

Norepinephrine acts at  $\alpha_1$ - and  $\alpha_2$ -adrenoceptors and  $\beta_1$ -,  $\beta_2$ -, and  $\beta_3$ -adrenoceptors. Research to date indicates that it has distinct actions at these receptors, improving PFC function through actions at  $\alpha_{2A}$ -receptors and impairing working memory through actions at  $\alpha_1$ - and  $\beta_1$ -receptors.

**$\alpha_{2A}$ -Adrenoceptor actions.** Norepinephrine improves PFC function through actions at postsynaptic<sup>53,54</sup>  $\alpha_{2A}$ -receptors<sup>55</sup> at both the cognitive and cellular levels (reviewed in Arnsten and Li<sup>55</sup>). The  $\alpha_{2A}$ -agonist guanfacine improves working memory, attention regulation, behavioral inhibition, and/or planning in rodents,<sup>55,56</sup> monkeys,<sup>57-60</sup> and humans.<sup>61</sup> These enhancing effects are most likely mediated through G<sub>i</sub>-mediated suppression of cAMP intracellular signaling.<sup>62</sup> Electrophysiologic studies have shown that  $\alpha_2$ -receptor stimulation increases delay-related fir-

ing,<sup>63</sup> the cellular measure of working memory and behavioral inhibition (see Prefrontal Cortex). This increase is especially observed for the preferred spatial direction, indicating that  $\alpha_{2A}$ -receptor stimulation increases "signals" in PFC (M. Wang, Ph.D.; A.F.T.A.; manuscript in preparation). Conversely, blocking  $\alpha_2$ -receptors in monkey PFC with yohimbine markedly reduces delay-related cell firing<sup>63</sup> and impairs working memory<sup>64</sup> as well as impulse control.<sup>65</sup> Yohimbine infusions into PFC have also been shown to induce locomotor hyperactivity.<sup>66</sup> Therefore, insufficient  $\alpha_2$ -receptor stimulation in monkey PFC can re-create the profile of ADHD. In this regard, it is of interest that ADHD has been associated with genetic alterations in dopamine  $\beta$ -hydroxylase (DBH), the enzyme that synthesizes norepinephrine. It is possible that weaker DBH would lead to insufficient endogenous stimulation of  $\alpha_{2A}$ -receptors in PFC, resulting in a profile similar to that of yohimbine-treated monkeys. Stimulant or  $\alpha_{2A}$ -agonist medications might correct this condition. Guanfacine is currently used for treating ADHD,<sup>67-69</sup> especially in patients who cannot take stimulants, e.g., those with tics or risk of drug abuse.

**$\alpha_1$ -Adrenoceptor actions.** In contrast to  $\alpha_2$ -receptor mechanisms, high levels of norepinephrine release (e.g., during stress) impair PFC function through actions at  $\alpha_1$ -receptors coupled to protein kinase C (PKC) intracellular signaling.<sup>70</sup> Thus, agonists such as phenylephrine (similar to pseudoephedrine) impair working memory when infused into the PFC in rats<sup>71</sup> and monkeys.<sup>59</sup> Similar effects are observed at the cellular level, where  $\alpha_1$ -receptor stimulation suppresses delay-related neuronal firing.<sup>70</sup> Conversely,  $\alpha_1$ -antagonists such as urapidil and prazosin protect PFC cognitive abilities, preventing stress-induced PFC impairment.<sup>72,73</sup> The PKC inhibitor chelerythrine also protects PFC function at the behavioral and cellular levels.<sup>70</sup> On the basis of this research in animals, prazosin is being successfully used to treat patients with posttraumatic stress disorder.<sup>74</sup> Interestingly, most effective antipsychotic medications, including the "atypical" neuroleptics, have potent  $\alpha_1$ -blocking properties, and overactivity of the PKC signaling pathway has been associated with mania<sup>75</sup> and possibly schizophrenia.<sup>76</sup> These mechanisms may be particularly relevant in children who have symptoms resembling ADHD that worsen with stimulant treatment and who are found to have childhood bipolar disorder.<sup>77</sup>

**$\beta_1$ -Adrenoceptor actions.** Recent studies suggest that stimulation of  $\beta_1$ -adrenoceptors impairs PFC function.<sup>78</sup> Therefore, systemic or local application of the  $\beta_1$ -antagonist betaxolol improved working memory in rats and monkeys. However, this treatment appeared to be associated with serious pancreatic side effects and thus may not be appropriate for human use.

### Summary

The PFC appears to thrive under conditions of moderate catecholamine release, when norepinephrine  $\alpha_{2A}$ -receptor



stimulation increases “signals” and optimal dopamine D<sub>1</sub>-receptor stimulation decreases “noise.” In contrast, PFC working memory functions are impaired under conditions of high catecholamine release that engage  $\alpha_1$ - and  $\beta$ -receptors and excessive D<sub>1</sub>-receptor stimulation. These neurochemical needs are opposite to those of posterior cortical and subcortical structures.<sup>79</sup> Therefore, catecholamines may act as a chemical switch, turning on PFC during normal waking and turning it off during drowsiness or stress. In contrast, high levels of catecholamines may turn on more primitive brain structures such as the amygdala for more automatic control of behavior under conditions of danger.<sup>80</sup>

### RELEVANCE TO ADHD MEDICATIONS

Drugs such as amphetamine and methylphenidate act to enhance the release and/or inhibit the reuptake of both dopamine and norepinephrine. Methylphenidate can improve PFC working memory function and enhance the efficiency of PFC activation in healthy adult humans<sup>81,82</sup> as well as in patients with ADHD.<sup>83</sup> Recent studies in rats show that low oral doses of methylphenidate that produce plasma drug levels similar to therapeutic doses in humans<sup>84</sup> also improve PFC function.<sup>85</sup> These enhancing effects in rodents depend on dopamine D<sub>1</sub>- and norepinephrine  $\alpha_2$ -receptor stimulation.<sup>85</sup> It is likely that both dopamine and norepinephrine actions contribute to the therapeutic effects of stimulants in patients with ADHD. In contrast, excessive doses of stimulant medication may produce cognitive inflexibility through  $\alpha_1$ -,  $\beta$ -, and high D<sub>1</sub>-receptor stimulation.

In summary, catecholamines have powerful influences on the brain circuits that appear altered in ADHD. Medications that optimize catecholamine transmission may normalize the function of these circuits and ameliorate ADHD symptomatology.

*Drug names:* amphetamine (Adderall and others), betaxolol (Kerlone and others), guanfacine (Tenex and others), methylphenidate (Ritalin, Concerta, and others), prazosin (Minipress and others), pseudoephedrine (Sudafed, Efidac, and others).

*Disclosure of off-label usage:* The author has determined that, to the best of her knowledge, guanfacine is not approved by the U.S. Food and Drug Administration for the treatment of attention-deficit/hyperactivity disorder, and prazosin is not approved for the treatment of posttraumatic stress disorder.

### REFERENCES

- Giedd JN, Blumenthal J, Molloy E, et al. Brain imaging of attention deficit/hyperactivity disorder. *Ann N Y Acad Sci* 2001;931:33–49
- Goldman-Rakic PS. Circuitry of the primate prefrontal cortex and the regulation of behavior by representational memory. In: Plum F, ed. *The Nervous System. Higher Functions of the Brain*. Bethesda, Md: American Physiological Society; 1987:373–417. *Handbook of Physiology*; vol 5
- Barbas H, Medalla M, Alade O, et al. Relationship of prefrontal connections to inhibitory systems in superior temporal areas in the rhesus monkey. *Cereb Cortex* 2005;15:1356–1370
- Rauschecker JP, Tian B. Mechanisms and streams for processing of “what” and “where” in auditory cortex. *Proc Natl Acad Sci U S A* 2000;97:11800–11806
- Desimone R. Neural mechanisms for visual memory and their role in attention. *Proc Natl Acad Sci U S A* 1996;93:13494–13499
- Kastner S, De Weerd P, Desimone R, et al. Mechanisms of directed attention in the human extrastriate cortex as revealed by functional MRI. *Science* 1998;282:108–111
- Mesulam M-M. *Principles of Behavioral and Cognitive Neurology*. New York, NY: Oxford University Press; 2000
- Coull JT, Nobre AC. Where and when to pay attention: the neural systems for directing attention to spatial locations and to time intervals as revealed by both PET and fMRI. *J Neurosci* 1998;18:7426–7435
- Crowe DA, Chafee MV, Averbeck BB, et al. Neural activity in primate parietal area 7a related to spatial analysis of visual mazes. *Cereb Cortex* 2004;14:23–34
- Snyder LH, Grieve KL, Brotchie P, et al. Separate body- and world-referenced representations of visual space in parietal cortex. *Nature* 1998;394:887–891
- Castellanos FX, Giedd JN, Marsh WL, et al. Quantitative brain magnetic resonance imaging in attention deficit/hyperactivity disorder. *Arch Gen Psychiatry* 1996;53:607–616
- Casey BJ, Castellanos FX, Giedd JN, et al. Implication of right frontostriatal circuitry in response inhibition and attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry* 1997;36:374–383
- Sowell ER, Thompson PM, Welcome SE, et al. Cortical abnormalities in children and adolescents with attention-deficit hyperactivity disorder. *Lancet* 2003;362:1699–1707
- Goldman-Rakic PS. The prefrontal landscape: implications of functional architecture for understanding human mentation and the central executive. *Philos Trans R Soc Lond B Biol Sci* 1996;351:1445–1453
- Godefroy O, Rousseaux M. Divided and focused attention in patients with lesion of the prefrontal cortex. *Brain Cogn* 1996;30:155–174
- Robbins TW. Dissociating executive functions of the prefrontal cortex. *Philos Trans R Soc Lond B Biol Sci* 1996;351:1463–1471
- Knight RT, Staines WR, Swick D, et al. Prefrontal cortex regulates inhibition and excitation in distributed neural networks. *Acta Psychol (Amst)* 1999;101:159–178
- Aron AR, Robbins TW, Poldrack RA. Inhibition and the right inferior frontal cortex. *Trends Cogn Sci* 2004;8:170–177
- Wilkins AJ, Shallice T, McCarthy R. Frontal lesions and sustained attention. *Neuropsychologia* 1987;25:359–365
- Bartus RT, Levere TE. Frontal decortication in rhesus monkeys: a test of the interference hypothesis. *Brain Res* 1977;119:233–248
- Birrell JM, Brown VJ. Medial frontal cortex mediates perceptual attentional set shifting in the rat. *J Neurosci* 2000;20:4320–4324
- Miller EK, Li L, Desimone R. Activity of neurons in anterior inferior temporal cortex during a short-term memory task. *J Neurosci* 1993;13:1460–1478
- Funahashi S, Chafee MV, Goldman-Rakic PS. Prefrontal neuronal activity in rhesus monkeys performing a delayed anti-saccade task. *Nature* 1993;365:753–756
- Dum RP, Li C, Strick PL. Motor and nonmotor domains in the monkey dentate. *Ann N Y Acad Sci* 2002;978:289–301
- Berquin PC, Giedd JN, Jacobsen LK, et al. Cerebellum in attention-deficit hyperactivity disorder: a morphometric MRI study. *Neurology* 1998;50:1087–1093
- Steiner H, Gerfen CR. Role of dynorphin and enkephalin in the regulation of striatal output pathways and behavior. *Exp Brain Res* 1998;123:60–76
- Cartford MC, Samec A, Fister M, et al. Cerebellar norepinephrine modulates learning of delay classical eyeblink conditioning: evidence for post-synaptic signaling via PKA. *Learn Mem* 2004;11:732–737
- Lewis DA, Cambell MJ, Foote SL, et al. The distribution of tyrosine hydroxylase-immunoreactive fibers in primate neocortex is widespread but regionally specific. *J Neurosci* 1987;7:279–290
- Lewis DA, Morrison JH. Noradrenergic innervation of monkey prefrontal cortex: a dopamine-beta-hydroxylase immunohistochemical study. *J Comp Neurol* 1989;282:317–330
- Mouradian RD, Sellar FM, Waterhouse BD. Noradrenergic potentiation of excitatory transmitter action in cerebrocortical slices: evidence of mediation by an  $\alpha_1$ -receptor-linked second messenger pathway. *Brain Res* 1991;546:83–95
- Witte EA, Marrocco RT. Alteration of brain noradrenergic activity in rhesus monkeys affects the alerting component of covert orienting. *Psychopharmacology (Berl)* 1997;132:315–323
- Coull JT, Nobre AC, Frith CD. The noradrenergic  $\alpha_2$  agonist clonidine modulates behavioural and neuroanatomical correlates of human attentional orienting and alerting. *Cereb Cortex* 2001;11:73–84
- Brozoski T, Brown RM, Rosvold HE, et al. Cognitive deficit caused by regional depletion of dopamine in prefrontal cortex of rhesus monkey. *Science* 1979;205:929–931

34. Arnsten AFT, Robbins TW. Neurochemical modulation of prefrontal cortical function in humans and animals. In: Stuss DT, Knight RT, eds. *Principles of Frontal Lobe Function*. New York, NY: Oxford University Press; 2002: 51–84
35. Arnsten AFT, Li B-M. Neurobiology of executive functions: catecholamine influences on prefrontal cortical function. *Biol Psychiatry* 2005;57: 1377–1384
36. Van Tol HHM, Bunzow JR, Guan H-C, et al. Cloning of the gene for a human dopamine D<sub>4</sub> receptor with high affinity for the antipsychotic clozapine. *Nature* 1991;350:610–614
37. Goldman-Rakic PS, Lidow MS, Gallager DW. Overlap of dopaminergic, adrenergic, and serotonergic receptors and complementarity of their subtypes in primate prefrontal cortex. *J Neurosci* 1990;10:2125–2138
38. Smiley JF, Levey AI, Ciliax BJ, et al. D<sub>1</sub> dopamine receptor immunoreactivity in human and monkey cerebral cortex: predominant and extrasynaptic localization in dendritic spines. *Proc Natl Acad Sci U S A* 1994;91:5720–5724
39. Arnsten AFT, Cai JX, Murphy BL, et al. Dopamine D<sub>1</sub> receptor mechanisms in the cognitive performance of young adult and aged monkeys. *Psychopharmacology (Berl)* 1994;116:143–151
40. Granon S, Passetti F, Thomas KL, et al. Enhanced and impaired attentional performance after infusion of D<sub>1</sub> dopaminergic receptor agents into rat prefrontal cortex. *J Neurosci* 2000;20:1208–1215
41. Lidow MS, Koh P-O, Arnsten AFT. D<sub>1</sub> dopamine receptors in the mouse prefrontal cortex: immunocytochemical and cognitive neuropharmacological analyses. *Synapse* 2003;47:101–108
42. Zahrt J, Taylor JR, Mathew RG, et al. Supranormal stimulation of dopamine D<sub>1</sub> receptors in the rodent prefrontal cortex impairs spatial working memory performance. *J Neurosci* 1997;17:8528–8535
43. Arnsten AFT, Goldman-Rakic PS. Noise stress impairs prefrontal cortical cognitive function in monkeys: evidence for a hyperdopaminergic mechanism. *Arch Gen Psychiatry* 1998;55:362–369
44. Mattay VS, Goldberg TE, Fera F, et al. Catechol O-methyltransferase val158-met genotype and individual variation in the brain response to amphetamine. *Proc Natl Acad Sci U S A* 2003;100:6186–6191
45. Kimberg DY, D'Esposito M, Farah MJ. Effects of bromocriptine on human subjects depend on working memory capacity. *Neuroreport* 1997;8: 3581–3585
46. Sawaguchi T, Goldman-Rakic PS. The role of D<sub>1</sub>-dopamine receptors in working memory: local injections of dopamine antagonists into the prefrontal cortex of rhesus monkeys performing an oculomotor delayed response task. *J Neurophysiol* 1994;71:515–528
47. Druzina MY, Kurzina NP, Malinina EP, et al. The effects of local application of D<sub>2</sub> selective dopaminergic drugs into the medial prefrontal cortex of rats in a delayed spatial choice task. *Behav Brain Res* 2000;109:99–111
48. Wang M, Vijayraghavan S, Goldman-Rakic PS. Selective D<sub>2</sub> receptor actions on the functional circuitry of working memory. *Science* 2004;303:853–856
49. Mrzljak L, Bergson C, Pappy M, et al. Localization of dopamine D<sub>1</sub> receptors in GABAergic neurons of the primate brain. *Nature* 1996;381:245–248
50. Wang X, Zhong P, Yan Z. Dopamine D<sub>4</sub> receptors modulate GABAergic signaling in pyramidal neurons of prefrontal cortex. *J Neurosci* 2002;22: 9185–9193
51. Tahir E, Yazgan Y, Cirakoglu B, et al. Association and linkage of DRD4 and DRD5 with attention deficit hyperactivity disorder (ADHD) in a sample of Turkish children. *Mol Psychiatry* 2000;5:396–404
52. Sunohara GA, Roberts W, Malone M, et al. Linkage of the dopamine D<sub>4</sub> receptor gene and attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry* 2000;39:1537–1542
53. Arnsten AFT, Goldman-Rakic PS. Alpha-2 adrenergic mechanisms in prefrontal cortex associated with cognitive decline in aged nonhuman primates. *Science* 1985;230:1273–1276
54. Cai JX, Ma Y, Xu L, et al. Reserpine impairs spatial working memory performance in monkeys: reversal by the alpha-2 adrenergic agonist clonidine. *Brain Res* 1993;614:191–196
55. Franowicz JS, Kessler L, Dailey-Borja CM, et al. Mutation of the  $\alpha_{2A}$ -adrenoceptor impairs working memory performance and annuls cognitive enhancement by guanfacine. *J Neurosci* 2002;22:8771–8777
56. Birnbaum SG, Arnsten AFT. The alpha-2A noradrenergic agonist, guanfacine, reverses the working memory deficits induced by pharmacological stress (FG7142) [abstract]. *Soc Neurosci Abstracts* 1996;22:1126
57. Arnsten AFT, Contant TA. Alpha-2 adrenergic agonists decrease distractibility in aged monkeys performing a delayed response task. *Psychopharmacology (Berl)* 1992;108:159–169
58. Steere JC, Arnsten AF. The alpha-2A noradrenergic receptor agonist guanfacine improves visual object discrimination reversal performance in aged rhesus monkeys. *Behav Neurosci* 1997;111:883–891
59. Mao Z-M, Arnsten AFT, Li B-M. Local infusion of alpha-1 adrenergic agonist into the prefrontal cortex impairs spatial working memory performance in monkeys. *Biol Psychiatry* 1999;46:1259–1265
60. Wang M, Ji JZ, Li BM. The  $\alpha_{2A}$ -adrenergic agonist guanfacine improves visuomotor associative learning in monkeys. *Neuropsychopharmacology* 2004;29:86–92
61. Jakala P, Riekkinen M, Sirvio J, et al. Guanfacine, but not clonidine, improves planning and working memory performance in humans. *Neuropsychopharmacology* 1999;20:460–470
62. Ramos B, Birnbaum SB, Lindenmayer I, et al. Dysregulation of protein kinase A signaling in the aged prefrontal cortex: new strategy for treating age-related cognitive decline. *Neuron* 2003;40:835–845
63. Li B-M, Mao Z-M, Wang M, et al. Alpha-2 adrenergic modulation of prefrontal cortical neuronal activity related to spatial working memory in monkeys. *Neuropsychopharmacology* 1999;21:601–610
64. Li B-M, Mei Z-T. Delayed response deficit induced by local injection of the alpha-2 adrenergic antagonist yohimbine into the dorsolateral prefrontal cortex in young adult monkeys. *Behav Neural Biol* 1994;62:134–139
65. Ma C-L, Qi X-L, Peng J-Y, et al. Selective deficit in no-go performance induced by blockade of prefrontal cortical  $\alpha_2$ -adrenoceptors in monkeys. *Neuroreport* 2003;14:1013–1016
66. Ma C-L, Arnsten AFT, Li B-M. Locomotor hyperactivity induced by blockade of prefrontal cortical alpha-2-adrenoceptors in monkeys. *Biol Psychiatry* 2005;57:192–195
67. Hunt RD, Arnsten AFT, Asbell MD. An open trial of guanfacine in the treatment of attention deficit hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry* 1995;34:50–54
68. Scahill L, Chappell PB, Kim YS, et al. Guanfacine in the treatment of children with tic disorders and ADHD: a placebo-controlled study. *Am J Psychiatry* 2001;158:1067–1074
69. Taylor FB, Russo J. Comparing guanfacine and dextroamphetamine for the treatment of adult attention deficit-hyperactivity disorder. *J Clin Psychopharmacol* 2001;21:223–228
70. Birnbaum SB, Yuan P, Bloom A, et al. Protein kinase C overactivity impairs prefrontal cortical regulation of working memory. *Science* 2004;306:882–884
71. Arnsten AFT, Mathew R, Ubriani R, et al. Alpha-1 noradrenergic receptor stimulation impairs prefrontal cortical cognitive function. *Biol Psychiatry* 1999;45:26–31
72. Arnsten AFT, Jentsch JD. The alpha-1 adrenergic agonist, cirazoline, impairs spatial working memory performance in aged monkeys. *Pharmacol Biochem Behav* 1997;58:55–59
73. Birnbaum SG, Gobeke KT, Auerbach J, et al. A role for norepinephrine in stress-induced cognitive deficits: alpha-1-adrenoceptor mediation in prefrontal cortex. *Biol Psychiatry* 1999;46:1266–1274
74. Raskind MA, Peskind ER, Kanter ED, et al. Prazosin reduces nightmares and other PTSD symptoms in combat veterans: a placebo-controlled study. *Am J Psychiatry* 2003;160:371–373
75. Manji HK, Lenox RH. Protein kinase C signaling in the brain: molecular transduction of mood stabilization in the treatment of manic-depressive illness. *Biol Psychiatry* 1999;46:1328–1351
76. Williams NM, Preece A, Spurlock G, et al. Support for RGS4 as a susceptibility gene for schizophrenia. *Biol Psychiatry* 2004;55:192–195
77. Biederman J, Mick E, Faraone SV, et al. Pediatric mania: a developmental subtype of bipolar disorder? *Biol Psychiatry* 2000;48:458–466
78. Ramos B, Colgan L, Nou E, et al. The beta-1 adrenergic antagonist, betaxolol, improves working memory performance in rats and monkeys. *Biol Psychiatry*. In press
79. Arnsten AFT. Through the looking glass: differential noradrenergic modulation of prefrontal cortical function. *Neural Plasticity* 2000;7:133–146
80. Arnsten AFT. Stress impairs PFC function in rats and monkeys: role of dopamine D<sub>1</sub> and norepinephrine alpha-1 receptor mechanisms. *Prog Brain Res* 2000;126:183–192
81. Elliott R, Sahakian BJ, Matthews K, et al. Effects of methylphenidate on spatial working memory and planning in healthy young adults. *Psychopharmacology (Berl)* 1997;131:196–206
82. Mehta MA, Owen AM, Sahakian BJ, et al. Methylphenidate enhances working memory by modulating discrete frontal and parietal lobe regions in the human brain. *J Neurosci* 2000;20:RC65
83. Turner DC, Blackwell AD, Dowson JH, et al. Neurocognitive effects of methylphenidate in adult attention-deficit/hyperactivity disorder. *Psychopharmacology (Berl)* 2005;178:286–295
84. Kuczenski R, Segal DS. Exposure of adolescent rats to oral methylphenidate: preferential effects on extracellular norepinephrine and absence of sensitization and cross-sensitization to methamphetamine. *J Neurosci* 2002;22: 7264–7271
85. Arnsten AFT, Dudley AG. Methylphenidate improves prefrontal cortical cognitive function through  $\alpha_2$  adrenoceptor and dopamine D<sub>1</sub> receptor actions: relevance to therapeutic effects in attention deficit hyperactivity disorder. *Behav Brain Funct [serial online]*. 2005;1:2