

**CME Background**

Articles are selected for credit designation based on an assessment of the educational needs of CME participants, with the purpose of providing readers with a curriculum of CME articles on a variety of topics throughout each volume. Activities are planned using a process that links identified needs with desired results.

To obtain credit, read the article, correctly answer the questions in the Posttest, and complete the Evaluation. A \$10 processing fee will apply.

**CME Objective**

After studying this article, you should be able to:

- Include screening for hepatitis C virus as part of the physical health workup in people with serious mental illness

**Accreditation Statement**

The CME Institute of Physicians Postgraduate Press, Inc., is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.



**Credit Designation**

The CME Institute of Physicians Postgraduate Press, Inc., designates this journal-based CME activity for a maximum of 1 *AMA PRA Category 1 Credit™*. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

*Note:* The American Nurses Credentialing Center (ANCC) and the American Academy of Physician Assistants (AAPA) accept certificates of participation for educational activities certified for *AMA PRA Category 1 Credit™* from organizations accredited by the ACCME.

**Release, Expiration, and Review Dates**

This educational activity was published in December 2021 and is eligible for *AMA PRA Category 1 Credit™* through February 29, 2024. The latest review of this material was November 2021.

**Financial Disclosure**

All individuals in a position to influence the content of this activity were asked to complete a statement regarding all relevant personal financial relationships between themselves or their spouse/partner and any commercial interest. The CME Institute has resolved any conflicts of interest that were identified. In the past 3 years, Marlene P. Freeman, MD, Editor in Chief, has received research funding from JayMac and Sage; has been a member of the Independent Data Safety and Monitoring Committee for Janssen (Johnson & Johnson) and Novartis; and has served on advisory boards for Eliem and Sage. As an employee of Massachusetts General Hospital (MGH), Dr Freeman works with the MGH National Pregnancy Registry, which receives funding from Alkermes, Aurobindo, AuroMedics, Johnson & Johnson/Janssen, Otsuka, Sage, Sunovion, Supernus, and Teva, and works with the MGH Clinical Trials Network and Institute, which receives research funding from multiple pharmaceutical companies and the National Institute of Mental Health. No member of the CME Institute staff reported any relevant personal financial relationships. **Faculty financial disclosure appears at the end of the article.**

# Determinants of Hepatitis C Virus Prevalence in People With Serious Mental Illness: A Systematic Review and Meta-Analysis

Michael Rudi Braude, MSc, MBBS<sup>a,\*</sup>;  
 Timothy Phan, MD<sup>a</sup>; Anouk Dev, MBBS, PhD<sup>a,b</sup>;  
 and William Sievert, BA(Hons), MD<sup>a,b</sup>

**ABSTRACT**

**Objective:** To perform a meta-analysis of hepatitis C virus (HCV) prevalence in people with serious mental illness (SMI) and to systematically review barriers to care with the contention that both individual complications and HCV community transmission can be reduced with enhanced health care strategies.

**Data Sources:** PubMed, Scopus, Embase, CINAHL, and Web of Science were searched for articles published in English between April 21, 1989, and July 1, 2020. The terms *Hepatitis C Virus*, *HCV*, *HCV seroprevalence*, and *HCV prevalence* were cross-referenced with *serious mental illness*, *severe mental illness*, *psychiatric illness*, *mental illness*, and *psychiatric patients*.

**Study Selection:** We identified 230 titles after removing duplicates. The final analysis included 36 publications drawn from prospective and large retrospective cohort studies that cross-sectionally screened for HCV in people with SMI ≥ 18 years of age.

**Data Extraction:** Pooled HCV prevalence was analyzed, with random effects modeling due to significant attributable study heterogeneity. Demographic data and HCV risk factors were subanalyzed. Qualitative and semiquantitative data relating to control cohort prevalence and the HCV care cascade were also extracted.

**Results:** The pooled HCV prevalence was 8.0% (95% CI, 6.0%–9.0%). Subanalysis of prospective studies (n = 9,015 individuals) demonstrated a similar prevalence, 8.0% (CI, 5.0%–11.0%), to retrospective studies (n = 289,247), 8.0% (CI, 6.0%–10.0%). HCV was 3.0- to 11.3-fold higher in people with SMI relative to controls. Semiquantitative analysis of seropositive cases showed that (1) 20.0%–58.1% did not have an identified HCV risk factor; (2) 12.5%–100% of cases were not previously known to have HCV; and (3) the majority, 57.0%–96.6%, of people with SMI were receptive to HCV screening.

**Conclusions:** People with SMI have high HCV seroprevalence and should be recognized as a priority group for HCV screening and health care access.

*J Clin Psychiatry* 2022;83(1):21r14079

**To cite:** Braude MR, Phan T, Dev A, et al. Determinants of hepatitis C virus prevalence in people with serious mental illness: a systematic review and meta-analysis. *J Clin Psychiatry*. 2022;83(1):21r14079.

**To share:** <https://doi.org/10.4088/JCP.21r14079>

© Copyright 2021 Physicians Postgraduate Press, Inc.

<sup>a</sup>Department of Gastroenterology and Hepatology, Monash Health, Level 3, 246 Clayton Rd, Clayton, Victoria, Australia

<sup>b</sup>School of Clinical Sciences, Monash University, Wellington Rd, Clayton, Victoria, Australia

\*Corresponding author: Michael Rudi Braude, MSc, MBBS, Department of Gastroenterology and Hepatology, Monash Health, 246 Clayton Rd, Clayton, Victoria, Australia, 3168 (mrh.braude@gmail.com).

### Clinical Points

- Global eradication of hepatitis C virus (HCV) is now achievable but requires intensified case identification among key populations, including people with serious mental illness (SMI).
- Cross-sectional screening for HCV has high acceptance rates among people with SMI and improves detection rates.
- HCV screening should therefore be considered as part of physical health workup and management in people with SMI.

Hepatitis C virus (HCV) is a high morbidity, high mortality condition that affects approximately 1% of the global population.<sup>1</sup> Approximately 399,000 deaths occur annually due to either HCV-related cirrhosis or hepatocellular carcinoma, and HCV presently accounts for 27% of cirrhosis and 25% of hepatocellular carcinoma globally.<sup>1,2</sup> Liver sequelae aside, HCV can impact quality of life as a consequence of neuropsychiatric and ancillary extrahepatic disease manifestations.<sup>3</sup> However, HCV is now curable, and the prospect of global HCV eradication is achievable through the advent of highly effective, highly tolerable, oral direct-acting antiviral (DAA) therapies.<sup>4</sup> Achieving HCV elimination is predicated upon universal health coverage, which requires equitable and financially sustainable health care.<sup>5</sup> However, significant barriers exist, particularly in high-risk transmission populations, such as people who inject drugs (PWID). Providing DAA treatment to PWID is considered an essential strategy toward interrupting HCV transmission.<sup>6</sup> Decentralized HCV elimination strategies in homeless, incarcerated, and opioid-substitution therapy (OST) cohorts have therefore been prioritized.<sup>7</sup>

In contrast, SMI has fundamentally been underrecognized as a priority cohort for HCV screening and management.<sup>8</sup> HCV seroprevalence in people with SMI ranges from 3.0% (95% CI, 1.8%–5.0%) in Central and South America to 17.4% (95% CI, 13.2%–22.6%) in North America.<sup>9</sup> Despite the high prevalence, many mental health settings do not screen for HCV or other blood-borne viruses (BBVs).<sup>8</sup> This stems from underrecognized HCV prevalence, lack of clinician confidence, ambiguity as to whether BBV screening falls within the mental health remit, and absence of established clinical care pathways.<sup>8,10,11</sup> People with SMI have a preponderance of HCV risk factors, including a 16%–22% estimated lifetime prevalence of injecting drug use (IDU), as well as heightened incarceration rates relative to non-SMI populations.<sup>12–15</sup> While these risk factors may tend to overlap with other key, decentralized HCV elimination strategies, such as through OST and prison programs, SMI populations as a whole require focused strategic engagement.<sup>16,17</sup>

We performed a meta-analysis of HCV prevalence in SMI populations and analyzed for heterogeneity between prospective versus retrospective cohort study designs.

We additionally systematically reviewed cross-sectional screening studies and examined data pertaining to HCV risk factors, screening consent, and barriers to care. Our contention was that tailored approaches to HCV screening and health care integration are required in order to achieve parity of care in people with SMI.

### METHODS

#### Search Strategy and Screening

A literature search of PubMed, Scopus, Embase, CINAHL, and Web of Science was performed. Articles, including abstracts, titles, and keywords, published between April 21, 1989, and July 1, 2020, were extracted using the search terms *Hepatitis C Virus*, *HCV*, *HCV seroprevalence*, and *HCV prevalence* cross-referenced with *serious mental illness*, *severe mental illness*, *psychiatric illness*, *mental illness*, and *psychiatric patients*. Details of the analysis were registered on PROSPERO (registration: CRD42021237509).

#### Inclusion and Exclusion Criteria

Prospective observational and retrospective cross-sectional studies that assessed HCV prevalence in adults with SMI aged  $\geq 18$  years were systematically reviewed. HCV prevalence was determined based on *International Classification of Diseases, Ninth Revision (ICD-9)* coded data as well as seroprevalence data. Many of the extracted articles did not elucidate a specific definition for SMI in the recruitment strategy; however, data were included if SMI, severe psychiatric illnesses, or an enduring mental illness with functional impairment was clearly stipulated as an inclusion criterion.


We aimed to analyze HCV prevalence in general SMI populations. Data from dual substance abuse and homeless populations were therefore excluded in order to reduce ascertainment bias. Where sufficient data were available, general SMI subcohorts were specifically analyzed from mixed population settings. Studies were also excluded if (1) found to be a subanalysis of previously published data, (2) the denominator of HCV screened individuals was not cited, or (3) non-serologic methodology, such as self-report, was used for prospective HCV case identification. ICD-coded HCV, rather than serologic diagnosis, was accepted for large registry studies. A cross-sectional BBV screening strategy was required for prospective studies to minimize risk of overestimating seroprevalence due to selection bias. Retrospective and large registry-derived study prevalence data were based on the entire SMI cohort, rather than the screened cohort, again to reduce potential HCV screening selection bias.

#### Extraction and Data Collection

A total of 613 abstracts were identified across each of the search libraries. After removal of duplicates and non-English publications, 230 articles remained. These were reviewed by M.R.B., with 48 relevant titles/abstracts identified. There were 45 full text articles available for analysis (as 3

It is illegal to post this copyrighted PDF on any website.

Figure 1. CONSORT Diagram of Study Inclusion



Abbreviation: HCV = hepatitis C virus.

publications could not be sourced). An additional 4 relevant references were found in the citations from the above articles and assessed for inclusion. Of the 49 full texts, 13 publications did not meet inclusion criteria (Figure 1). This led to a total of 36 publications for analysis. Eligibility and inclusion criteria of full-text publications were verified by T.P. Key variables were recorded by M.R.B. and T.P., including study setting, recruitment strategy, participant demographics, SMI diagnosis, HCV risk factors, HCV screening modality, percentage of population screened, seroprevalence, and associations thereof. The quality of each study was recorded by M.B. and T.P. using the Quality Assessment Tool for Systematic Reviews of Observational Studies (Supplementary Table 1).<sup>18</sup> Interuser concordance was very high, and third-party review was not required. An additional customized global assessment of study quality was performed based on 3 criteria, including study size (< 200 participants = 1, 200–500 = 1, > 500 = 2); cohort selection and reproducibility (single site with minimal participant description = 0, single site with well-described participants/recruitment or large multinational dataset = 1, multisite/national prospective study with clear selection methodology = 2); and risk factor evaluation (no assessment = 0, thorough assessment without analysis of covariables/confounders = 1, thorough risk evaluation with linear regression analysis = 2). Scores were aggregated and assigned a ranking: 0–1 = poor, 2–3 = fair, and 4–6 = good/excellent.

## Analysis

Estimates of prevalence with 95% confidence intervals (CIs) were determined for each individual study. Pooled estimates of prevalence were calculated using the random effects model based on the DerSimonian and Laird method.<sup>19</sup> The random effects model assumes that each study is estimating a different effect but that the effects come from the same underlying distribution of effects. This model fundamentally incorporates differences between studies. Heterogeneity was assessed using the  $I^2$  statistic. Subgroup analysis was performed comparing pooled prevalence between prospective and retrospective studies. A further sensitivity analysis was performed assessing prevalence based on a customized global assessment of study quality. Descriptive analysis was used for evaluation of HCV risk factors. All analyses were performed using Stata Version 16 (StataCorp, College Station, Texas).


## RESULTS

### Prevalence of HCV in SMI: Scale of the Problem

Most prospective seroprevalence data were drawn from studies that recruited consecutive inpatient or community participants from single health services, though 1 study utilized random probability selection to recruit across a large geographical region.<sup>20</sup> Retrospective prevalence data were drawn from single-center seroprevalence studies as

You are prohibited from making this PDF publicly available.

**Figure 2. Pooled Prevalence of HCV in SMI Based on Publications Included in Systematic Review and Meta-Analysis**



Abbreviations: ES = effect size, HCV = hepatitis C virus, SMI = serious mental illness.

well as from large registry studies that used ICD coding for HCV reporting. In total, 36 studies with 298,262 individuals were included.<sup>12,20–54</sup> This included 3 studies that were not necessarily designed to examine HCV seroprevalence but that nonetheless provided cross-sectional seroprevalence data, as well as 2 additional studies that examined HCV prevalence in the broader context of generalized medical comorbidity analysis.<sup>22,29,32,38,48</sup> Key demographics, including country, participant age, and mental health diagnosis, are shown in Supplementary Table 2. Also shown in this table is the absolute number of people with diagnosed HCV in each study. A wide HCV prevalence range was noted (Figure 2). In prospective evaluations, a 1.3% HCV seroprevalence was seen at the lowest end of the spectrum in an Ethiopian SMI cohort.<sup>51</sup> This was in stark contrast to 31.0% seroprevalence in a US-based cohort.<sup>38</sup> Pooled HCV seroprevalence across all included studies was 8.0% (CI, 6.0%–9.0%) (Figure 2). Subanalysis of pooled prevalence

based on study type showed an 8.0% (CI, 5.0%–11.0%) prevalence in prospective studies (n = 9,015 individuals) and an 8.0% (CI, 6.0%–10.0%) prevalence among the retrospective studies (289,247 individuals). When analyzed according to study quality using a customized global assessment of study quality tool, the pooled estimated seroprevalence for good-, fair-, and poor-quality studies was 7.0% (CI, 5.0%–9.0%), 8.0% (5.0%–11.0%), and 10.0% (1.0%–25.0%), respectively.

### HCV Seroprevalence Relative to Control Populations

To contextualize the attributable risk of SMI to HCV, we specifically examined studies that included a non-SMI comparator cohort. Control groups were derived either from non-SMI individuals within large cohort studies or, when reported, from general population data, such as healthy blood donors. A total of 7 studies were included (Table

You are prohibited from making this PDF publicly available.

It is illegal to post this copyrighted PDF on any website.

**Table 1. SMI HCV Prevalence Relative to Control Populations**

Author	Study type	Setting	n	Control group	SMI HCV Ab+, %	Control HCV, %
Sawayama et al <sup>24</sup>	Prospective	Institution	196	Matched serum	10.2	1.5
Nakamura et al <sup>30</sup>	Retrospective	Inpatient	1,193	Healthy blood donors	9.1	1.2
Carney et al <sup>32</sup>	Retrospective	Insurance data	1,074	Cohort without SMI	0.7	0.1
Himelhoch et al <sup>41</sup>	Retrospective	VA dataset	155,172	VA cohort without SMI	7.6	2.5
Fuller et al <sup>53</sup>	Retrospective	VA dataset	11,570	VA cohort without SMI	16.5	1.9
Durotoye et al <sup>49</sup>	Prospective	Inpatient	350	Healthy blood donors <sup>a</sup>	12.4	1.1
Bauer-Staeb et al <sup>12</sup>	Retrospective	National data	97,797	Cohort without SMI	4.6	0.61

<sup>a</sup>Blood donors not propensity/age-matched.

Abbreviations: Ab+ = antibody-positive, HCV = hepatitis C virus, SMI = serious mental illness, VA = Veterans Affairs.

**Table 2. Documented HCV Risk Factors, History of Prior HCV Screening, and Previously Known HCV Diagnosis Among Individuals Included in HCV Prevalence Studies**

Author	Year	Country	Participants, n	HCV Ab+, n	% of HCV Ab+ individuals		
					Nil risk factors	Previously screened	Known HCV
Klinkenberg et al <sup>57</sup>	2003	US	114	34	...	...	17.6
Huckans et al <sup>33</sup>	2006	US	8,836	1,158	...	58.3	...
Freudenreich et al <sup>36</sup>	2007	US	98	8	50.0	...	12.5
Lacey et al <sup>37</sup>	2007	Australia	71	14	...	9.0	...
Tabibian et al <sup>56</sup>	2008	US	129	40	20.0	...	...
Goldberg and Seth <sup>38</sup>	2008	US	100	31	58.1	32.0	...
Rothbard et al <sup>44</sup>	2009	US	588	117	...	50.4	...
Sockalingam et al <sup>46</sup>	2010	Canada	110	3	33.3	...	...
Sanger et al <sup>48</sup>	2013	UK	57	5	...	...	100.0
Ramachandran et al <sup>54</sup>	2018	Australia	260	28	...	100.0	...
Williams et al <sup>52</sup>	2020	Australia	97	7	...	...	57.1
Ayano et al <sup>51</sup>	2020	Ethiopia	309	4	...	25.0	25.0

Abbreviations: Ab+ = antibody-positive, HCV = hepatitis C virus.

1).<sup>12,24,30,32,41,49,53</sup> The studies varied in their respective data collection and comparator populations, though 4 of the studies used large data sets to compare ICD-coded HCV between people with SMI and matched controls. Across each of the studies, HCV seroprevalence was 3.0- to 11.3-fold higher in people with SMI relative to control cases. The range was tighter, 6.8- to 8.7-fold, after excluding lower and upper limit data.<sup>12,24,30,32</sup>

### Risk Factors for HCV in SMI

Given the increased prevalence of HCV in people with SMI, risk factor data were appraised. Lifetime IDU was reported in 10 studies, each prospective, with a pooled size of 4,541 individuals. A wide range (0%–50%) of whole population IDU was noted across these studies (Supplementary Table 3). Studies that subanalyzed IDU among people with seropositive HCV found that 50.0%–78.6% of people with HCV had IDU as a risk factor, although it is notable that these data were drawn from US and Australian cohorts (Supplementary Table 3).<sup>22,25,37,46,54</sup> The apparent heterogeneity in HCV seroprevalence in PWID therefore likely reflects the nature and demographic of the sample cohorts as well as the methodology of data recording and reporting. Illicit substance abuse and/or history of general substance abuse disorder was more widely reported across included studies and, for the most part, mirrored HCV seroprevalence (Supplementary Figure 1). Other reported HCV risk factors included high-risk sexual

encounters, shared skin piercing equipment, transfusions, health care work exposures, and shared razor blades in institutions.<sup>24,37,39</sup> Depression was more strongly associated with HCV compared to other primary mental health conditions based on multivariable analysis in 3 out of 4 studies that evaluated this.<sup>26,34,55,56</sup> Quantitative analysis of risk factors relative to pooled seroprevalence was precluded by variable and lack of standardized reporting between individual studies.

### Risk Factors, Prior Screening, and HCV Awareness Among Seropositive Individuals

As a means of informing screening practices, we appraised the percentage of seropositive individuals from cross-sectional screening programs who had (1) documented HCV risk factors, (2) a history of prior HCV screening, or (3) previously diagnosed HCV. At least 20.0%–58.1% of seropositive individuals did not have a reported HCV risk factor in studies that comprehensively cataloged potential exposures (Table 2).<sup>36,38,46,56</sup> Prior HCV screening in seropositive individuals varied widely (9.0%–100%) but was less than 50% in 3 of 6 studies. In 5 studies that cross-sectionally screened consecutive individuals for BBVs, 27.6% (16/58) of seropositive individuals had been previously diagnosed with HCV.<sup>36,51,52,57</sup> The range of previously undiagnosed cases between studies was broad, likely reflecting sample size, local screening policies, and population-specific factors (Table 2).

**Table 3. Consent for HCV Testing in Prospective SMI Prevalence Studies**

Author	Year	Country	Location	Cohort, n	Participant screening	Consent provided, %
Klinkenberg et al <sup>57</sup>	2003	US	Community	114	Consecutive	57
Tabibian et al <sup>56</sup>	2008	US	Inpatient	188	Consecutive	68.4
Guimarães et al <sup>20</sup>	2009	Brazil	In- and outpatient	2,300	Probability selection	83.2
Lacey et al <sup>37</sup>	2007	Australia	Inpatient	73	Consecutive	18 <sup>a</sup>
Gunewardene et al <sup>45</sup>	2010	Australia	Inpatient	334	Consecutive and at-risk	80
Sanger et al <sup>48</sup>	2013	UK	Inpatient	57	Consecutive	65.5
Ramachandran et al <sup>54</sup>	2018	Australia	Inpatient	260	Consecutive	72.2
Williams et al <sup>52</sup>	2020	Australia	Community	97	Consecutive	81.7
Ayano et al <sup>51</sup>	2020	Ethiopia	Community	309	Consecutive	96.6

<sup>a</sup>This study assessed an HCV education and counseling program, and the consent rate may therefore not correlate with HCV screening alone.

Abbreviations: HCV = hepatitis C virus, SMI = serious mental illness.

### Acceptability of BBV Screening

We next reviewed whether obtaining consent for HCV screening among people with SMI was a potential barrier to accurate delineation of HCV prevalence. We identified 7 prospective studies from inpatient and community settings that documented HCV consent rate and found that most people with SMI, 57.0%–96.6%, agreed to HCV screening (Table 3).<sup>20,45,48,51,52,54,56,57</sup> There was an outlying study in which 18% of participants agreed to HCV screening. This was drawn from an HCV education and screening study and may not reflect general HCV screening acceptability.<sup>37</sup> These data highlight that excellent screening participation rates can be achieved through comprehensive screening programs.

### Liver Function Tests and HCV

Few prevalence studies analyzed liver function test derangement. Freudenreich et al<sup>36</sup> reported an elevated alanine aminotransferase (ALT) in 25% (2/8) HCV seropositive individuals, while Hung et al<sup>47</sup> reported 72.7% (8/11) and Dinwiddie et al<sup>27</sup> reported 39.4% (26/66). In each study, a statistically significant increase in ALT was noted among HCV seropositive individuals compared to HCV seronegative individuals. However, these data indicate that liver function tests alone are incompletely sensitive as a screening tool for HCV screening in SMI. Beyond liver function tests, severity and staging of liver disease have not been prospectively assessed in the SMI context despite the high prevalence of HCV.

### HCV Treatment Uptake and SVR in SMI

We next sought to assess linkages to care and treatment in HCV-positive people with SMI. Several studies have demonstrated excellent DAA treatment outcomes in people with psychiatric disorders.<sup>58,59</sup> However, these data were drawn from individuals who were treated via DAA registry studies or specialist clinic services and therefore likely enriched for people with well-controlled mental health issues. Importantly, these studies did not indicate the denominator of people who did not access care following an HCV diagnosis.

Small prospective studies have shown that linking viremic individuals with SMI to HCV treatment can be complex. Among 72 individuals with viremic HCV

(pooled from 3 prospective seroprevalence studies), 38 were successfully linked to care, and 9 individuals were ultimately treated, mostly with DAA therapies.<sup>36,38,54</sup> The individuals who were successfully treated required integrated health service delivery via “nontraditional” services including support from hepatitis outreach nurses and mental health case managers. Cited barriers to care included lack of referral to specialist services and patient-related factors including refusal to attend traditional outpatient specialist care services and refusal to commence DAA therapies.<sup>36,54</sup>

Notwithstanding the potential challenges of translating HCV diagnosis to successful treatment, retrospective US Veterans Affairs data have shown that Veterans with SMI achieve similar linkage to treatment when compared to an internal Veterans Affairs control cohort (11.9% versus 13.9%).<sup>33</sup> In a second study, 33% of Veterans who were diagnosed with HCV during an inpatient mental health admission were successfully commenced on interferon following referral to outpatient specialist services.<sup>35</sup> This compares favorably to general population real-world interferon data, whereby a pooled rate of 19% interferon treatment initiation has been estimated.<sup>60</sup>

## DISCUSSION

### What Drives Increased HCV Risk in People With SMI?

In our meta-analysis, pooled HCV seroprevalence in people with SMI was 8.0%, that is, at least 8-fold higher than the estimated global prevalence.<sup>1</sup> The overrepresentation of HCV in people with SMI appears to be increasingly driven by substance abuse, a problem deeply embedded within social disadvantage and one that may be perpetuated by cognitive issues and positive symptoms related to SMI.<sup>12–15,61,62</sup> In terms of relative effect size, Bauer-Staeb et al<sup>12</sup> demonstrated that the odds ratio of HCV seropositivity in a Swedish national cross-sectional analysis of people with SMI decreased from 6.18 to 1.72 after adjusting for substance abuse. Substance abuse is common among people with SMI, and in a previous systematic review of HIV risk factors in SMI, IDU was found to have a weighted mean lifetime prevalence of 21.7% (14.0%–37.0%), albeit there was a preponderance of US data.<sup>15</sup> The relevance of this is that a history of IDU correlates with a >40% HCV

prevalence (in 61 out of 77 countries).<sup>63</sup> Substance abuse, and in particular IDU, may explain much of the disproportionate increase in HCV seroprevalence in people with SMI and therefore requires strong consideration when screening for HCV.

### Rationale for BBV Screening in SMI and Barriers to Care

We show that BBV screening delivered through population-based programs has high acceptability among people with SMI. Nonetheless, real-world data suggest that people within mental health services are underserved from an HCV screening perspective. In a cross-sectional analysis of 57,170 SMI outpatients, 4.7% were screened for HCV over a 12-month period as compared to 12.4% in a general population comparator cohort, with the strongest predictor of HCV screening in the SMI group being non-psychiatric health care utilization.<sup>64</sup> Importantly, person-centered blood-borne virus screening delivered through collocated specialist services has been shown to significantly bolster HCV screening rates in community-managed individuals with SMI.<sup>65</sup> However, even with positive case identification, significant attrition in follow-up and treatment is commonly encountered.<sup>65-67</sup> Few publications in this systematic review addressed linkage to care in HCV-positive individuals; however, known barriers exist as a composite of complex patient-related, clinician, and health system factors.<sup>8,10,66</sup> For example, issues pertaining to access and retention within traditional specialist health care models may be experienced at a patient level. This may be compounded by perceived or actual stigma.<sup>66</sup> From a clinician perspective, legacy concerns stemming from interferon-era therapies coupled with competing mental health care priorities may hinder advocacy and referral to HCV treatment services.<sup>66,68</sup>

### Looking to Other Models of Care

Mental health has lagged other high-prevalence HCV populations in terms of decentralization of screening programs,<sup>8,69</sup> although some data exist regarding intensified support within traditional outpatient models.<sup>70</sup> In other high-risk settings, such as the prison and opioid-substitution contexts, decentralized models of HCV care service delivery augmented by community nurse engagement, case management, and enhanced motivational support have demonstrated efficacy in engaging and improving HCV treatment access when compared to traditional health care models.<sup>17</sup> People with SMI are often integrated within community mental health services, and a remodeling of the current care paradigm bolstered by enhanced clinician education could translate to effective BBV screening and treatment.<sup>35,70-72</sup> These pathways could be supported through point-of-care HCV antigen assays, integration of nurse practitioners, improved clinician education, streamlined screening strategies, and horizontal integration of specialists, through either visiting sites or telehealth platforms.<sup>70,73</sup> Notwithstanding potential barriers in the SMI setting, decentralization of traditional models of care could

yield markedly improved access to highly tolerable curative therapies.<sup>58,59,74</sup>

### Recommendations for Screening

With regard to the nature of BBV screening, a patient-centered, codesign screening approach would likely yield enhanced patient acceptability and engagement. Screening programs should ideally be applied cross-sectionally in order to both reduce perceived stigma and increase HCV diagnostic yield relative to individualized risk factor-based screening. Indeed, several cross-sectional screening studies included in this analysis show that HCV risk factors are not always ascertained in seropositive individuals and that many people with seropositive HCV were previously undiagnosed via traditional screening algorithms.<sup>33,36,38,44,46,54,56</sup> Cross-sectional screening has the advantage of bypassing issues like incomplete risk factor appraisal by clinicians, patient recall bias, and/or stigma-related underreporting. Overall, a broader approach to screening would serve to change cultural attitudes toward BBV screening and improve HCV identification rates.

### Study Limitations

Convenience samples included in the meta-analysis may have enriched for individuals at greater risk of HCV due to higher acuity psychosocial issues. We attempted to adjust for this in a number of ways. First, SMI cohorts drawn from dual substance abuse disorders settings were excluded. Furthermore, we specifically included retrospective studies, including those from large registry datasets, given that these data generally have a greater cross-sectional reach. Pooled seroprevalence was very similar between prospective and retrospective subcohorts despite methodological heterogeneity and remained similar in a study quality subanalysis, and therefore appears to be robust.

Notwithstanding this, marked variability in HCV seroprevalence between studies was noted. Background population HCV prevalence is likely to be a large contributing factor. For example, the general population HCV seroprevalence in Belgium is 0.57%–0.90% as compared to 1.2%–2.4% in the US.<sup>12,75</sup> The effect size of background seroprevalence heterogeneity is likely amplified by risk factors preponderant to specific SMI cohorts. These risks may vary in frequency between regions and may even be idiosyncratic to a local cohort, such as in the case of shared razor blades as a dominant driver of HCV transmission in an institutionalized cohort of people with SMI in Japan.<sup>24</sup> Other ascertainment biases related to study design and data acquisition may exaggerate the observed heterogeneity in HCV SMI seroprevalence. In particular, small cohort prospective data may not represent national SMI HCV seroprevalence, particularly where few representative studies exist. Regardless of absolute seroprevalence, HCV was reproducibly 3.0 to 11.3 times higher in SMI cohorts compared to controls despite heterogeneity between individual studies.<sup>12,24,30,32,33,41,51,53</sup> These data demonstrate that SMI is associated with preponderant HCV and

It is illegal to post this copyrighted PDF on any website. reinforces that people with SMI should be considered as a higher risk cohort from a BBV perspective.

## CONCLUSIONS

People with SMI have high HCV seroprevalence, and a major risk factor is high-prevalence substance abuse. Given the availability of highly effective DAA therapies, tailored HCV screening and treatment programs would invariably reduce avoidable liver and extrahepatic complications at an individual level and would contribute toward global efforts

toward HCV elimination. A first step is to recognize SMI as a priority group for HCV screening and to adopt broad population screening with an emphasis on a patient-centered approach. This is acknowledging that BBV screening has high acceptability among individuals with SMI and that cross-sectional screening is effective in identifying previously undiagnosed HCV. In terms of bridging existing barriers, innovative models of care that maximize opportunity for destigmatized screening and minimize complexity of follow-up will likely translate to an improved cascade of care and enhanced health outcomes.

**Submitted:** May 12, 2021; accepted September 10, 2021.

**Published online:** December 14, 2021.

**Disclosure of off-label usage:** The authors have determined that, to the best of their knowledge, no investigational information about pharmaceutical agents or device therapies that is outside US Food and Drug Administration–approved labeling has been presented in this article.

**Financial disclosure:** Drs Braude, Phan, Dev, and Sievert have no conflicts to disclose.

**Funding/support:** None.

**Supplementary material:** Available at PSYCHIATRIST.COM.

## REFERENCES

- World Health Organization. Global hepatitis report, 2017. Accessed Nov 2020. <https://www.who.int/publications/i/item/global-hepatitis-report-2017>
- Perz JF, Armstrong GL, Farrington LA, et al. The contributions of hepatitis B virus and hepatitis C virus infections to cirrhosis and primary liver cancer worldwide. *J Hepatol*. 2006;45(4):529–538.
- Dammacco F, Sansonno D, Piccoli C, et al. The lymphoid system in hepatitis C virus infection: autoimmunity, mixed cryoglobulinemia, and overt B-cell malignancy. *Semin Liver Dis*. 2000;20(2):143–157.
- Dore GJ, Martinello M, Alavi M, et al. Global elimination of hepatitis C virus by 2030: why not? *Nat Med*. 2020;26(2):157–160.
- World Health Assembly. Resolutions and decisions, annexes. 69th World Health Assembly: May 23–28, 2016. <https://apps.who.int/iris/handle/10665/259134>
- Grebel J, Dore GJ. Treatment of HCV in persons who inject drugs: treatment as prevention. *Clin Liver Dis (Hoboken)*. 2017;9(4):77–80.
- Hellard M, Rolls DA, Sacks-Davis R, et al. The impact of injecting networks on hepatitis C transmission and treatment in people who inject drugs. *Hepatology*. 2014;60(6):1861–1870.
- Wainberg M, Dixon L. Ending HIV, hepatitis B, and hepatitis C: what about people with severe mental illness? *Lancet Psychiatry*. 2017;4(9):651–653.
- Hughes E, Bassi S, Gilbody S, et al. Prevalence of HIV, hepatitis B, and hepatitis C in people with severe mental illness: a systematic review and meta-analysis. *Lancet Psychiatry*. 2016;3(1):40–48.
- De Hert M, Cohen D, Bobes J, et al. Physical illness in patients with severe mental disorders, 2: barriers to care, monitoring and treatment guidelines, plus recommendations at the system and individual level. *World Psychiatry*. 2011;10(2):138–151.
- Lagios K, Deane F. Psychiatrists' knowledge and practices in screening and assessment of hepatitis C for inpatients with severe mental illness. *Australas Psychiatry*. 2011;19(2):156–159.
- Bauer-Staeb C, Jørgensen L, Lewis G, et al. Prevalence and risk factors for HIV, hepatitis B, and hepatitis C in people with severe mental illness: a total population study of Sweden. *Lancet Psychiatry*. 2017;4(9):685–693.
- Lagios K, Deane FP. Severe mental illness is a new risk marker for blood-borne viruses and sexually transmitted infections. *Aust N Z J Public Health*. 2007;31(6):562–566.
- Sheidow AJ, McCart M, Zajac K, et al. Prevalence and impact of substance use among emerging adults with serious mental health conditions. *Psychiatr Rehabil J*. 2012;35(3):235–243.
- Meade CS, Sikkema KJ. HIV risk behavior among adults with severe mental illness: a systematic review. *Clin Psychol Rev*. 2005;25(4):433–457.
- Fragomeli V, Weltman M. Addressing viral hepatitis in the opiate substitution setting: an integrated nursing model of care. *J Gastroenterol Hepatol*. 2015;30(suppl 2):6–11.
- Papaluca T, McDonald L, Craigie A, et al. Outcomes of treatment for hepatitis C in prisoners using a nurse-led, statewide model of care. *J Hepatol*. 2019;70(5):839–846.
- Quality assessment tool for observational cohort and cross-sectional studies. National Heart, Lung, and Blood Institute. 2014. Retrieved Sept 16, 2020. <https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools>
- DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials*. 1986;7(3):177–188.
- Guimarães MD, Campos LN, Melo AP, et al; PESSOAS Project Network Group. Prevalence of HIV, syphilis, hepatitis B and C among adults with mental illness: a multicenter study in Brazil. *Br J Psychiatry*. 2009;31(1):43–47.
- Chang TT, Lin H, Yen YS, et al. Hepatitis B and hepatitis C among institutionalized psychiatric patients in Taiwan. *J Med Virol*. 1993;40(2):170–173.
- Di Nardo V, Petrosillo N, Ippolito G, et al. Prevalence and incidence of hepatitis B virus, hepatitis C virus and human immunodeficiency virus among personnel and patients of a psychiatric hospital. *Eur J Epidemiol*. 1995;11(2):239–242.
- Cividini A, Pistorio A, Regazzetti A, et al. Hepatitis C virus infection among institutionalized psychiatric patients: a regression analysis of indicators of risk. *J Hepatol*. 1997;27(3):455–463.
- Sawayama Y, Hayashi J, Kakuda K, et al. Hepatitis C virus infection in institutionalized psychiatric patients: possible role of transmission by razor sharing. *Dig Dis Sci*. 2000;45(2):351–356.
- Rosenberg SD, Goodman LA, Osher FC, et al. Prevalence of HIV, hepatitis B, and hepatitis C in people with severe mental illness. *Am J Public Health*. 2001;91(1):31–37.
- Al Jurdi RK, Burruss JW. Prevalence of hepatitis C in psychiatric institutions. *Psychosomatics*. 2003;44(5):439–440.
- Dinwiddie SH, Shicker L, Newman T. Prevalence of hepatitis C among psychiatric patients in the public sector. *Am J Psychiatry*. 2003;160(1):172–174.
- Meyer JM. Prevalence of hepatitis A, hepatitis B, and HIV among hepatitis C-seropositive state hospital patients: results from Oregon State Hospital. *J Clin Psychiatry*. 2003;64(5):540–545.
- Kilbourne AM, Cornelius JR, Han X, et al. Burden of general medical conditions among individuals with bipolar disorder. *Bipolar Disord*. 2004;6(5):368–373.
- Nakamura Y, Koh M, Miyoshi E, et al. High prevalence of the hepatitis C virus infection among the inpatients of schizophrenia and psychoactive substance abuse in Japan. *Prog Neuropsychopharmacol Biol Psychiatry*. 2004;28(3):591–597.
- Pirl WF, Greer JA, Weissgarber C, et al. Screening for infectious diseases among patients in a state psychiatric hospital. *Psychiatr Serv*. 2005;56(12):1614–1616.
- Carney CP, Jones L, Woolson RF. Medical comorbidity in women and men with schizophrenia: a population-based controlled study. *J Gen Intern Med*. 2006;21(11):1133–1137.
- Huckans MS, Blackwell AD, Harms TA, et al. Management of hepatitis C disease among VA patients with schizophrenia and substance use disorders. *Psychiatr Serv*. 2006;57(3):403–406.
- Raja M, Azzoni A, Pucci D. Characteristics of HCV positive patients in an Italian urban psychiatric unit. *Clin Pract Epidemiol Ment Health*. 2006;2(1):26.
- Rifai MA, Moles JK, Short DD. Hepatitis C treatment eligibility and outcomes among patients with psychiatric illness. *Psychiatr Serv*. 2006;57(4):570–572.
- Freudenreich O, Gandhi RT, Walsh JP, et al. Hepatitis C in schizophrenia: screening experience in a community-dwelling clozapine cohort. *Psychosomatics*. 2007;48(5):405–411.
- Lacey C, Ellen S, Devlin H, et al. Hepatitis C in psychiatry inpatients: testing rates, prevalence and risk behaviours. *Australas Psychiatry*. 2007;15(4):315–319.
- Goldberg RW, Seth P. Hepatitis C services and individuals with serious mental illness. *Community Ment Health J*. 2008;44(5):381–384.
- Alvarado-Esquivel C, Arreola-Valenzuela MA, Rodríguez-Briones A, et al. Seroprevalence of



# It is illegal to post this copyrighted PDF on any website.

- selected viral, bacterial and parasitic infections among inpatients of a public psychiatric hospital of Mexico. *Rev Inst Med Trop São Paulo*. 2008;50(3):161–164.
40. De Hert M, Wampers M, Van Eyck D, et al. Prevalence of HIV and hepatitis C infection among patients with schizophrenia. *Schizophr Res*. 2009;108(1–3):307–308.
  41. Himelhoch S, McCarthy JF, Ganoczy D, et al. Understanding associations between serious mental illness and hepatitis C virus among veterans: a national multivariate analysis. *Psychosomatics*. 2009;50(1):30–37.
  42. Kakisi OK, Grammatikos AA, Karageorgopoulos DE, et al. Prevalence of hepatitis B, hepatitis C, and HIV infections among patients in a psychiatric hospital in Greece. *Psychiatr Serv*. 2009;60(9):1269–1272.
  43. Mamani M, Hashemi SH, Niayesh A, et al. Study on the frequency of hepatitis B and C infection in chronic psychiatric patients in Hamedan in 2006–2007. *J Pak Med Assoc*. 2009;59(8):505–507.
  44. Rothbard AB, Blank MB, Staab JP, et al. Previously undetected metabolic syndromes and infectious diseases among psychiatric inpatients. *Psychiatr Serv*. 2009;60(4):534–537.
  45. Gunewardene R, Lampe L, Ilchef R. Prevalence of hepatitis C in two inpatient psychiatry populations. *Australas Psychiatry*. 2010;18(4):330–334.
  46. Sockalingam S, Shammi C, Powell V, et al. Determining rates of hepatitis C in a clozapine treated cohort. *Schizophr Res*. 2010;124(1–3):86–90.
  47. Hung CC, Loh W, Hu TM, et al. Prevalence of hepatitis B and hepatitis C in patients with chronic schizophrenia living in institutions. *J Chin Med Assoc*. 2012;75(6):275–280.
  48. Sanger C, Hayward J, Patel G, et al. Acceptability and necessity of HIV and other blood-borne virus testing in a psychiatric setting. *Br J Psychiatry*. 2013;202(4):307–308.
  49. Durotoye IA, Issa BA, Fadeyi A, et al. Seroprevalence of hepatitis B and C among mentally ill patients attending a tertiary hospital in Nigeria. *Ann Afr Med*. 2014;13(4):210–216.
  50. Karabulut N. Prevalence of HBV, HCV and HIV in inpatients of a mental health hospital in Turkey, 2011–2013. *Iran J Public Health*. 2015;44(7):1026–1028.
  51. Ayano G, Haile K, Tesfaye A, et al. Undiagnosed HIV, hepatitis B, and hepatitis C infections in people with severe psychiatric disorders in Ethiopia. *BMC Infect Dis*. 2020;20(1):180.
  52. Williams J, Barclay M, Omana C, et al. Universal blood-borne virus screening in patients with severe mental illness managed in an outpatient clozapine clinic: uptake and prevalence. *Australas Psychiatry*. 2020;28(2):186–189.
  53. Fuller BE, Rodriguez VL, Linke A, et al. Prevalence of liver disease in veterans with bipolar disorder or schizophrenia. *Gen Hosp Psychiatry*. 2011;33(3):232–237.
  54. Ramachandran J, Budd S, Slattery H, et al. Hepatitis C virus infection in Australian psychiatric inpatients: a multicenter study of seroprevalence, risk factors and treatment experience. *J Viral Hepat*. 2019;26(5):609–612.
  55. Himelhoch S, Goldberg R, Calmes C, et al. Screening for and prevalence of HIV and hepatitis C among an outpatient urban sample of people with serious mental illness and co-occurring substance abuse. *J Community Psychol*. 2011;39(2):231–239.
  56. Tabibian JH, Wirshing DA, Pierre JM, et al. Hepatitis B and C among veterans on a psychiatric ward. *Dig Dis Sci*. 2008;53(6):1693–1698.
  57. Klinkenberg WD, Caslyn RJ, Morse GA, et al. Prevalence of human immunodeficiency virus, hepatitis B, and hepatitis C among homeless persons with co-occurring severe mental illness and substance use disorders. *Compr Psychiatry*. 2003;44(4):293–302.
  58. Back D, Belperio P, Bondin M, et al. Efficacy and safety of glecaprevir/pibrentasvir in patients with chronic HCV infection and psychiatric disorders: an integrated analysis. *J Viral Hepat*. 2019;26(8):951–960.
  59. Gayam V, Jegede O, Tiongson B, et al. Outcomes of direct-acting antiviral treatment of psychiatric patients with comorbid hepatitis C virus infection. *Dig Dis*. 2020;38(3):232–239.
  60. North CS, Hong BA, Adewuyi SA, et al. Hepatitis C treatment and SVR: the gap between clinical trials and real-world treatment aspirations. *Gen Hosp Psychiatry*. 2013;35(2):122–128.
  61. Draine J, Salzer MS, Culhane DP, et al. Role of social disadvantage in crime, joblessness, and homelessness among persons with serious mental illness. *Psychiatr Serv*. 2002;53(5):565–573.
  62. Parks J, Svendsen D, Singer P, et al. *Morbidity and Mortality in People with Serious Mental Illness*. Alexandria, VA: National Association of State Mental Health Program Directors (NASMHPD) Medical Directors Council; 2006.
  63. Nelson PK, Mathers BM, Cowie B, et al. Global epidemiology of hepatitis B and hepatitis C in people who inject drugs: results of systematic reviews. *Lancet*. 2011;378(9791):571–583.
  64. Trager E, Khalili M, Masson CL, et al. Hepatitis C screening rate among underserved adults with serious mental illness receiving care in California community mental health centers. *Am J Public Health*. 2016;106(4):740–742.
  65. Rosenberg SD, Goldberg RW, Dixon LB, et al. Assessing the STIRR model of best practices for blood-borne infections of clients with severe mental illness. *Psychiatr Serv*. 2010;61(9):885–891.
  66. Chasser Y, Kim AY, Freudenreich O. Hepatitis C treatment: clinical issues for psychiatrists in the post-interferon era. *Psychosomatics*. 2017;58(1):1–10.
  67. McLaren M, Garber G, Cooper C. Barriers to hepatitis C virus treatment in a Canadian HIV-hepatitis C virus coinfection tertiary care clinic. *Can J Gastroenterol*. 2008;22(2):133–137.
  68. Dixon LB, Adler DA, Berlant JL, et al. Psychiatrists and primary caring: what are our boundaries of responsibility? *Psychiatr Serv*. 2007;58(5):600–602.
  69. Rifai MA, Gleason OC, Sabouni D. Psychiatric care of the patient with hepatitis C: a review of the literature. *Prim Care Companion J Clin Psychiatry*. 2010;12(6):PCC.09r00877whi.
  70. Knott A, Dieperink E, Willenbring ML, et al. Integrated psychiatric/medical care in a chronic hepatitis C clinic: effect on antiviral treatment evaluation and outcomes. *Am J Gastroenterol*. 2006;101(10):2254–2262.
  71. Morgan VA, Waterreus A, Jablensky A, et al. People living with psychotic illness in 2010: the second Australian national survey of psychosis. *Aust N Z J Psychiatry*. 2012;46(8):735–752.
  72. Bonner JE, Barritt AS 4th, Fried MW, et al. Time to rethink antiviral treatment for hepatitis C in patients with coexisting mental health/substance abuse issues. *Dig Dis Sci*. 2012;57(6):1469–1474.
  73. Arora S, Kalishman S, Thornton K, et al. Expanding access to hepatitis C virus treatment—Extension for Community Healthcare Outcomes (ECHO) project: disruptive innovation in specialty care. *Hepatology*. 2010;52(3):1124–1133.
  74. de Gennaro N, Diella L, Monno L, et al. Efficacy and tolerability of DAAs in HCV-monoinfected and HCV/HIV-coinfected patients with psychiatric disorders. *BMC Infect Dis*. 2020;20(1):196.
  75. Gower E, Estes C, Blach S, et al. Global epidemiology and genotype distribution of the hepatitis C virus infection. *J Hepatol*. 2014;61(1 suppl):S45–S57.

See supplementary material for this article at [PSYCHIATRIST.COM](https://www.psychiatrist.com).

For the CME Posttest, see next page.



## POSTTEST

To obtain credit, go to [PSYCHIATRIST.COM](http://PSYCHIATRIST.COM) to take this Posttest and complete the Evaluation. A \$10 processing fee is required.

1. Regarding the pooled global prevalence of hepatitis C virus (HCV) in people with serious mental illness (SMI), which of the following is most correct?
  - a. 1 in 1,200 people with SMI will have HCV
  - b. 1 in 120 people with SMI will have HCV
  - c. 1 in 12 people with SMI will have HCV
  - d. 1 in 6 people with SMI will have HCV
2. A history of injecting drug use (IDU) correlates with a > 40% HCV prevalence. True or false?
  - a. True
  - b. False
3. Paul is a 42-year-old man who was diagnosed with bipolar I disorder in his early 20s. He has been prescribed various medications to treat the disorder over the years, although he has often been nonadherent to his treatment. Paul has a history of substance abuse, and in particular IDU. In a recent appointment at your clinical practice, you recognize Paul's risk factors for HCV and would like to recommend HCV screening to him. However, you are unsure of the outcome of your recommendation. According to this meta-analysis, which of the following is most correct and should be considered in your decision to suggest screening to Paul and other patients? HCV screening in people with SMI is:
  - a. Acceptable among most patients
  - b. Not accepted by most patients
  - c. Is contraindicated due to risk of destabilizing underlying mental health issues
  - d. Is contraindicated in most cases due to lack of patient autonomy to provide consent

You are prohibited from making this PDF publicly available.



THE JOURNAL OF  
**CLINICAL PSYCHIATRY**

THE OFFICIAL JOURNAL OF THE AMERICAN SOCIETY OF CLINICAL PSYCHOPHARMACOLOGY

## **Supplementary Material**

**Article Title:** Determinants of Hepatitis C Virus Prevalence in People With Serious Mental Illness: A Systematic Review and Meta-Analysis

**Author(s):** Michael Rudi Braude, MSc, MBBS; Timothy Phan, MD; Anouk Dev, MBBS, PhD; and William Sievert, BA(Hons), MD

**DOI Number:** 10.4088/JCP.21r14079

### **List of Supplementary Material for the article**

1. [Table 1](#) Quality Analysis of Studies Using the NIH Quality Assessment Tool and a Customized Quality Assessment Tool
2. [Table 2](#) Key Demographics and Seroprevalence From Individual Prevalence Studies
3. [Table 3](#) Rates of IDU in Prospective HCV Seroprevalence Studies
4. [Figure 1](#) Trend in HCV Antibody Prevalence Relative to Prevalence of Reported Substance Abuse Across 14 BBV Prevalence Studies in the Context of SMI

### **Disclaimer**

This Supplementary Material has been provided by the author(s) as an enhancement to the published article. It has been approved by peer review; however, it has undergone neither editing nor formatting by in-house editorial staff. The material is presented in the manner supplied by the author.

**Supplementary Table 1:** Quality analysis of studies using the NIH quality assessment tool and a customized quality assessment tool<sup>‡</sup>

Author	Quality based on NIH score	Customized quality assessment score				Overall assessment
		Study size	Patient selection	Risk factor assessment	Total	
Chang et al	8	2	1	1	4	Good
Di Nardo et al	10	0	1	1	2	Fair
Cividini et al	10	1	2	0	3	Fair
Sawayama et al	9	0	1	2	3	Fair
Rosenberg et al	11	2	2	2	6	Good
Al Jurdi and Burruss	7	1	1	1	3	Fair
Dinwiddie et al	6	2	0	0	2	Fair
Meyer	6	2	0	0	2	Fair
Nakamura et al	6	2	0	0	2	Fair
Kilbourne et al	5	2	1	0	3	Fair
Pirl et al	8	2	1	2	5	Good
Rifai et al	6	2	0	0	2	Fair
Raja et al	10	2	1	1	4	Good
Carney et al	7	2	1	0	3	Fair
Huckans et al	8	2	1	1	4	Good
Freudenreich et al	9	0	0	1	1	Poor
Lacey et al	9	0	1	1	2	Fair
Goldberg et al	8	0	0	0	0	Poor
Rothbard et al	9	2	1	0	3	Fair
Alvarado-Esquivel et al	9	0	1	1	2	Fair
Guimarães et al	10	2	2	0	4	Good
De Hert et al	7	2	0	0	2	Fair
Mamani et al	8	0	0	1	1	Poor
Himelhoch et al	11	2	1	1	4	Good
Kakisi et al	5	2	0	0	2	Fair
Sockalingam et al	9	0	1	1	2	Fair
Gunewardene et al	10	1	0	1	2	Fair
Fuller et al	9	2	1	1	4	Good
Hung et al	8	2	0	0	2	Fair
Sanger et al	7	0	1	0	1	Poor
Durotoye et al	7	1	0	1	2	Fair
Karabulut	4	2	0	0	2	Fair
Bauer-Staeb et al	10	2	2	2	6	Good
Ramachandran et al	12	1	1	2	4	Good
Williams et al	8	0	1	1	2	Fair
Ayano et al	11	1	2	1	4	Good

<sup>‡</sup>The customized quality assessment score is outlined in the Extraction and Data Collection section.

**Supplementary Table 2:** Key demographics and seroprevalence from individual prevalence studies.

Abbreviations: P = prospective; R = retrospective; SCZ = schizophrenia, BAD = bipolar affective disorder; MDD = major depressive disorder

Author	Year	Country	(n)	Study Type	Male (%)	Age (years)	Mental Health Diagnosis			
							SCZ (%)	BAD (%)	MDD (%)	Other (%)
Chang et al	1993	Taiwan	780	P	66.5	42.0	96.2	-	-	3.8
Di Nardo et al	1995	Italy	176	P	-	-	-	-	-	-
Cividini et al*	1997	Italy	423	P	-	55.0	-	-	-	-
Sawayama et al*	2000	Japan	163	P	52.0	57.9	-	-	-	-
Rosenberg et al	2001	USA	751	P	-	-	44.9	16.8	11.7	19.9
Al Jurdi and Burruss	2003	USA	238	P	50.4	-	44.1	12.6	31.1	-
Dinwiddie et al	2003	USA	1,556	R	73.3	38.4	9.8	26.5	18.7	45.0
Meyer	2003	USA	535	R	74.8	42.7	74.8	6.8	4.8	-
Nakamura et al	2004	Japan	1,193	R	71.1	50.4	37.6	-	9.2	40.7
Kilbourne et al*	2004	USA	4,310	R	90.0	53.0	-	100	-	-
Pirl et al‡	2005	USA	655	R	-	-	34.0	20	-	-
Rifai et al§	2006	USA	3,470	R	-	-	-	-	-	-
Raja et al	2006	Italy	1,492	P	44.5	41.9	9.0	34.1	9.0	9.0
Carney et al*	2006	USA	1,074	R	47.0	40.2	100	-	-	-
Huckans et al	2006	USA	4,644	R	93.3	48.9	100	-	-	-
Freudenreich et al	2007	USA	98	P	75.0	44.7	100	-	-	-
Lacey et al	2007	Australia	71	P	-	30.0	73.0	-	-	-
Goldberg et al§	2008	USA	100	P	54.0	45.1	-	-	-	-
Rothbard et al	2009	USA	656	R	48.0	42.0	33.0	-	57.0	-
Alvarado-Esquivel et al	2009	Mexico	105	P	71.4	46.5	25.7	4.8	1.0	-
Guimarães et al	2009	Brazil	2,300	P	48.1	-	47.3	9.0	13.3	-
De Hert et al	2009	Belgium	595	R	65.0	36.7	81.4	-	-	18.6
Mamani et al	2009	Iran	170	P	56.0	-	49.0	-	-	-
Himelhoch et al	2009	USA	155,172	R	92.5	54.8	57.5	42.5	-	-
Kakisi et al	2009	Turkey	793	R	74.7	-	-	-	-	-
Sockalingam et al	2010	Canada	110	P	-	-	100	-	-	-
Gunewardene et al	2010	Australia	334	P	-	-	-	-	-	-
Fuller et al	2011	USA	11,570	R	91.5	55.7	54.0	46.0	-	-
Hung et al	2012	Taiwan	590	P	58.4	42.5	100	-	-	-
Sanger et al§	2013	UK	57	P	71.9	-	-	-	-	-
Durotoye et al	2014	Nigeria	350	P	51.1	36.5	-	-	-	-
Karabulut	2015	Turkey	5,227	R	78.3	35.5	-	-	-	-
Bauer-Staeb et al	2017	Sweden	97,797	R	48.1	52.0	21.7	-	-	6.3
Ramachandran et al	2018	Australia	260	P	70.0	44.0	19.0	10.0	26.0	-
Williams et al	2020	Australia	97	P	-	-	100	-	-	-
Ayano et al	2020	Ethiopia	309	P	65.4	36.2	43.7	17.5	29.8	9.1


\*Patients with psychosis - sub-type not explicated

\*Study designed to assess for comorbidity, not BBVs or HCV specifically

§Study not designed for prevalence assessment, but sufficient cross-sectional data available

**Supplementary Table 3:** Rates of IDU in prospective HCV seroprevalence studies

<b>Author</b>	<b>Year</b>	<b>Country</b>	<b>(n)</b>	<b>Study Type</b>	<b>HCV Ab (%)</b>	<b>Lifetime IDU (%)</b>	<b>IDU in HCV Ab +ve (%)</b>
Chang et al	1993	Taiwan	780	Prospective	6.8	0.0	-
Di Nardo et al	1995	Italy	176	Prospective	11.4	0.0	-
Sawayama et al	2000	Japan	196	Prospective	10.2	1.0	-
Rosenberg et al	2001	USA	751	Prospective	16.1	12.1	75.2
Al Jurdi and Burruss	2003	USA	238	Prospective	16.0	13.9	-
Lacey et al	2007	Australia	71	Prospective	19.7	50.0	50.0
Guimarães et al	2009	Brazil	2300	Prospective	2.63	3.0	-
Sockalingham et al	2010	Canada	110	Prospective	2.7	8.0	66.7
Gunewardene et al	2010	Australia	334	Prospective	5.6	14.4	-
Ramachandran et al	2018	Australia	260	Prospective	10.8	28.0	78.6



**Supplementary Figure 1:** Trend in HCV antibody prevalence relative to prevalence of reported substance abuse across 14 BBV prevalence studies in the context of SMI. HCV antibody prevalence (blue bars) was categorised in ascending order, and rates of substance abuse (orange bars) charted relative to this. Most studies (57.9%) were drawn from the USA. Both retrospective and prospective studies were included, though the same trend was seen in assessment of retrospective and prospective study designs respectively. <sup>1-14</sup>