

It is illegal to post this copyrighted PDF on any website.

Suicidal Ideation Trends and Associated Factors in Different Large Spanish Samples During the First Year of the COVID-19 Pandemic

Pilar A. Sáiz, PhD^{a,b,c,d,e}; Francesco Dal Santo, MD^{a,c,e}; Leticia García-Alvarez, PhD^{b,c,d,f}; María Teresa Bobes-Bascarán, PhD^{b,c,d,f}; Luis Jiménez-Treviño, PhD^{a,b,c,d,e}; Elisa Seijo-Zazo, PhD^{c,e}; Julia Rodríguez Revuelta, PhD^{a,c,d,e}; Leticia González-Blanco, PhD^{a,b,c,d,e}; María P. García-Portilla, PhD^{a,b,c,d,e,*}; and Julio Bobes, PhD^{a,b,c,d,e}

Data suggest an increased prevalence of depressive disorder and suicidal ideation (SI) (both closely connected with suicidal behavior) associated with the COVID-19 pandemic.¹

The main study objective was to determine the prevalence of passive SI (PSI) and active SI (ASI) in Spanish general population surveys conducted at 3 points in time during the COVID-19 pandemic. We also characterize the main factors associated with ASI.

Methods

Three cross-sectional online surveys, designed to assess the psychological impact of the pandemic in the Spanish population 18 years and older, were conducted April 16–23, 2020 (S1: peak of first wave); October 14–November 8, 2020 (S2: peak of second wave); and March 16–31, 2021 (S3: after third wave). Study procedures were similar to those described elsewhere.^{2,3} Studies were conducted according to the Declaration of Helsinki, using an ad hoc sociodemographic and clinical questionnaire and the Spanish versions of the Depression, Anxiety and Stress Scale (DASS-21)⁴ and Paykel Suicide Scale (PSS).⁵ It should be noted that COVID-19 vaccination status and specific questions about sleep problems, complaints, and satisfaction were included only in S3. PSI was defined as positive answers to PSS items 1 and/or 2, and ASI, as positive answers to PSS items 3 and/or 4. In each survey, the comparison between groups (with and

without ASI) was carried out using an exploratory χ^2 test to establish different groups for categorical variables and a 1-way analysis of variance with a Bonferroni post hoc analysis for continuous variables. Forward stepwise regression models were estimated to determine independent factors associated with “active suicidal ideation during past month” (no/yes). Odds ratios and 95% confidence intervals were calculated. IBM SPSS Version 24.00 (IBM Corp, Armonk, NY) was used for all data analyses, and the *P* value was set at $<.05$.

Results

Sociodemographic and clinical characteristics of the samples are described in Supplementary Table 1. At S1, PSI was detected in 317 respondents (5.2%; mean [SD] age, 41.64 [14.55] years; males vs females: 76 [4.2%] vs 241 [5.6%], *P* = .021) and ASI in 223 respondents (3.7%; age, 39.25 [13.03]; males vs females: 54 [3.0%] vs 169 [3.9%], *P* = .068). At S2, PSI was detected in 1,024 respondents (16.0%; age, 32.03 [10.07]; males vs females: 69 [10.0%] vs 955 [16.7%], *P* < .001) and ASI in 590 respondents (9.2%; age, 31.27 [10.95]; males vs females: 57 [8.3%] vs 533 [9.3%], *P* = .429). At S3, PSI was detected in 590 respondents (10.4%; age, 36.69 [11.95]; males vs females: 55 [5.1%] vs 535 [11.7%], *P* < .001) and ASI in 415 respondents (7.3%; age, 34.89 [11.36]; males vs females: 58 [5.4%] vs 357 [7.8%], *P* < .007). In all surveys, subjects with PSI and ASI were younger than subjects without suicidal ideation (*P* < .001), and no age differences were found between subjects with PSI and those with ASI. Logistic regression models were run to assess variables associated with ASI in all surveys (Table 1).

Discussion

To our knowledge, this study is one of the first to show the evolution of SI in large Spanish population samples during the COVID-19 pandemic. Data show a high prevalence of PSI and ASI at the 3 points in time. In all surveys, the prevalence of prior-month SI was much higher than reported in the adult general population during the 12 months before (2001/2002) and after (2011/2012) the economic crisis⁶ (0.69% and 0.89%, respectively), and even during the very early stage of the pandemic.⁷ However, prior data suggest that only suicide attempts, and not SI, were more common in psychiatric admissions during the COVID-19 pandemic than before.⁸ The large increase in prevalence of both PSI

^aDepartment of Psychiatry, Universidad de Oviedo, Oviedo, Spain

^bCentro de Investigación Biomédica en Red de Salud Mental (CIBERSAM), Spain

^cInstituto de Investigación Sanitaria del Principado de Asturias (ISPA), Oviedo, Spain

^dInstituto Universitario de Neurociencias del Principado de Asturias (INEUROPA), Oviedo, Spain

^eServicio de Salud del Principado de Asturias (SESPE), Oviedo, Spain

^fDepartment of Psychology, Universidad de Oviedo, Oviedo, Spain

*Corresponding author: María P. García-Portilla, PhD, Department of Psychiatry-CIBERSAM, School of Medicine, University of Oviedo, Julián Clavería, s/n, 33006 Oviedo, Spain (albert@uniovi.es).

J Clin Psychiatry 2022;83(2):21br14061

To cite: Sáiz PA, Dal Santo F, García-Alvarez L, et al. Suicidal ideation trends and associated factors in different large Spanish samples during the first year of the COVID-19 pandemic. *J Clin Psychiatry*. 2022;83(2):21br14061.

To share: <https://doi.org/10.4088/JCP.21br14061>

© Copyright 2022 Physicians Postgraduate Press, Inc.

It is illegal to post this copyrighted PDF on any website

Table 1. Factors Associated With Active Suicidal Ideation (Past Month) in Surveys of the Spanish General Population Conducted at 3 Points in Time During the COVID-19 Pandemic

Variable	Survey 1 (April 16–22, 2020)			Survey 2 (October 14–November 8, 2020)			Survey 3 (March 16–31, 2021)		
	β	P value	OR (95% CI)	β	P value	OR (95% CI)	β	P value	OR (95% CI)
Sex (female)				-0.462	.005	0.630 (0.455–0.872) ^a			
Age							-0.027	<.001	0.974 (0.963–0.984) ^a
Educational level									
Primary (reference)					.001				
Secondary				-0.570	.146	0.566 (0.262–1.220)			
Higher				-0.896	.022	0.408 (0.189–0.880) ^b			
Work status									
Unemployed (reference)					<.001				
Employed/retired				-0.630	<.001	0.533 (0.398–0.713) ^b			
Self-employed				-0.178	.436	0.837 (0.534–1.311)			
Other				-0.373	.019	0.688 (0.505–0.939) ^a			
Income (€)									
No income (reference)								.039	
500–1,500							-0.173	.281	0.841 (0.615–1.152)
> 1,500							-0.472	.007	0.624 (0.443–0.879) ^a
Prefer not to answer							-0.335	.265	0.716 (0.398–1.288)
Changes in income due to COVID-19									
No changes (reference)		.013						.002	
Reduction, ≤ 50%	0.324	.074	1.382 (0.969–1.973)				0.303	.039	1.354 (1.016–1.807) ^a
Reduction, 51%–100%	0.636	.003	1.899 (1.245–2.865) ^b				0.656	.002	1.926 (1.267–2.928) ^b
Increase	0.652	.303	1.920 (0.555–6.649)				0.480	.027	1.616 (1.055–2.475) ^a
COVID-19 diagnosis (yes)	0.433	.030	1.542 (1.042–2.283) ^a						
Living situation									
Alone (reference)								<.001	
With 1 other person							-0.662	<.001	0.516 (0.370–0.719) ^b
With more than 1 other person							-0.532	.001	0.588 (0.430–0.803) ^b
Dependent children									
None (reference)					.001				
1				-0.391	.014	0.677 (0.496–0.922) ^a			
More than 1				-0.485	.002	0.616 (0.454–0.835) ^a			
Past history of mental disorder									
No lifetime mental disorder (reference)		<.001			<.001			<.001	
Past mental disorder	0.042	.852	1.043 (0.671–1.620)	0.323	.044	1.382 (1.008–1.893) ^a	0.399	.015	1.490 (1.081–2.053) ^a
Current mental disorder	0.782	<.001	2.186 (1.529–3.124) ^b	1.247	<.001	3.479 (2.785–4.347) ^b	1.160	<.001	3.191 (2.489–4.091) ^b
Personal history of suicide attempt (yes)	1.171	<.001	3.224 (2.158–4.817) ^b	1.104	<.001	3.017 (2.278–3.996) ^b	1.350	<.001	3.856 (2.736–5.434) ^c
Able to enjoy free time (yes)	-0.552	.002	0.576 (0.404–0.821) ^b				-0.370	.001	0.691 (0.557–0.857) ^a
DASS-21 depression (yes)	1.669	<.001	5.469 (3.299–9.065) ^c	1.372	<.001	3.943 (2.819–5.515) ^c			
DASS-21 anxiety (yes)	0.835	<.001	2.305 (1.602–3.315) ^b	0.487	<.001	1.628 (1.285–2.062) ^b			
DASS-21 stress (yes)	0.728	<.001	2.070 (1.445–2.966) ^b	0.666	<.001	1.946 (1.515–2.499) ^b			
Insomnia (yes) ^d							0.604	<.001	1.829 (1.387–2.412) ^b
Sleep satisfaction ^d									
Dissatisfaction (reference)								<.001	
Neutral							-0.405	.004	0.667 (0.505–0.880) ^a
Satisfaction							-0.743	<.001	0.475 (0.351–0.644) ^b
Constant	-5.096	<.001	0.006	-2.910	<.001	0.054	-1.005	.001	0.366
Logistic regression									
Cox and Snell R		.079			.132			.076	
Nagelkerke R ²		.292			.288			.187	
Hosmer-Lemeshow χ ² [df], P		5.729 [6], .454			5.697 [8], .681			14.410 [8], .072	
Correct predictions		96.3%			91.0%			92.7%	

^aCohen *d* equivalence: insignificant (OR < 1.68). ^bCohen *d* equivalence: small (OR = 1.68–3.47). ^cCohen *d* equivalence: medium (OR = 3.48–6.71). ^dInsomnia and sleep satisfaction assessment was included only in Survey 3. Abbreviations: CI = confidence interval; DASS-21 = Depression, Anxiety and Stress Scale; OR = odds ratio.

You are prohibited from making this PDF publicly available.

It is illegal to post this copyrighted PDF on any website.

and ASI, especially during the second and third waves, may be due to the different baseline sample characteristics, but possibly also to psychological fatigue after several months of living with COVID-19 pandemic risks and restrictions.³ Personal history of suicide attempt; current or past history of mental disorder; insomnia; DASS-21 depression, anxiety, or stress; and reduction in income (> 50%) are consistent risk factors for ASI. However, the main protective factor was being able to enjoy free time. Our data suggest that risk and protective factors are dynamic and change slightly depending on the time point and stage of the pandemic.

The main limitations are the cross-sectional online nature of the surveys, which precludes establishing causality, and the use of 3 independent convenience samples, which precludes establishing a rigorous longitudinal SI trend and does not exclude the possibility that the data reflect cohort effects. Further, the included samples are smaller than other samples recently reported. Our data suggest some risk and protective factors associated with ASI during different stages of the COVID-19 pandemic in Spain that could contribute to better adaptation of evidence-based suicide prevention strategies during and after the pandemic.⁹

Published online: March 7, 2021.

Author contributions: Concept and design: García-Alvarez, García-Portilla, Sáiz, and Bobes. Acquisition, analysis, or interpretation of data: All authors. Drafting of the manuscript: Sáiz. Statistical analysis: All authors. Critical revision of the manuscript for important intellectual content: All authors reviewed all drafts and gave final approval.

Potential conflicts of interest: None reported.

Funding/support: This work was partly supported by the Government of the Principality of Asturias PCTI-2018-2022 IDI/2018/235, CIBERSAM, and Fondos Europeos de Desarrollo Regional (FEDER). Financial support was also provided by Fundación para la Investigación e Innovación Biosanitaria del Principado de Asturias (FINBA).

Role of the sponsor: The funding organizations had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, and approval of the manuscript; or decision to submit the manuscript for publication.

Acknowledgment: The authors thank Sharon Grevet for English assistance. Ms Grevet reports no potential conflict of interest.

Supplementary material: Available at PSYCHIATRIST.COM

REFERENCES

1. Czeisler MÉ, Lane RI, Petrosky E, et al. Mental health, substance use, and suicidal ideation during the COVID-19 pandemic—United States, June 24–30, 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69(32):1049–1057.
2. García-Álvarez L, de la Fuente-Tomás L, García-Portilla MP, et al. Early psychological impact of the 2019 coronavirus disease (COVID-19) pandemic and lockdown in a large Spanish sample. *J Glob Health.* 2020;10(2):020505.
3. Ruiz FJ, Sáiz PA, García-Portilla MP, et al. Does COVID-19 psychological fatigue exist? results of three cross-sectional studies conducted in Spain from April 2020 to March 2021. *J Glob Health.* 2021. Accepted on October 22, 2021.
4. Bados A, Solanas A, Andres R. Psychometric properties of the Spanish version of Depression, Anxiety and Stress Scales (DASS). *Psicothema.* 2005;17(4):679–683.
5. Paykel ES, Myers JK, Lindenthal JJ, et al. Suicidal feelings in the general population: a prevalence study. *Br J Psychiatry.* 1974;124(0):460–469.
6. Miret M, Caballero FF, Huerta-Ramírez R, et al. Factors associated with suicidal ideation and attempts in Spain for different age groups. Prevalence before and after the onset of the economic crisis. *J Affect Disord.* 2014;163:1–9.
7. Sáiz PA, de la Fuente-Tomas L, García-Alvarez L, et al. Prevalence of passive suicidal ideation in the early stage of the coronavirus disease 2019 (COVID-19) pandemic and lockdown in a large Spanish sample. *J Clin Psychiatry.* 2020;81(6):2013421.
8. Berardelli I, Sarubbi S, Rogante E, et al. The impact of the COVID-19 pandemic on suicide ideation and suicide attempts in a sample of psychiatric inpatients. *Psychiatry Res.* 2021;303:114072.
9. Wasserman D, Iosue M, Wuestefeld A, et al. Adaptation of evidence-based suicide prevention strategies during and after the COVID-19 pandemic. *World Psychiatry.* 2020;19(3):294–306.

See supplementary material for this brief report at PSYCHIATRIST.COM.



THE JOURNAL OF CLINICAL PSYCHIATRY

THE OFFICIAL JOURNAL OF THE AMERICAN SOCIETY OF CLINICAL PSYCHOPHARMACOLOGY

Supplementary Material

Article Title: Suicidal Ideation Trends and Associated Factors in Different Large Spanish Samples During the First Year of the COVID-19 Pandemic

Authors: Pilar A. Sáiz, PhD; Francesco Dal Santo, MD; Leticia García-Alvarez, PhD; María Teresa Bobes-Bascarán, PhD; Luis Jiménez-Treviño, PhD; Elisa Seijo-Zazo, PhD; Julia Rodríguez Revuelta, PhD; Leticia González-Blanco, PhD; María P. García-Portilla, PhD; and Julio Bobes, PhD

DOI Number: 10.4088/JCP.21br14061

List of Supplementary Material for the article

1. [Table 1](#) Sociodemographic and Clinical Characteristics of the Spanish General Population Samples Included in Three Surveys Conducted at Three Points in Time During the COVID-19 Pandemic

Disclaimer

This Supplementary Material has been provided by the author(s) as an enhancement to the published article. It has been approved by peer review; however, it has undergone neither editing nor formatting by in-house editorial staff. The material is presented in the manner supplied by the author.

Supplementary Table 1. Sociodemographic and clinical characteristics of the Spanish general population samples included in three surveys conducted at three points in time during the COVID-19 pandemic

	Survey 1 (April 16-22, 2020) Total sample (n = 6,108) ¹	Survey 2 (October 14 – November 8, 2020) Total sample (n = 6,418) ²	Survey 3 (March 16-31, 2021) Total sample (n = 5,654) ³	P Value (Statistically different groups)
Age [mean (SD)]	45.78 (14.15)	34.71 (11.74)	39.65 (12.65)	< .001 (1≠2≠3)
Sex (female) [n(%)]	4,280 (70.1)	5,731 (89.3)	4,575 (80.9)	< .001 (1≠2≠3)
Marital status [n(%)]				
Never married	2,080 (34.1)	3,488 (54.3)	2,351 (41.6)	< .001 (1≠2≠3)
Married / Living as married	3,365 (55.1)	2,645 (41.2)	2,947 (52.1)	< .001 (1≠2 & 2≠3); .001 (1≠3)
Separated / Divorced / Widowed	663 (10.9)	285 (4.4)	356 (6.3)	< .001 (1≠2≠3)
Educational level [n(%)]				
Primary	99 (1.6)	53 (0.8)	36 (0.6)	< .001 (1≠2 & 1≠3)
Secondary	1,763 (28.9)	1,983 (30.9)	1,634 (28.9)	.014 (1≠2); .018 (2≠3)
Higher	4,246 (69.5)	4,382 (68.3)	3,984 (70.5)	.010 (2≠3)
Work status [n(%)]				
Unemployed	518 (8.5)	506 (7.9)	358 (6.3)	< .001 (1≠3); .001 (2≠3)
Employed / Retired	2,748 (45.0)	3,954 (61.6)	3,906 (69.1)	< .001 (1≠2≠3)
Self-employed	2,098 (34.3)	422 (6.6)	455 (8.0)	< .001 (1≠2 & 1≠3); .002 (2≠3)
Other	744 (12.2)	1,536 (23.9)	935 (16.5)	< .001 (1≠2≠3)
Income (€) [n(%)]				
No income	592 (9.7)	1,293 (20.1)	784 (13.9)	< .001 (1≠2≠3)
500-1500	2,008 (32.9)	2,343 (36.5)	1,855 (32.8)	< .001 (1≠2 & 2≠3)
> 1500	3,038 (49.7)	2,376 (37.0)	2,729 (48.3)	< .001 (1≠2 & 2≠3)
Prefer not to answer	470 (7.7)	406 (6.3)	286 (5.1)	.003 (1≠2 & 2≠3); < .001 (1≠3)
Changes in income due to COVID-19 [n(%)]				
No changes	4,334 (71.0)	4,602 (71.7)	4,279 (75.7)	< .001 (1≠2≠3)
Reduction, ≤ 50%	1,182 (19.4)	1,256 (19.6)	861 (15.2)	< .001 (1≠2≠3)
Reduction, 51-100%	533 (8.7)	341 (5.3)	201 (3.6)	< .001 (1≠2≠3)
Increase	59 (1.0)	219 (3.4)	313 (5.5)	< .001 (1≠2≠3)
Living situation [n(%)]				
Alone	783 (12.8)	643 (10.0)	620 (11.0)	< .001 (1≠2); .002 (1≠3)
With one other person	2,325 (38.1)	2,219 (34.6)	2,016 (35.7)	< .001 (1≠2); .007 (1≠3)
With more than one	3,000 (49.1)	3,556 (55.4)	3,018 (53.4)	< .001 (1≠2 & 1≠3); < .027 (2≠3)
Dependent children [n(%)]				
None	3,789 (62.0)	4,441 (69.2)	3,393 (60.0)	< .001 (1≠2 & 2≠3); .026 (1≠3)
One	1,092 (17.9)	878 (13.7)	926 (16.4)	< .001 (1≠2 & 2≠3); .033 (1≠3)
More than one	1,227 (20.1)	1,099 (17.1)	1,335 (23.6)	< .001 (1≠2≠3)
Elderly dependents [n(%)]				

None	5,428 (88.9)	5,842 (91.0)	4,964 (87.8)	< .001 (1#2 & 2#3)
One	523 (8.6)	400 (6.2)	472 (8.3)	< .001 (1#2 & 2#3)
Two or more	157 (2.6)	176 (2.7)	218 (3.9)	< .001 (1#3); .001 (2#3)
Chronic physical disease (Yes) [n(%)]	1,295 (21.2)	1,081 (16.8)	1,279 (22.6)	< .001 (1#2 & 2#3)
Diagnosed with COVID-19 disease (Yes) [n(%)]	566 (9.3)	1,013 (15.8)	696 (12.3)	< .001 (1#2#3)
Family / Friends diagnosed with COVID-19 disease (Yes) [n(%)]	2,398 (39.3)	4,195 (65.4)	-	< .001 (1#2)
Family / Friends died of COVID-19 (Yes) [n(%)]	705 (11.5)	819 (12.8)	1,272 (22.5)	.040 (1#2); < .001 (1#3 & 2#3)
Frontline worker [n(%)]*				
No	-	3,836 (59.8)	3,231 (57.1)	.004 (2#3)
Healthcare worker	-	1,575 (24.5)	1,316 (23.3)	NS
First responder	-	28 (0.4)	254 (4.5)	< .001 (2#3)
Other	-	979 (15.3)	853 (15.1)	NS
Past history of mental disorder [n(%)]				
No lifetime mental disorder	4,812 (78.8)	4,014 (62.5)	4,344 (76.8)	< .001 (1#2 & 2#3); .012 (1#3)
Past mental disorder	786 (12.9)	914 (14.2)	606 (10.7)	< .001 (1#3 & 2#3); .027 (1#2)
Current mental disorder	510 (8.3)	1,490 (23.2)	704 (12.5)	< .001 (1#2#3)
Personal history of suicide attempt (Yes) [n(%)]	264 (4.3)	329 (5.1)	215 (3.8)	< .001 (2#3); .038 (1#2)
Able to enjoy free time (Yes) [n(%)]	5,690 (93.2)	5,453 (85.0)	3,353 (59.3)	< .001 (1#2#3)
DASS-21 Depression (Yes) [n(%)]	2,523 (41.3)	3,623 (56.5)	2,552 (45.1)	< .001 (1#2#3)
DASS-21 Anxiety (Yes) [n(%)]	1,056 (17.3)	2,374 (37.0)	2,443 (43.2)	< .001 (1#2#3)
DASS-21 Stress (Yes) [n(%)]	1,151 (18.8)	2,634 (41.0)	2,440 (43.2)	< .001 (1#2 & 1#3); .020 (2#3)
Insomnia (Yes) [n(%)]**	-	-	2,320 (41.0)	-
Sleep complaints (Yes) [n(%)]**	-	-	3,456 (61.1)	-
Sleep satisfaction [n(%)]**				
Dissatisfied	-	-	1,523 (26.9)	-
Neutral	-	-	1,218 (21.5)	-
Satisfied	-	-	2,913 (51.5)	-

*Family / Friends diagnosed with COVID-19 disease was assessed only in Surveys 1 and 2; **Sleep problems, complaints, and satisfaction assessment was included only in Survey 3.

Abbreviations: DASS-21: Depression, Anxiety and Stress Scale; NS: Not statistically significant