

It is illegal to post this copyrighted PDF on any website.

You are prohibited from making this PDF publicly available.

**CME Background Information**

Articles are selected for credit designation based on an assessment of the educational needs of CME participants, with the purpose of providing readers with a curriculum of CME articles on a variety of topics throughout each volume. Activities are planned using a process that links identified needs with desired results.

To obtain credit, read the article, correctly answer the questions in the Posttest, and complete the Evaluation.

This commentary was prepared and independently developed by the CME Institute of Physicians Postgraduate Press, Inc., and supported by an educational grant from Alexion Pharmaceuticals, Inc. Dr Brown acknowledges Michael R. Page, PharmD, RPh, for editorial assistance in developing the manuscript.

**Robert S. Brown, Jr, MD, MPH**, is affiliated with Weill Cornell Medicine, New York, New York.

**CME Objective**

After completing this educational activity, you should be able to:

- Implement an individualized treatment strategy for the patient with Wilson disease, while watching for new options that could resolve unmet needs

**Accreditation Statement**

The CME Institute of Physicians Postgraduate Press, Inc., is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.



**Release, Review, and Expiration Dates**

This Neuroscience Commentary activity was published in November 2022 and is eligible for *AMA PRA Category 1 Credit* through December 31, 2023. The latest review of this material was June 2022.

**Financial Disclosure**

The CME Institute adheres to the Standards for Integrity and Independence in Accredited Continuing Education of the Accreditation Council for Continuing Medical Education (ACCME). Any individuals in a position to control the content of a continuing education activity, including faculty, content developers, reviewers, staff, and others, are required to disclose to learners the presence or absence of any relevant financial relationships with an ACCME-defined ineligible company within the preceding 24 months of the activity. The ACCME defines an “ineligible company” as one whose primary business is producing, marketing, selling, re-selling, or distributing healthcare products used by or on patients.

The CME Institute has mitigated all relevant conflicts of interest prior to the commencement of the activity. None of the individuals involved in the content have relevant financial relationships with ineligible companies except the following:

Larry Culpepper, MD, MPH, Editor in Chief, Boston, Massachusetts, has been an advisor for AbbVie, Eisai, and Supernus; has been a stock shareholder of M-3 Information; and has received royalties from UpToDate. Michael R. Page, PharmD, RPh, Independent Medical Director/Medical Writer, Plainsboro, New Jersey, serves as a consultant for BioCentric, Inc., and American Medical Communications, Inc.

None of the other planners, reviewers, and CME Institute staff for this educational activity have relevant financial relationships with ineligible companies to disclose. All relevant financial relationships have been mitigated. **Dr Brown’s financial disclosure appears on the next page.**

*Prim Care Companion CNS Disord* 2022;24(6):AN21065COM1C

**To cite:** Brown RS Jr. An update on the future of Wilson disease management. *Prim Care Companion CNS Disord.* 2022;24(6):AN21065COM1C.

**To share:** <https://doi.org/10.4088/PCC.AN21065COM1C>

© Copyright 2022 Physicians Postgraduate Press, Inc.

# An Update on the Future of Wilson Disease Management

Robert S. Brown, Jr, MD, MPH

**W**ilson disease (WD) is a rare autosomal recessive genetic disorder of copper metabolism. Although WD was first described as progressive lenticular degeneration in 1912,<sup>1</sup> and established as a condition related to elevated copper levels in 1948,<sup>2</sup> it was not until 1956 that the copper chelator penicillamine was introduced<sup>3</sup> and 1993 that mutations in the *ATP7B* gene were implicated as the causative underlying genetic abnormality.<sup>4</sup> With a broad range of symptoms, including dysarthria, dystonia, tremor, chorea, athetosis, and, in later stages of disease, hepatic failure and neurologic disability, the differential diagnosis of WD may be challenging, leading to delays in initiation of effective therapy.<sup>5</sup> Although heterozygous mutations of *ATP7B* may be present in up to 2.5% of the population,<sup>6</sup> clinical disease manifests in approximately 1 in 30,000 to 100,000 individuals.<sup>7</sup> WD usually presents in childhood into early adulthood; however, a wider range of ages at onset is recognized.<sup>8</sup> While it is extremely rare for WD to present after the age of 35, late-onset cases are reported. Given the myriad manifestations of WD, its rarity, and its diversity of presentations, diagnostic acumen and a knowledge of the latest updates in the field, as well as novel therapies in late-stage development, are crucial in ensuring appropriate management.

From a pathophysiologic perspective, WD is caused by dysregulation of the tightly regulated system of transport, retention, and excretion of copper from the diet. Of the roughly 2 to 5 mg of dietary copper ingested daily by an average individual, approximately 2% is retained in the body for use in enzymatic activity, and the remaining portion is excreted in bile.<sup>9</sup> This process involves multiple transporters. Copper is first absorbed within the small intestines through nonspecific metal uptake transporters, exported from the small intestine to the portal circulation via P-type ATPase copper transporters, and absorbed from portal circulation to the liver via the *CTR1* transporter. Within the liver, the *ATP7B* transporter regulates incorporation of copper into the carrier protein ceruloplasmin, as well as excretion of copper through the biliary system.<sup>9</sup> In patients with *ATP7B* mutations, this crucial transport function is impaired, resulting in deficient levels of ceruloplasmin and an accumulation of unbound copper within the liver that ultimately reaches other vital organs, including the brain, eyes, and kidney, resulting in clinical signs and symptoms.<sup>10</sup>



**Case Practice Question**

*Discussion of best response can be found at the end of the activity.*

Charles is a 12-year-old male with suspected Wilson disease presenting with ataxia and other neurologic symptoms. Which of the following features would be suggestive of Wilson disease in this patient?

- a. Elevated levels of ceruloplasmin
- b. The patient cannot have Wilson disease, as cases usually present after age 40
- c. Mutations in the *ATP7B* gene
- d. Wilson disease can be ruled out due to the lack of liver disease

**Credit Designation**

The CME Institute of Physicians Postgraduate Press, Inc., designates this enduring material for a maximum of 0.5 *AMA PRA Category 1 Credit*. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

*Note:* The American Nurses Credentialing Center (ANCC) and the American Academy of Physician Assistants (AAPA) accept certificates of participation for educational activities certified for *AMA PRA Category 1 Credit* from organizations accredited by the ACCME.

**Financial Disclosure**

**Dr Brown** has served as a consultant for Alexion, AbbVie, Gilead, Intercept, Mallinckrodt, Antios, and Ambys and has received grant/research support from AbbVie, Gilead, Intercept, and Mallinckrodt.

**Review Process**

The author agreed to provide a balanced and evidence-based presentation and discussed the topic and CME objective during the planning sessions. The author's submitted content was validated by CME Institute staff, and the activity was evaluated for accuracy, use of evidence, and fair balance by a peer reviewer who is without conflict of interest.

The opinions expressed herein are those of the faculty and do not necessarily reflect the opinions of the CME provider and publisher or Alexion Pharmaceuticals, Inc.

The management of WD has evolved over the years. Current guidelines<sup>10</sup> for diagnosis and treatment of WD in adults published by the American Association for the Study of Liver Diseases in 2008 and pediatric guidance<sup>11</sup> published in 2018 recommend use of chelation therapies to decrease total body copper stores as well as zinc to reduce intestinal copper absorption. While penicillamine therapy remains a standard treatment for WD,<sup>11</sup> trientine is becoming a preferred first-line treatment for WD due to lower rates of side effects and superior tolerability compared with penicillamine.<sup>12</sup> Another chelating agent, tetrathiomolybdate (TTM), was not widely available at the time of publication due to chemical instability of the ammonium salt used initially for WD treatment<sup>12</sup> but is mentioned in guidelines<sup>10</sup> as an experimental decoppering therapy in the United States and Canada. However, a more stable form of this potent and multimodal decoppering agent, bis-choline TTM, has been developed and promising results were reported in a phase 2 clinical trial.<sup>13</sup> A phase 3 clinical study<sup>14</sup> is underway in patients with WD with a 48-week double-blind phase and an extension period of up to 60 months.

You are prohibited from making this PDF publicly available.

**Table 1. Common Therapies Used in the Treatment of Wilson Disease<sup>a</sup>**

Therapy	Year of introduction	Mechanism of action	Summary of efficacy and adverse events leading to treatment discontinuation
Penicillamine	1956	Copper chelation	<ul style="list-style-type: none"> <li>• Promotes copper urinary excretion of 200-500 µg per 24-hour period in the maintenance phase, with excretion rates exceeding 1000 µg per 24-hour period in the initial phase</li> <li>• AEs are reported in 20% to 30% of patients during treatment and include "early" AEs (eg, fever, cutaneous manifestations, arthralgia) and "late" manifestations (eg, paradoxical neurological worsening, renal insufficiency, lupus-like syndrome)</li> </ul>
Zinc salts	1961	Decreased GI absorption of copper	<ul style="list-style-type: none"> <li>• Copper urinary excretion &lt; 75 µg per 24-hour period, with NCC levels ranging from 5-15 µg/dL (&gt; 12 months of treatment)</li> <li>• AEs reported in 3% to 7% of patients, with the most common being gastritis, biochemical pancreatitis, immunosuppression, and bone marrow depression</li> </ul>
Trientine	1982	Copper chelation	<ul style="list-style-type: none"> <li>• Promotes copper urinary excretion of 200-500 µg per 24-hour period in the maintenance phase, with excretion rates exceeding 1000 µg per 24-hour period in the initial phase</li> <li>• AEs in 7.1% of patients, with the most common being gastritis, sideroblastic anemia, lupus-like reactions, and loss of taste</li> </ul>
Bis-choline tetrathiomolybdate	In phase 3 evaluation as of 2022	Copper chelation and decreased GI absorption of copper	<ul style="list-style-type: none"> <li>• On a primary endpoint of change in baseline NCC concentration at 24 weeks and with maintenance of normalized NCC levels of 2.3 µmol/L or lower, or at least a 25% reduction from baseline, 71% of patients met criteria for treatment success. Mean NCC levels were reduced by 72% from baseline to week 24 (<i>P</i> &lt; .0001)</li> <li>• AEs were reported (psychiatric disorders, gait disturbance, elevated liver aminotransferases, and decline in neurological functioning), although only 4 of 11 AEs observed were considered possibly or probably related to study medication</li> </ul>

Abbreviations: AE = adverse event, GI = gastrointestinal, NCC = non-ceruloplasmin-bound copper.

<sup>a</sup>Based on Czlonkowska et al<sup>6</sup> and Weiss et al.<sup>13</sup>

**It is illegal to post this copyrighted PDF on any website.**

In a systematic review and meta-analysis<sup>15</sup> rating the comparative effectiveness and safety of common WD therapies, although D-penicillamine was associated with lower mortality versus no treatment, the drug was not associated with a lower risk of mortality and improved rates of prevention or amelioration of clinical symptoms compared with zinc alone. In a 10-year follow-up study of 22 children with WD, treatment with zinc sulfate adjusted by age and weight, starting in presymptomatic pediatric patients, resulted in 73% of children (16/22) having normal alanine aminotransferase levels, concurrent with increased urinary copper excretion.<sup>16</sup> A randomized controlled trial comparing trientine hydrochloride plus zinc with TTM plus zinc in primarily newly diagnosed patients with Wilson disease with neurologic symptoms identified a higher risk of neurologic deterioration in patients receiving trientine versus TTM (6 of 23 patients [26%] vs 1 of 25 patients [4%],  $P < .05$ ).<sup>17</sup> Treatment with another form of trientine—trientine tetrahydrochloride—has also been trialed, with noninferiority to D-penicillamine over 24 weeks of follow-up established in terms of reductions in non-ceruloplasmin copper levels.<sup>18</sup> Clinical efficacy and safety of current therapies and the latest data on bis-choline TTM are reviewed in Table 1.<sup>6,13</sup>

In the modern management of WD, consideration of the latest evidence in both diagnosis and long-term treatment are important in optimizing clinical management. In 2022, clinicians should be aware of the role of copper and mutations of the *ATP7B* gene in disease pathophysiology, as well as the clinical manifestations that may raise clinical suspicion of this rare disease. Through an awareness of key clinical data and novel therapies in late-stage development, clinicians can screen for and maintain awareness of the latest multimodal mechanisms for the management of WD.

**Published online:** November 29, 2022.

**Unlabeled and investigational usage:** The faculty of this educational activity may include discussions of products or devices that are not currently labeled for use by the FDA. Faculty members have been advised to disclose to the audience any reference to an unlabeled or investigational use. No endorsement of unapproved products or uses is made or implied by coverage of these products or uses. Please refer to the official prescribing information for each product for discussion of approved indicators, contraindications and warnings.

**REFERENCES**

- Wilson S. Progressive lenticular degeneration: a familial nervous disease associated with cirrhosis of the liver. *Brain*. 1912;34(4):295–507.
- Cumings JN. The copper and iron content of brain and liver in the normal and in hepato-lenticular degeneration. *Brain*. 1948;71(Pt. 4):410–415.
- Walshe JM. Penicillamine, a new oral therapy for Wilson’s disease. *Am J Med*. 1956;21(4):487–495.
- Bull PC, Thomas GR, Rommens JM, et al. The Wilson disease gene is a putative copper transporting P-type ATPase similar to the Menkes gene. *Nat Genet*. 1993;5(4):327–337.
- Patil M, Sheth KA, Krishnamurthy AC, et al. A review and current perspective on Wilson disease. *J Clin Exp Hepatol*. 2013;3(4):321–336.
- Czlonkowska A, Litwin T, Dusek P, et al. Wilson disease. *Nat Rev Dis Primers*. 2018;4(1):21.
- Kalita J, Kumar V, Chandra S, et al. Worsening of Wilson disease following penicillamine therapy. *Eur Neurol*. 2014;71(3-4):126–131.
- Lorincz MT. Neurologic Wilson’s disease. *Ann N Y Acad Sci*. 2010;1184(1):173–187.
- Kathawala M, Hirschfield GM. Insights into the management of Wilson’s disease. *Therap Adv Gastroenterol*. 2017;10(11):889–905.
- Roberts EA, Schilsky ML; American Association for Study of Liver Diseases (AASLD). Diagnosis and treatment of Wilson disease: an update. *Hepatology*. 2008;47(6):2089–2111.
- Socha P, Janczyk W, Dhawan A, et al. Wilson’s disease in children: a position paper by the hepatology committee of the European Society for paediatric gastroenterology, hepatology and nutrition. *J Pediatr Gastroenterol Nutr*. 2018;66(2):334–344.
- Moini M, To U, Schilsky ML. Recent advances in Wilson disease. *Transl Gastroenterol Hepatol*. 2021;6:21.
- Weiss KH, Askari FK, Czlonkowska A, et al. Bis-choline tetrathiomolybdate in patients with Wilson’s disease: an open-label, multicentre, phase 2 study. *Lancet Gastroenterol Hepatol*. 2017;2(12):869–876.
- A Phase 3, Randomized, Rater-Blinded, Multi-Center Study to Evaluate the Efficacy and Safety of ALXN1840 Administered for 48 Weeks Versus Standard of Care in Patients With Wilson Disease Aged 12 Years and Older With an Extension Period of up to 60 Months. *ClinicalTrials.gov* website. <https://clinicaltrials.gov/ct2/show/NCT03403205>. Published February 15, 2018. Accessed March 2, 2022.
- Appenzeller-Herzog C, Mathes T, Heeres MLS, et al. Comparative effectiveness of common therapies for Wilson disease: A systematic review and meta-analysis of controlled studies. *Liver Int*. 2019;39(11):2136–2152.
- Marcellini M, Di Ciommo V, Callea F, et al. Treatment of Wilson’s disease with zinc from the time of diagnosis in pediatric patients: a single-hospital, 10-year follow-up study. [published correction appears in *J Lab Clin Med*. 2005 Jul;146(1):44. Carelli, Francesco [corrected to Carelli, Giovanni]. *J Lab Clin Med*. 2005;145(3):139–143.
- Schilsky ML, Cassiman D, Ott P, et al. Trientine tetrahydrochloride versus d-penicillamine for the treatment of patients with Wilson disease. AASLD Oral Abstracts website. <https://aasldpubs.onlinelibrary.wiley.com/doi/10.1002/hep.32187?af=R>. Published October 14, 2021. Accessed July 18, 2022.
- Brewer GJ, Askari F, Lorincz MT, et al. Treatment of Wilson disease with ammonium tetrathiomolybdate, IV: comparison of tetrathiomolybdate and trientine in a double-blind study of treatment of the neurologic presentation of Wilson disease. *Arch Neurol*. 2006;63(4):521–527.

You are prohibited from making this PDF publicly available.



**Discussion of Case Practice Question**

**Preferred response: c. Mutations in the *ATP7B* gene**

Wilson disease symptoms include dysarthria, dystonia, tremor, chorea, athetosis, and ataxia and usually presents in childhood into early adulthood. Heterozygous mutations of *ATP7B* gene are implicated as the causative underlying genetic abnormality, causing impairment in *ATP7B* transport function which regulates incorporation of copper into the carrier protein ceruloplasmin as well as excretion of copper through the biliary system. This results in deficient levels of ceruloplasmin and an accumulation of unbound copper within the liver that ultimately reaches other vital organs.

For the CME Posttest, see next page.

It is illegal to post this copyrighted PDF on any website.



## CME INSTITUTE POSTTEST

To obtain credit, go to [CMEInstitute.com](https://www.cmeinstitute.com) to complete the Posttest and Evaluation.

1. Charles is referred for therapy with trientine plus zinc salts but does not experience regression of neurologic symptoms following 8 weeks of treatment. Which of the following therapies might be an alternative in clinical trials?
  - a. Bis-choline tetrathiomolybdate
  - b. Tetrathiomolybdate
  - c. Addition of D-penicillamine
  - d. British anti-Lewisite
2. Which of the following treatments for Wilson disease uses both copper chelation and decreased GI absorption of copper as mechanisms of action and has resulted in a decrease in non-ceruloplasmin-bound copper concentration by 72% from baseline over the course of 2 years?
  - a. Trientine tetrahydrochloride
  - b. Zinc sulfate
  - c. Tetrathiomolybdate
  - d. Bis-choline tetrathiomolybdate

Scan this  
QR code  
to access.



You are prohibited from making this PDF publicly available.