

Table 1. Performance on Cognitive Tasks by Fibromyalgia Patients, Age-Matched Controls, and Older Controls^{a,b}

Task	Fibromyalgia Patients (N = 23)	Age-Matched Controls (N = 23)	Older Controls (N = 22)	p Value ^c
Information-processing speed	139.45 ± 29.55	139.23 ± 29.55	118.50 ± 19.15	.975
Working memory	22.22 ± 7.85	26.30 ± 1.67	22.09 ± 6.27	.042
Free recall	23.56 ± 7.80	27.83 ± 6.43	23.91 ± 6.77	.005
Recognition memory	2.53 ± 1.19	2.95 ± 1.07	2.80 ± 1.19	.035
Verbal fluency	49.78 ± 11.63	56.08 ± 15.65	49.43 ± 13.74	.055
Verbal knowledge	43.17 ± 7.62	51.26 ± 6.01	50.56 ± 7.93	< .001

^aAdapted with permission from Park et al.⁹

^bValues shown as mean ± SD.

^cp Values are for fibromyalgia patients compared with age-matched controls.

memory that are troublesome for fibromyalgia patients are working (short-term) memory, episodic long-term memory, and semantic memory access. Short-term memory refers to memory of less than about 30 seconds, and working memory combines short-term memory with other mental processes. Mental arithmetic is an example of a task that makes heavy use of the working memory system. Episodic long-term memory is the ability to remember particular episodes, such as items on a grocery list or personal experiences. Semantic memory involves facts and knowledge about the world. Also problematic for fibromyalgia patients is the ability to pay attention to things that are important while ignoring distracting items.

Objective Tests of Memory

Working memory. One objective test of working memory is the Reading Span Test,⁸ which colleagues and I used in a study of patients with fibromyalgia.⁹ In this test, the experimenter reads a sentence aloud, for example, "The fans at the football game wore hats and scarves." On a computer screen, the participant sees a question—in this case, "Where were the fans?" Then the participant selects one of the multiple choice answers. Next, the experimenter reads another sentence, for example, "The Grants told their guests from France about the city." Again, the participant selects a multiple choice answer to a question about the sentence. At the end of the trial, the subject is prompted to recall the last word of each sentence in the order in which the words appeared, so in this case, the participant would need to remember *scarves* and *city*. This task is challenging because it includes a lot of competing information that requires storage and manipulation. This test starts with 2 sentences per trial and moves up to 3 sentences and more until the participant can no longer successfully recall all the words. When the patients with fibromyalgia were compared with education-matched controls who were either age-matched or 20 years older, the fibromyalgia patients performed like the older controls, less well ($p = .042$) than the age-matched controls (Table 1).

A standardized test of working memory is the Paced Auditory Serial-Addition Test (PASAT).¹⁰ This test measures memory competence when a source of distraction

is introduced. A tape recorder is normally used to present digits one at a time and the subject's task is to add the most recent digit to the one that preceded it and state the answer aloud. Participants have to continually update the digit they are holding in working memory, which is very challenging. Although Suhr¹¹ found no between-group differences on PASAT scores in a study of patients with fibromyalgia or chronic pain versus healthy controls, Leavitt and Katz¹² found that fibromyalgia patients performed at a level below normal on this task. Furthermore, using the Auditory Consonant Trigram,¹³ Leavitt and Katz¹² found impairment in 82.6% of fibromyalgia patients.

Episodic long-term memory (free recall). Colleagues and I also investigated episodic long-term memory in patients with fibromyalgia compared with age-matched controls and controls who were 20 years older.⁹ A word-list, free recall test¹⁴ was used, in which subjects are presented with a list of words to learn and then about 5 minutes later are asked to recall the words. Participants are allowed to recall the words in any order. As shown in Table 1, the fibromyalgia patients performed like older controls, at a significantly ($p = .005$) lower level than age-matched controls.

Semantic memory (verbal fluency). A standard verbal fluency test can be used to test semantic memory. In the study of patients with fibromyalgia compared with older controls and age-matched controls,⁹ colleagues and I used a standard verbal fluency test¹⁵ in which participants are given paper with the letters *f*, *a*, and *s* at the top and in 90 seconds must write as many words as possible that start with each letter. Again, as shown in Table 1, the fibromyalgia patients performed like the older controls, at a lower level than the age-matched controls.

These results show impairment in working memory, episodic memory, and semantic memory access in fibromyalgia patients that mimics about 20 years of aging.

Objective Tests of Attention

Results of attention tests in patients with fibromyalgia are consistent with the working memory results already discussed because the attention tests require people to store some information for a short period of time and also process distracting information.

Dick and colleagues⁷ studied attentional functioning in patients with fibromyalgia using the Test of Everyday Attention,¹⁶ a computerized battery that is meant to mimic the kinds of things that people do in everyday life. They found that fibromyalgia patients scored significantly ($p < .005$) worse than pain-free controls in selective attention and auditory-verbal working memory.

Leavitt and Katz¹² suggested that the memory difficulties seen in fibromyalgia patients may be due to attentional distraction. They found that when patients with fibromyalgia were taking standardized tests of memory function, higher levels of impairment were observed for tasks that involved distraction.

Colleagues and I conducted a study¹⁷ in which we specifically looked at distraction during memory performance. We asked participants to memorize a list of 24 words, and sometimes while they were memorizing and/or recalling a list of words, they also had to perform a "distractor" task. The distractor task was a simple digit monitoring test in which the participant held a counter in his or her nondominant hand and listened to a tape recording of digits. Half of the participants were asked to press the counter whenever they heard an odd digit and the other half of the subjects were asked to press the counter whenever they heard an even digit.

For one list of words there was full attention at learning and at recall.¹⁷ For another list, attention was divided while learning the list but not while recalling the words. For a third word list, attention was divided while recalling the list but not while learning it. For a final list of words, attention was divided while learning and while recalling the list. The overall results showed that healthy controls on average recalled 15.7 words, whereas the fibromyalgia patients recalled on average 13.3 words ($p = .006$). Dividing attention had costs for both controls and fibromyalgia patients, but the greatest cost incurred was among the fibromyalgia patients when attention was divided during both learning and recall.

These results may explain why fibromyalgia patients report poor memory performance in real-world environments. Distracting information competes for attention, people rarely get a chance to concentrate on one thing at a time, and memory performance is therefore highly disrupted.

Task switching/selective attention. My colleagues and I¹⁸ studied the ability of fibromyalgia patients to switch attention between different aspects of a task. We used a computerized version of a card sorting task in which a card appears at the bottom of a screen and, above the card, 4 piles of cards appear in a row and the top card on each pile has a different shape on it. The size, shading, and number of shapes can differ. The participant has to press keys on the keyboard to indicate which pile the card at the bottom of the screen should join. The rules for sorting the cards can be simple, for example, by shape, size, shading,

or number of objects on the card, or the rules can be complex so that participants have to put 2 variables together. The same rules may apply throughout the whole test, or participants may have to switch back and forth between rules. The data showed that the cost in accuracy was greatest when fibromyalgia patients had to switch between complex rules.

These results confirm patients' accounts that they are most aware of cognitive problems when tasks are difficult, multiple tasks are required, and there is a lot of distraction. Overall, the results of these objective tests of memory and attention suggest that fibromyalgia patients have lower memory performance and are more susceptible to distraction than those without fibromyalgia.

FACTORS THAT CONTRIBUTE TO FIBROFOG

Several factors may contribute to cognitive problems in patients with fibromyalgia. These potential causes include comorbid depression and anxiety, sleep problems, neuroendocrine abnormalities, pain, and central augmentation of sensory input.

Psychiatric Comorbidity

Symptoms of depression and anxiety frequently occur in fibromyalgia patients.¹¹ Depression and anxiety have been associated with lowered memory performance.^{19,20} Colleagues and I⁹ did not find a correlation between depression and anxiety symptoms and results on cognitive tests, but other researchers^{11,21,22} have reported a correlation between psychiatric symptoms and cognitive function in patients with fibromyalgia. Depression or anxiety could be related to cognitive function in some fibromyalgia patients, but these symptoms do not explain all of their memory and attention problems.

Sleep Problems

Fibromyalgia patients usually have sleep problems.^{1,23} In healthy subjects, insufficient sleep has been found to impair learning and memory.^{24,25} Colleagues and I included some measures of sleep when studying patients with fibromyalgia, but we did not find any correlation with cognitive performance. As is the case with depression, poor sleep undoubtedly contributes to some of the cognitive difficulties of fibromyalgia patients but does not tell the whole story.

Neuroendocrine Abnormalities

Neuroendocrine disturbances or abnormalities and sensitivity to stress occur in fibromyalgia patients.²⁶ Stress-level cortisol doses have been shown to decrease memory function in healthy adults,²⁷ and elevated levels of cortisol were associated with lower performance on cognitive tests in those with Cushing's disease.²⁸ However, in a study by Sephton and colleagues,²⁹ higher salivary cortisol levels

correlated with better visual-spatial memory, the opposite of other findings. Neuroendocrine function does seem to have some effect on memory function, but exactly how this may affect fibromyalgia patients is still unclear.

Pain

Fibromyalgia patients experience pain, which disrupts cognition, especially attention, and may contribute to fibro-fog. Painful stimulation is attention-demanding and activates brain areas associated with cognitive attention. Studies³⁰ in different chronic pain patient groups have shown that pain is related to cognitive function. Colleagues and I⁹ found that although scores on the McGill Pain Questionnaire³¹ were not related to cognitive function, scores on the Arthritis Impact Measurement Scales³² pain subscale were. Correlations have been shown between cognitive function and both self-reported and evoked pain in fibromyalgia,³³ so it seems likely that pain contributes to the cognitive problems experienced by fibromyalgia patients.

Central Augmentation of Sensory Input

Evidence suggests that almost all patients with fibromyalgia have central processing abnormalities³⁴; central augmentation of sensory input may interfere with other activities involved in processing sensory input and the ability to ignore items that are distracting and maintain focus on what is important. Gracely and colleagues³⁵ used functional magnetic resonance imaging (fMRI) to examine the pattern of cerebral activation during the application of painful pressure in patients with fibromyalgia compared with controls and found central nervous system augmentation of pain. The fMRI results showed that when similar levels of moderate pressure were applied to the patients and the controls, no common regions of activation were seen and greater effect was seen in patients. When the stimulation was increased to deliver a subjective level of pain in the control group similar to that experienced by fibromyalgia patients, similar activation patterns were seen in patients and controls. Enhanced processing of sensory signals has implications for attention in fibromyalgia because it may lead to increased distractibility.

FEASIBILITY OF TESTING VERSUS PATIENT SELF-REPORT IN CLINICAL PRACTICE

Physicians often ask how cognitive function in fibromyalgia patients should be measured in clinical practice. Generally, neuropsychological and cognitive tests are extensive and time-consuming, which may not be feasible in practice. Many of the tests that are sensitive to the problems that fibromyalgia patients experience are not standardized, and some of the standard neuropsychological tests may miss the attention difficulties of fibromyalgia patients. No quick and easy test of cognitive function in fibromyalgia patients exists as yet.

In the clinic, self-report may be the best and quickest measure of cognitive problems in fibromyalgia patients. Using the MIA questionnaire,⁵ colleagues and I⁴ found that performance on an objective memory recall task correlated significantly with fibromyalgia patients' perceived capacity ($p < .01$), achievement motivation ($p < .05$), and self-efficacy ($p < .01$), but no significant correlations were found between the recall task performance and the perceptions about memory in either control group.

The high correlation between objective and subjective measures of cognitive function that my colleagues and I found⁴ is exceptional. Typically, in all kinds of groups, including healthy young adults, self-reported cognitive function is not correlated well with actual cognitive function. Lack of specificity between the self-report measure and the specific performance measure used frequently accounts for the lack of correspondence. Lack of correspondence may also occur when people with high ability are tested because these individuals may be more sensitive to changes in cognitive function than those of lesser ability. High-ability patients may report a large memory loss when they still appear to perform well on a memory test. Additionally, self-report of cognitive problems may track effort instead of actual performance.³⁶ A patient may perform at normal levels, but has to put a great deal of effort into attaining that performance. Future neuroimaging studies could examine whether brain activity correlates well with self-report of task difficulty.

CONCLUSION

Research using objective tests has verified fibromyalgia patients' self-report of cognitive difficulties. As no simple and quick tests exist for cognitive problems in fibromyalgia patients, patient self-report can be used instead of complex testing in clinical practice. More research is needed to delineate the exact nature of these cognitive difficulties, including memory and attention problems. Lastly, uncovering causes of cognitive difficulties in fibromyalgia patients may lead to treatments that will improve cognitive function.

Disclosure of off-label usage: The author has determined that, to the best of her knowledge, no investigational information about pharmaceutical agents that is outside U.S. Food and Drug Administration–approved labeling has been presented in this article.

REFERENCES

1. Zachrisson O, Regland B, Jahreskog M, et al. A rating scale for fibromyalgia and chronic fatigue syndrome (the FibroFatigue scale). *J Psychosom Res* 2002;52:501–509
2. Bennett RM, Jones J, Turk DC, et al. An internet survey of 2596 people with fibromyalgia. *BMC Musculoskelet Disord* 2007;9:27
3. Katz RS, Heard AR, Mills M, et al. The prevalence and clinical impact of reported cognitive difficulties (fibrofog) in patients with rheumatic disease with and without fibromyalgia. *J Clin Rheumatol* 2004;10:53–58
4. Glass JM, Park DC, Minear M, et al. Memory beliefs and function in

- fibromyalgia patients. *J Psychosom Res* 2005;58:263–269
5. Dixon RA, Hultsch DR, Hertzog C. The Metamemory in Adulthood (MIA) Questionnaire. *Psychopharmacol Bull* 1988;25:671–688. Correction 1989;25:157
 6. Glass JM, Park DC. Cognitive dysfunction in fibromyalgia. *Curr Rheumatol Rep* 2001;3:123–127
 7. Dick B, Eccleston C, Crombez G. Attentional functioning in fibromyalgia, rheumatoid arthritis, and musculoskeletal pain patients. *Arthritis Rheum* 2002;47:639–644
 8. Salthouse TA, Babcock RL. Decomposing adult age differences in working memory. *Dev Psychol* 1991;27:763–777
 9. Park DC, Glass JM, Minear M, et al. Cognitive function in fibromyalgia patients. *Arthritis Rheum* 2001;44:2125–2133
 10. Gronwall DM. Paced auditory serial-addition task: a measure of recovery from concussion. *Percept Mot Skills* 1977;44:367–373
 11. Suhr JA. Neuropsychological impairment in fibromyalgia: relation to depression, fatigue, and pain. *J Psychosom Res* 2003;55:321–329
 12. Leavitt F, Katz RS. Distraction as a key determinant of impaired memory in patients with fibromyalgia. *J Rheumatol* 2006;33:127–132
 13. Stuss DT, Stethem LL, Pelchat G. Three tests of attention and rapid information processing: an extension. *Clin Neuropsychol* 1988;2:246–250
 14. Park DC, Smith AD, Lautenschlager G, et al. Mediators of long-term memory performance across the life span. *Psychol Aging* 1996;11:621–637
 15. Thurstone IL, Thurstone T. Examiner Manual for the SRA Primary Mental Abilities. Chicago, Ill: Scientific Resources Associates; 1949
 16. Robertson IH, Ward T, Ridgeway V, et al. The Test of Everyday Attention. Suffolk, Va: Thames Valley Test Company; 1994
 17. Glass JM, Park DC, Minear M. Memory performance with divided attention in fibromyalgia patients. *Arthritis Rheum* 2004;50(suppl 9):S489
 18. Glass JM, Park DC, Crofford LJ, et al. Fibromyalgia patients show reduced executive/cognitive control in a task-switching test. *Arthritis Rheum* 2006;54(suppl 9):S610
 19. Austin MP, Mitchell P, Goodwin GM. Cognitive deficits in depression: possible implications for functional neuropathology. *Br J Psychiatry* 2001;178:200–206
 20. Zakzanis KK, Leach L, Kaplan E. On the nature and pattern of neurocognitive function in major depressive disorder. *Neuropsychiatry Neuropsychol Behav Neurol* 1998;11:111–119
 21. Landrø NI, Stiles TC, Sletvold H. Memory functioning in patients with primary fibromyalgia and major depression and healthy controls. *J Psychosom Res* 1997;42:297–306
 22. Grace GM, Nielson WR, Hopkins M, et al. Concentration and memory deficits in patients with fibromyalgia syndrome. *J Clin Exp Neuropsychol* 1999;21:477–487
 23. Coté KA, Moldofsky H. Sleep, daytime symptoms, and cognitive performance in patients with fibromyalgia. *J Rheumatol* 1997;24:2014–2023
 24. Blatter K, Graw P, Münch M, et al. Gender and age differences in psychomotor vigilance performance under differential sleep pressure conditions. *Behav Brain Res* 2006;168:312–317
 25. Banks S, Dinges DF. Behavioral and physiological consequences of sleep restriction. *J Clin Sleep Med* 2007;3:519–528
 26. Staud R. Treatment of fibromyalgia and its symptoms. *Expert Opin Pharmacother* 2007;8:1629–1642
 27. Newcomer JW, Selke G, Melson AK, et al. Decreased memory performance in healthy humans induced by stress-level cortisol treatment. *Arch Gen Psychiatry* 1999;56:527–533
 28. Starkman MN, Giordani B, Berent S, et al. Elevated cortisol levels in Cushing's disease are associated with cognitive decrements. *Psychosom Med* 2001;63:985–993
 29. Sephton SE, Studts JL, Hoover K, et al. Biological and psychological factors associated with memory function in fibromyalgia syndrome. *Health Psychol* 2003;22:592–597
 30. Hart RP, Martelli MF, Zasler ND. Chronic pain and neuropsychological functioning. *Neuropsychol Rev* 2000;10:131–149
 31. Melzack R. The McGill Pain Questionnaire: major properties and scoring methods. *Pain* 1975;1:277–299
 32. Meenan RF, Mason JH, Anderson JJ, et al. AIMS2: the content and properties of a revised and expanded Arthritis Impact Measurement Scales Health Status Questionnaire. *Arthritis Rheum* 1992;35:1–10
 33. Glass J, Williams D, Gracely R, et al. Myofascial pain and fibromyalgia: relationship of self-reported pain, tender point count, and evoked pressure pain sensitivity to cognitive function in fibromyalgia. *J Pain* 2004;5(suppl 1):S38
 34. Staud R, Domingo M. Evidence for abnormal pain processing in fibromyalgia syndrome. *Pain Med* 2001;2:208–215
 35. Gracely RH, Petzke F, Wold JM, et al. Functional magnetic resonance imaging evidence of augmented pain processing in fibromyalgia. *Arthritis Rheum* 2002;46:1333–1343
 36. Glass JM. Cognitive dysfunction in fibromyalgia and chronic fatigue syndrome: new trends and future directions. *Curr Rheumatol Rep* 2006;8:425–429